

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

#### **About Google Book Search**

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/

# PHYSIOLOGICAL ECONOMY IN NUTRITION CHITTENDEN

M Ch

Boston Society of Natural History.

to juncted arti

Received me 9,191.6.

alle or with

H 5.m.1905.3

Harvard University

Library of
The Medical School
and

The School of Public Health



The Gift of

Boston Society of

Matural History

GLARKE CO

	,	

### PHYSIOLOGICAL ECONOMY

IN

### NUTRITION

WITH SPECIAL REFERENCE TO THE MINIMAL PROTEID REQUIREMENT OF THE HEALTHY MAN

### AN EXPERIMENTAL STUDY

BY

# RUSSELL H. CHITTENDEN,

PH.D., LL.D., Sc.D.

DIRECTOR OF THE SHEFFIELD SCIENTIFIC SCHOOL OF TALE UNIVERSITY AND PROFESSOR OF PHYSIOLOGICAL CHEMISTRY; MEMBER OF THE NATIONAL ACADEMY OF SCIENCES; PRESIDENT OF THE AMERICAN PHYSIOLOGICAL SOCIETY; MEMBER OF THE AMERICAN PHILOSOPHICAL SOCIETY, ETC.

> NEW YORK FREDERICK A. STOKES COMPANY 1905

# HARVARD UNIVERSITY SCHOOL OF MEDICINE AND PUBLIC HEALTH LIBRARY

Gift Boston Society of Natural History

/ 5. M. 1905.3

Copyright, 1904,
By Frederick A. Stokes Company

Published in November, 1904

## LIST OF ILLUSTRATIONS

						F	CI	ŧG.	PAGE
Group of soldiers at work in the Gymnasium	ı	•	•	•	•				136
Side view of Fritz									198
Back view of Fritz	,						•,		204
Front view of Coffman and Steltz	,								212
Back view of Coffman and Steltz								•	220
Side view of Zooman and Cohn									234
Back view of Zooman and Cohn									240
Side view of Loewenthal and Morris									<b>2</b> 58
Group of soldiers exercising in the Gymnasic	um	ı							262
Front view of Sliney								•	272
Soldiers exercising in the Gymnasium									<b>2</b> 84
Soldiers exercising in the Gymnasium .								•	296
Side view of Stapleton									<b>32</b> 8
Back view of Stapleton									866
Front view of W. L. Anderson and Bellis								•	440
Back view of W. L. Anderson and Bellis									442



### ACKNOWLEDGMENTS

# OF FINANCIAL AID IN MEETING THE EXPENSE OF THE EXPERIMENTS HEREIN DESCRIBED

THE writer has been most generously aided by substantial grants from the Bache Fund of the National Academy of Sciences, and from the Carnegie Institution of Washington; also by large donations from Mr. Horace Fletcher of Venice, and from Mr. John H. Patterson of Dayton, Ohio. In addition, the War Department of the United States met in large measure the expense of maintaining at New Haven the Detachment of Volunteers from the Hospital Corps of the United States Army, detailed here through the courtesy of Surgeon-General Robert Maitland O'Reilly.

#### ACKNOWLEDGMENTS

#### OF AID IN THE CONDUCT OF THE EXPERIMENTS

THE successful carrying out of the experiments in all their details, especially the chemical work, has been rendered possible by the active and continuous co-operation of the writer's colleague, Lafayette B. Mendel, Ph.D., Professor of Physiological Chemistry in the Sheffield Scientific School.

Efficient aid in the routine chemical and other work of the laboratory in connection with the experiments has been rendered by Frank P. Underhill, Ph.D., Arthur L. Dean, Ph.D., Harold C. Bradley, B.A., Robert B. Gibson, Ph.B., Oliver E. Closson, Ph.B., and Charles S. Leavenworth, Ph.B.

Dr. William G. Anderson, Director of the Yale Gymnasium, with the co-operation of his assistants, has rendered valuable aid in looking after the physical development of the men

under experiment, in arranging for frequent strength tests, as well as in prescribing the character and extent of their work in the Gymnasium. The greater portion of the training of the soldiers was under the personal supervision of William H. Callahan, M.D., Medical Assistant at the Gymnasium, while Messrs. William Chase, Anton Muller, John Stapleton, and H. R. Gladwin, Assistant Instructors in the Gymnasium, led the drills and looked after the actual muscular training of the men.

In the study of "Reaction Time" and other matters of psychological interest the work was under the direction of Charles H. Judd, Ph.D., in charge of the Yale Psychological Laboratory, aided by Warren M. Steele, B.A., and Cloyd N. McAllister, Ph.D.

In the morphological study of the blood, etc., Dr. Wallace DeWitt, Lieutenant in command of the Army detail, rendered valuable aid. Dr. DeWitt likewise co-operated in all possible ways during his stay in New Haven to maintain the integrity of the conditions necessarily imposed on the soldier detail in an experiment of this character.

Further, acknowledgments are due the several non-commissioned officers of the Hospital Corps for their intelligent co-operation and interest. Finally, to the men of the Hospital Corps who volunteered for the experiment, our thanks are due for their cheerful compliance with the many restrictions placed upon them during their six months' sojourn in New Haven, and for the manly way in which they conducted themselves under conditions not always agreeable.

To the students of the University who volunteered as subjects of experiment our acknowledgments are due for their intelligent co-operation, keen interest, and hearty compliance with the conditions imposed.

### PREFACE

THERE is no subject of greater physiological importance, or of greater moment for the welfare of the human race, than the subject of nutrition. How best to maintain the body in a condition of health and strength, how to establish the highest degree of efficiency, both physical and mental, with the least expenditure of energy, are questions in nutrition that every enlightened person should know something of, and yet even the expert physiologist to-day is in an uncertain frame of mind as to what constitutes a proper dietary for different conditions of life and different degrees of activity. We hear on all sides widely divergent views regarding the needs of the body, as to the extent and character of the food requirements, contradictory statements as to the relative merits of animal and vegetable foods; indeed, there is great lack of agreement regarding many of the fundamental questions that constantly arise in any consideration of the nutrition of the human body. Especially is this true regarding the so-called dietary standards, or the food requirements of the healthy adult. Certain general standards have been more or less widely adopted, but a careful scrutiny of the conditions under which the data were collected leads to the conclusion that the standards in question have a very uncertain value, especially as we see many instances of people living, apparently in good physical condition, under a régime not at all in harmony with the existing standards.

Especially do we need more definite knowledge of the true physiological necessities of the body for proteid or albuminous foods, *i.e.*, those forms of foods that we are accustomed to speak of as the essential foods, since they are absolutely requisite for life. If our ideas regarding the daily quantities of these foods necessary for the maintenance of health and

strength are exaggerated, then a possible physiological economy is open to us, with the added possibility that health and vigor may be directly or indirectly increased. Further, if through years and generations of habit we have become addicted to the use of undue quantities of proteid foods, quantities way beyond the physiological requirements of the body, then we have to consider the possibility that this excess of daily food may be more or less responsible for many diseased conditions, which might be obviated by more careful observance of the true physiological needs of the body.

First, however, we must have more definite information as to what the real necessities of the body for proteid food are, and this information can be obtained only by careful scientific experimentation under varying conditions. This has been the object of the present study, and the results obtained are now placed before the public with the hope that they will prove not only of scientific interest and value, but that they will also serve to arouse an interest in the minds of thoughtful people in a subject which is surely of primary importance for the welfare of mankind. That the physical condition of the body exercises an all-powerful influence upon the mental state, and that a man's moral nature even is influenced by his bodily condition are equally certain; hence, the subject of nutrition, when once it is fully understood and its precepts obeyed, bids fair to exert a beneficial influence not only upon · bodily conditions, but likewise upon the welfare of mankind in many other directions.

In presenting the results of the experiments, herein described, the writer has refrained from entering into lengthy discussions, preferring to allow the results mainly to speak for themselves. They are certainly sufficiently convincing and need no superabundance of words to give them value; indeed, such merit as the book possesses is to be found in the large number of consecutive results, which admit of no contradiction and need no argument to enhance their value. The results presented are scientific facts, and the conclusions they justify are self-evident.

# CONTENTS

	Page
ACKNOWLEDGMENTS	. v
Preface	vii
Introductory	. 1
INTRODUCTORI	-
<b>I.</b>	
Experiments with Professional Men.	
Chittenden: Daily Record of Nitrogen Excretion, etc	. 24
First Nitrogen Balance, with comparison of income and output	•
amount and character of the daily food	
Second Nitrogen Balance, with composition of daily food, etc.	43
Mendel: Daily Record of Nitrogen Excretion, etc	53
First Nitrogen Balance, with comparison of income and output	)
amount and character of the daily food	. 60
Second Nitrogen Balance, with composition of daily food, etc.	67
Underhill: Daily Record of Nitrogen Excretion, etc	79
First Nitrogen Balance, with comparison of income and output	,
composition of the daily food, etc	87
Second Nitrogen Balance, with composition of daily food, etc.	93
Dean: Daily Record of Nitrogen Excretion, etc	98
Nitrogen Balance, with comparison of income and output, amoun	j
and character of the daily food	103
Beers: Daily Record of Nitrogen Excretion, etc	111
First Nitrogen Balance, with comparison of income and output	
amount and character of the daily food	114
Second Nitrogen Balance, with composition of daily food, etc.	121
Summary of Results; True Proteid Requirements	127
TY	
II.	
EXPERIMENTS WITH VOLUNTEERS FROM THE HOSPITAL COR	P8
OF THE UNITED STATES ARMY.	
Description of the Men	134
Daily Routine of Work	135

	Page
Daily Record of Nitrogen Excretion, etc., for each of the thirteen	
men under experiment	139
Average Daily Output of Nitrogen	199
Nitrogen Metabolized per kilo of Body-Weight	201
Changes in Body-Weight during the Experiment	202
First Nitrogen Balance, with comparison of income and output,	
amount and character of the daily food	203
Second Nitrogen Balance, with composition of daily food, etc	223
Third Nitrogen Balance, with composition of daily food, etc	242
Summary regarding Nitrogen Requirement	254
Physical Training of the Men-Report by Dr. Anderson of the	
Yale Gymnasium	255
Body Measurements	261
Strength or Dynamometer Tests	262
Comparison of the Total Strength of the Men at the beginning and	
end of the Experiment	274
Reaction Time Experiments - Report by Dr. Judd of the Yale	
Psychological Laboratory	276
Character and Composition of the Blood	283
General Conclusions	285
General Conclusions	288
•	
III.	
111.	
Experiments with University Students, trained in	
ATHLETICS.	
Consumption of Proteid Food by Athletes	<b>327</b>
Description of the Men	329
Daily Record of Nitrogen Excretion, etc., for each of the eight men	
under Experiment	<b>332</b>
Average Daily Excretion of Metabolized Nitrogen	364
Metabolized Nitrogen per kilo of Body-Weight	365
Daily Diet Prescribed	366
Nitrogen Balance, with comparison of income and output, and	
amount and character of the daily food, etc	375
The Physical Condition of the Men	434
Strength or Dynamometer Tests	436
Report by Dr. Anderson of the Yale Gymnasium	439
Reaction Time — Report by Dr. Judd of the Yale Psychological	
Laboratory	442
General Summary; True Physiological Requirements for Proteid	
Food	454

### IV.

THE SYSTEMIC VALUE OF PHYSIOLOGICAL ECONOMY IN	
te Products of Proteid Metabolism may be Dangerous to Health 456 n and Significance of Uric Acid	
	_
•	-
	<b>4</b> 58
Modification of Uric Acid Excretion by diminishing the amount of	
Proteid Food	463
Tables showing Excretion of Uric Acid by the three groups of men	
under observation; Uric Acid per kilo of Body-Weight, etc	467
V.	
ECONOMIC AND SOCIOLOGICAL IMPORTANCE OF THE RESULTS .	471
VI.	
General Conclusions	474
VII.	
DESCRIPTION OF ILLUSTRATIONS	477

1



# PHYSIOLOGICAL ECONOMY IN NUTRITION

### INTRODUCTORY

AS the result of many years of observation and experiment certain general conclusions have been arrived at regarding the requisite amounts of food necessary for the maintenance of health and strength. Certain dietary standards have been set up which have found more or less general acceptance in most parts of the civilized world; standards which have been reinforced and added to by man's aptitude for self-indulgence. Carl Voit, of Munich, whose long and successful life as a student of Nutrition renders his conclusions of great value, considers that an adult man of average body-weight (70-75 kilos) doing moderate muscular work requires daily 118 grams of proteid or albuminous food, of which 105 grams should be absorbable, 56 grams of fat, and 500 grams of carbohydrate, with a total fuel value of over 3000 large calories, in order to maintain the body in equilibrium. The Voit standard or daily diet is accepted more or less generally as representing the needs of the body under normal conditions of life, and

Note. — For the benefit of lay readers, metabolism, a word frequently made use of, may be defined as a term applied to the collective chemical changes taking place in living matter. When these metabolic changes are constructive, as in the building up of tissue protoplasm from the absorbed food material, they are termed anabolic; when they are destructive, as in the breaking down of living matter or in the decomposition of the materials stored up in the tissues and organs, they are termed katabolic. Proteid metabolism, or more exactly proteid katabolism, therefore, means the destructive decomposition of proteid or albuminous matter in the living body and is practically synonymous with nitrogenous metabolism, since the entire nitrogen income is mainly supplied by the proteids or albuminous matters of the food. The chief carbon income, on the other hand, is supplied by fats and carbohydrates, such as starches and sugars.

the conclusions arrived at by other investigators along these same lines have been more or less in accord with Voit's figures. In confirmation of this statement the following data may be quoted:

AVERAGE I	IETS.
-----------	-------

	Moleschott.	Ranke.	Forster.	Hultgrent and Landergren.	Atwater.	Studemund.‡	Schmidt.§
	grams	grams	grams	grams	grams	grams	grams
Proteid	130	100	131	134	125	114	105
Carbohydrates .	550	240	494	523	400	551	541
Fats	40	100	68	79	125	54	63
Fuel value (calories)	8160	2824	3195	3436	3315	3229	3235

In many of these diets it is to be noted that the proteid requirement is placed at even a higher figure than Voit's standard. Similarly, Erisman, studying the diets of Russian workmen having a free choice of food and doing moderately hard work, found the daily diet to be composed of 181.8 grams of proteid, 79.7 grams of fat, and 583.8 grams of carbohydrate, with a total fuel value of 3675 large calories. Further, Hultgren and Landergren ¶ found that Swedish laborers doing hard work had as their daily diet 189 grams of proteid, 110 grams of fat, and 714 grams of carbohydrate, with a total fuel value of 4726 large calories. Voit found that German soldiers on active service consumed daily 145 grams of proteid, 100 grams of fat, and 500 grams of carbohydrate, with a total fuel value

<sup>\*</sup> Die Ernährung des Menschen. Munich, 1876.

<sup>†</sup> Untersuchungen ü. d. Ernähr. schwedischer Arbeiter. Stockholm, 1891.

<sup>†</sup> Ein Beitrag zur Lehre vom Eiweissbedarf des gesunden Menschen. Pflüger's Archiv f. d. gesammte Physiologie. Band 48, p. 578.

<sup>§</sup> D. Militärkrztl. Zeitschr., 1901, p. 622. Quoted by Hirschfeld, Archiv. f. Physiologie, 1903, p. 881.

<sup>||</sup> Large calories, or kilogram-degree units of heat.

<sup>¶</sup> Jahresbericht für Thierchemie, 1891, p. 369. The figures given in the above table represent the diet for men doing a moderate amount of work.

of 3574 large calories. Lichtenfelt,\* studying the nutrition of Italians, states that an Italian laborer doing a moderate amount of work requires 110.5 grams of proteid and a total fuel value for the daily food of 2698 calories, while at hard labor he needs 146 grams of proteid daily, with carbohydrates and fat sufficient to give 3088 large calories. In our own country Atwater,† who has made many valuable observations upon the dietetic habits of different classes of people and under different conditions of life, has stated that a somewhat more liberal allowance of proteid would seem desirable, say 125 grams, with a total fuel value of 3500 large calories for a man doing severe muscular labor.

In what is perhaps the latest book on alimentation, Armand Gautier,‡ writing of the French people, states that the ordinary man in that climate needs daily 110 grams of albuminous food, 68 grams of fat, and about 428 grams of amylaceous or saccharine food. It is possible, however, says Gautier, that the quantity of albuminous food can be reduced, if necessary, to 78 grams per day in case a man is not doing work and takes in addition at least 50 grams of fat and 485 grams of carbohydrate food. Where, however, an individual works eight to ten hours a day, the ration, says Gautier, must be increased to at least 135 grams of albuminous food, with 85 to 100 grams of fat, and with from 500 to 900 grams of starchy food.

While these figures may be taken as showing quite conclusively the dietetic standards adopted by mankind, there is no evidence whatever that they represent the real needs or requirements of the body. We may even question whether simple observation of the kinds and amounts of food consumed by different classes of people under different conditions of life have any very important bearing upon this question. They

<sup>\*</sup> Ueber die Ernährung der Italiener. Pflüger's Archiv. f. d. gesammte Physiologie, Band 99, p. 1 (1903).

<sup>†</sup> Bulletin No. 46. United States Department of Agriculture, p. 63.

<sup>‡</sup> L'Alimentation et les Régimes chez l'Homme sain et chez les Malades, Paris, 1904.

### 4 PHYSIOLOGICAL ECONOMY IN NUTRITION

throw light upon dietetic habits, it is true, but such observations give no information as to how far the diets in question serve the real needs of the body. We may find, for example, that under certain given conditions of diet the people in question have the appearance of being well nourished, and that they do their work with apparent ease and comfort; but might not these same results follow with smaller amounts of food? If so, there must of necessity be a certain amount of physiological economy under the more restricted diet, and a consequent ultimate gain to the body through diminished wear and tear of the bodily machinery.

Indeed, experimental work and observations scattered through the last few years have suggested the possibility of much lower standards of diet sufficing to meet the real physiological needs of the body. Thus, Hirschfeld,\* in 1887, found in experimenting on himself (24 years of age and weighing 73 kilos) that it was possible to maintain nitrogen equilibrium on a diet containing only 5 to 7.5 grams of nitrogen per day, or 35 to 45 grams of proteid, for a period of ten to fifteen days. The amount of non-nitrogenous food consumed, however, was fairly large, especially the amount of butter, - frequently 100 grams a day - the average fuel value ranging from 3750 to 3916 large calories daily. In 1888 Hirschfeld,† again experimenting on himself, maintained nitrogen equilibrium for several days on 7.5 grams of nitrogen per day, with fats and carbohydrate sufficient to yield a total fuel value of 3462 large calories as the daily average. The chief criticism of Hirschfeld's experiments is that he failed to obtain in all cases definite analytical data of the food-stuffs employed and failed to determine the nitrogen of the fæces. Still his results are of value as indicating the possibility of maintaining nitrogenous equilibrium for a brief time at least on a low proteid intake.

<sup>\*</sup> Untersuchungen über den Eiweissbedarf des Menschen. Pflüger's Archiv f. die gesammte Physiologie. Band 41, p. 583.

<sup>†</sup> Beiträge zur Ernährungslehre des Menschen. Virchow's Archiv für pathol. Anat. u. Physiol. Band 114, p. 311.

Kumagawa, \* studying especially the diet of the Japanese and experimenting on himself (27 years old and weighing 48 kilos), found with a purely vegetable diet, containing per day 54.7 grams of proteid, 2.5 grams of fat, and 569.8 grams of carbohydrate, that he showed for a period of nine days a plus balance of nitrogen, indicating that his body was laying on about 4 grams of proteid per day. The nitrogen excreted per urine and fæces amounted to 8.09 grams per day, while the nitrogen in the daily food amounted to 8.75 grams. It is interesting to observe in these experiments, as indicating the degree of absorption of the vegetable food (composed in large measure of rice) that the daily average of nitrogen in the urine amounted to 6.069 grams and in the fæces 2.029 grams. In other words, of the 54.7 grams of nitrogen-containing food only 37.8 grams were absorbed, 12.69 grams passing out with the fæces. The total fuel value of the absorbed food per day was 2478 large calories. Similarly, Hirschfeld † has called attention to the fact that with many vegetable foods especially, not more than 75 per cent of the ingested proteid can be digested and absorbed, thus emphasizing the necessity of paying heed to the character of the proteid food in considering the nutritive value of a given diet.

In some experiments reported by C. Voit ‡ in 1889, on the diet of vegetarians, E. Voit and Constantinidi found that nitrogenous equilibrium was established in one man with about 8 grams of nitrogen, corresponding to 48.5 grams of proteid as the daily diet, with large amounts of starchy foods and some fat. Similarly, Nakahama § in the same year, studying the diet (mostly vegetable) and nutritive condition of thirteen German laborers in Leipzig, found that their daily

<sup>\*</sup> Vergleichende Untersuchungen über die Ernährung mit gemischter und rein vegetablischer Kost mit Berücksichtigung des Eiweissbedarfes. Virchow's Archiv für pathol. Anat. u. Physiol. Band 116, p. 370.

<sup>†</sup> Die Ernährung der Soldaten vom physiologischen und volkswirthschaftlichen Standpunkt. Achiv f. Physiologie 1903, p. 380.

t Ueber die Kost eines Vegetariers. Zeitschr. f. Biologie. Band 25, p. 281.

<sup>§</sup> Ueber den Eiweissbedarf des Erwachsenen mit Berücksichtigung der Beköstigung in Japan. Archiv f. Hygiene. Band 8, p. 78.

food contained on an average 85 grams of proteid, but Carl Voit criticising these results states that the men were of comparatively light body-weight—about 60 kilos—and not well nourished.

Kellner and Mori,\* studying the nutrition of a Japanese (weighing 52 kilos and 23 years of age) state that on a purely vegetable diet containing 11.34 grams of nitrogen, of which only 8.58 grams were digested, there was a distinct loss of body-weight, with a daily loss to the body of 1.16 grams of nitrogen. On a mixed diet, however, containing fish, it was possible to establish nitrogenous equilibrium with a daily diet containing 17.48 grams of nitrogen, of which 15.27 grams were digested and utilized. Similarly, Caspari, † 29 years old and weighing 66.2 kilos, found that while he could maintain his body in nitrogenous equilibrium on 13.26 grams of nitrogen per day, he could not accomplish it on 10.1 grams of nitrogen, though his daily food contained 3200 large calories.

Other investigators, however, have found no great difficulty in establishing nitrogenous equilibrium in man with much lower quantities of proteid food. Thus, Klemperer ‡ found in the case of two young men of 64 and 65.5 kilos body-weight respectively, in an experiment lasting eight days, that nitrogenous equilibrium was established on 4.38 and 3.58 grams of nitrogen per day, but with a daily diet containing in addition to the small amount of proteid 264 grams of fat, 470.4 grams of carbohydrate, and 172 grams of alcohol, with a total fuel value of 5020 large calories.

Peschel, \$ too, has reported experimental results showing that he was able to establish nitrogenous equilibrium for a

<sup>\*</sup> Untersuchungen über die Ernährung der Japaner. Zeitschr. f. Biologie. Band 25, p. 102.

<sup>†</sup> Ein Beitrag zur Frage der Ernährung bei verringerter Eiweisszufuhr. Archiv f. Physiologie, Jahrgang 1901, p. 323.

<sup>†</sup> Untersuchungen über Stoffwechsel und Ernährung in Krankheiten. Zeitschr. f. klin, Medizin. Band 16, p. 550.

<sup>§</sup> Untersuchungen über den Eiweissbedarf des gesunden Menschen. Berlin, 1891.

brief period with 7 grams of nitrogen daily, 5.31 grams appearing in the urine and 1.58 grams in the fæces.

Caspari and Glaessner,\* in a five-days' experiment with two vegetarians, found that the wife consumed daily, on an average, 5.33 grams of nitrogen, with fats and carbohydrates to equal 2715 calories, while the man took in 7.82 grams of nitrogen and 4559 calories. Both persons laid on nitrogen in spite of the low intake of proteid food.

Siven's † experiments, however, are perhaps worthy of more careful consideration. Of 60 kilos body-weight and 301 years of age, his experiments conducted on himself extended through thirty-two days with establishment of nitrogenous equilibrium on 6.26 grams of nitrogen. Moreover, in another experiment he was in nitrogen equilibrium for a day or two at least on 4.5 grams of nitrogen. In Siven's experiment, the most noticeable feature is the added fact that the total intake of food per day was comparatively low, with a fuel value of only 2444 large calories. In this connection we may call attention to the recent experiments of Landergren, t who found with four individuals fed on a daily diet containing only 2.1 to 2.4 grams of nitrogen, but with a large amount of carbohydrate, some fat and alcohol, that on the fourth day of this "specific nitrogen hunger" only 3 to 4 grams of nitrogen were metabolized and appeared in the urine. In other words, a healthy adult man having a sufficient intake of nonnitrogenous food seemingly need not metabolize more proteid than suffices to yield 3 to 4 grams of nitrogen per day.

Such data as these, of which many more might be quoted, surely warrant the question, how far are we justified in assuming the necessity for the rich proteid diet called for by the Voit standard? Voit, however, with many other physiol-

<sup>\*</sup> Ein Stoffwechselversuch an Vegetarianern. Biochemisches Centralblatt Band 2, p. 144 (1903).

<sup>†</sup> Ueber das Stickstoffgleichgewicht beim erwachsenen Menschen. Skandinavisches Archiv f. Physiol. Band 10, p. 91.

<sup>†</sup> Untersuchungen über die Eiweissumsetzung des Menschen. Skandinavisches Archiv f. Physiol. Band 14, p. 121 (1903).

j

ogists would apparently object to any diminution of the daily 118 grams of proteid for the moderate worker, on the ground that an abundance of proteid in the food is a necessity for the maintenance of physical vigor and muscular activity. This view is certainly reinforced by the customs and habits of mankind; but we may well query whether our dietetic habits will bear criticism, and in the light of modern scientific inquiry we may even express doubt as to whether a rich proteid diet adds anything to our muscular energy or bodily strength.

How far can our natural instinct be trusted in the choice of diet? We are all creatures of habit, and our palates are pleasantly excited by the rich animal foods with their high content of proteid, and we may well question whether our dietetic habits are not based more upon the dictates of our palates than upon scientific reasoning or true physiological needs. There is a prevalent opinion that to be well nourished the body must have a large excess of fat deposited throughout the tissues, and that all bodily ills and weaknesses are to be · met and combated by increased intake of food. constant temptation to increase the daily ration, and there is almost universal belief in the efficacy of a rich and abundant diet to strengthen the body and to increase bodily and mental Is there any justification for these beliefs? apparently, other than that which comes from the customs of generations of high living.

It is self-evident that the smallest amount of food that will serve to keep the body in a state of high efficiency is physiologically the most economical, and hence the best adapted for the needs of the organism. Any excess over and above what is really needed is not only uneconomical, but may be directly injurious. This is especially true of the proteid or albuminous foods. It is, however, quite proper to question whether a brief experiment of a few days in which nitrogenous equilibrium is perhaps established at the low level of 4 to 5 grams of nitrogen, the equivalent of 25 to 35 grams of proteid, is to be accepted as fixing the daily requirements of the healthy man, offsetting the customs or habits of a lifetime. Voit himself,

however, has clearly emphasized the general principle that the smallest amount of proteid, with non-nitrogenous food added, that will suffice to keep the body in a state of continual vigor is the ideal diet. Proteid decomposition products are a constant menace to the well-being of the body; any quantity of proteid or albuminous food beyond the real requirements of the body may prove distinctly injurious. We see the evil effects of uric acid in gout, but there are many other nitrogenous waste products of proteid katabolism, which with excess of proteid food are liable to be unduly conspicuous in the fluids and tissues of the body, and may do more or less damage prior to their excretion through the kidneys. Further, it requires no imagination to understand the constant strain upon the liver and kidneys, to say nothing of possible influence upon the central and peripheral parts of the nervous system, by these nitrogenous waste products which the body ordinarily gets rid of as speedily as possible. They are an ever present evil, but why increase them unnecessarily? This question brings us back to the starting-point. What is the minimal proteid requirement for the healthy man, or rather, how far can we safely and advantageously diminish our proteid intake below the commonly accepted standards?

The question of safety is a pertinent one. Thus, Munk \* some years ago (1893) sounded a warning on this point which was later confirmed by Rosenheim.† Both of these observers reported that in dogs fed for some time on a low proteid diet, but with an abundance of carbohydrate and fat, there was after some weeks (6-8) a loss of the power of absorption from the alimentary tract, dependent not alone upon a changed condition of the epithelial cells of the intestine, but also upon a diminished secretion of the digestive juices, loss of bodyweight, strength, and vigor, followed speedily by death. If

<sup>\*</sup> Ueber die Folgen einer ausreichenden, aber eiweissarmen Nahrung. Ein Beitrag zur Lehre vom Eiweissbedarf. Virchow's Archiv für pathol. Anat. u. Physiol. Band 132, p. 91.

<sup>†</sup> Weitere Untersuchungen über die Schädlichkeit eiweissarmer Nahrung. Pflüger's Archiv f. die gesammte Physiol. Band 54, p. 61.

these results were really due to the low proteid diet, they suggest a grave danger which must not be lightly passed by. Jägerroos \* has likewise observed, experimenting on dogs, that there was, after some months, a striking disturbance of the intestines on a low proteid intake, which, however, was eventually traced to a distinct infection, and probably in no manner connected with the diminished amount of proteid in In these various experiments on dogs carried out by Munk, Rosenheim, and by Jägerroos, there was of necessity great monotony in the diet, and in Munk's experiments no fresh meat at all was fed, but simply dried food. In other words, if the diet was in any sense responsible for the poor health of the animals, it is fully as plausible to attribute the results to the abnormal conditions under which the animals were kept as to any specific effect due to the low proteid intake. It is very essential that the food of dogs, as of men, shall fulfil all ordinary hygienic conditions. It must be not only of sufficient quantity for the true needs of the body, but it should also have the necessary variety with reasonable degree of digestibility, and proper volume or bulk. these qualities are lacking, it is not strange if deviations from the normal gradually develop. That the low intake of proteid food could be responsible for the condition existing in Munk's and Rosenheim's experiments is not plausible; a view which is strongly reinforced by many observations, notably those of Albu t on a woman thirty-seven years old and weighing 37.5 kilos, who had followed a vegetarian diet for six years, and who while under Albu's care for two years consumed only 34 grams of proteid per day, the total fuel value of the food being only 1400 calories per day. This woman was in nitrogenous equilibrium on 5.4 grams of nitrogen, and on this diet had freed herself from the illness to which she had long been subject.

<sup>\*</sup> Ueber die Folgen einer ausreichenden, aber eiweissarmen Nahrung. Skandinavisches Archiv f. Physiol. Band 13, p. 375.

<sup>†</sup> Zur Bewertung der vegetarischen Diät. Berliner klin. Wochenschr. 1901. p. 647 and 670. See also, Albu, die vegetarische Diät. Leipzig, 1902. p. 65.

Voit's \* vegetarian is described by Voit himself as a man twenty-eight years old, weighing 57 kilos, well nourished, with well developed muscles, etc. He had lived on a purely vegetable diet for three years, and was found to be in nitrogenous equilibrium on 8.2 grams of nitrogen. No mention is made of any disagreeable effects connected with this low proteid ration, although persisted in for several years. Jaffa's t experiments and observations on the fruitarians and nutarians of California "showed in every case (two women and three children) that though the diet had a low protein and energy value, the subjects were apparently in excellent health and had been so during the five to eight years they had been living in this manner." In comparing the income and outgo of nitrogen on a diet composed mainly of nuts and fruits, it was observed in two subjects that 8 grams of nitrogen were sufficient to bring about nitrogen equilibrium, while with two other subjects on a like diet the nitrogen required daily for equilibrium was about 10 grams. The diet used in these experiments, however, was of necessity more or less restricted in variety, and was without doubt somewhat monotonous. Jaffa appears to agree with Caspari that the minimum amount of proteid required daily varies with the individual, and may even vary with the same individual at different times. Further, Jaffa, in harmony with Siven, believes that after the body has suffered a loss of nitrogen, there is at once an effort to attain nitrogenous equilibrium, and that any gain of nitrogenous body material is a comparatively slow process. If this is true, it is obvious that the living substance of the tissue protoplasm must be slowly formed from the proteid of the diet. This, says Jaffa, should serve as a warning to anyone contemplating any appreciable decrease in the proteid of the daily diet.

Another statement made by Jaffa may be quoted in this

<sup>\*</sup> See Zeitschr. f. Biologie. Band 25, p. 255.

<sup>†</sup> Further Investigations among Fruitarians at the California Agricultural Experiment Station. 1901–1902. U. S. Department of Agriculture. Bulletin No. 182.

connection, since it illustrates the attitude taken by many physiologists on this question. "Even if it could be proved," says Jaffa, "by a large number of experiments that nitrogen equilibrium can be maintained on a small amount of protein, it would still be a great question whether or not it would be wise to do so. There must certainly be a constant effort on the part of the human organism to attain this condition, and with a low protein supply it might be forced to do so under conditions of strain. In such a case the bad results might be slow in manifesting themselves, but might also be serious and lasting. It has also been suggested that when living at a fairly high protein level the body is more resistant to disease and other strains than when the protein level is low." While these suggestions demand careful consideration, it is equally evident that there is another side to the question, viz., the possible danger to the body from the physiological action of the larger amounts of nitrogenous waste products which result from an excess of proteid food, and which float about through the system prior to their excretion. In addition, we must not overlook the great loss of energy to the body in handling and getting rid of the surplus of unnecessary food of whatever kind introduced into the alimentary tract, to say nothing of the danger of intestinal putrefaction and toxemia when from any cause the system loses its ability to digest and absorb the excess of food consumed. Further, the possible strain on the kidneys and other organs must not be overlooked. Hence we may well query on which side lies the greater danger. unprejudiced observer, one not wedded to old-time tradition, it would seem as if great effort was being made to sustain the claims of a high-proteid intake. It is surely well to be careful, but it is certainly not necessary to magnify imaginary dangers to the extent of suppressing all efforts toward the establishment of possible physiological economy.

In a paper read before the Physiological Section of the British Medical Association in 1901 by Dr. van Someren, claim is made of the existence of a reflex of deglutition, the proper working of which protects from the results of mal-

nutrition by preventing the intake of any excess of food. Thorough mastication and insalivation aid in the more complete utilization of the food and render possible great economy, so that body-weight and nitrogen equilibrium are both maintained on an exceptionally small amount of food. This principle had been worked out by Mr. Horace Fletcher on himself in an attempt to restore his health to a normal condition, with such beneficial results that he was speedily restored to a state of exceptional vigor and well-being. Deliberation in eating, necessitated by the habit of thorough insalivation, it is claimed results in the occurrence of satiety on the ingestion of comparatively small amounts of food, and hence all excess of food is avoided.

In the autumn of 1901, Mr. Fletcher and Dr. van Someren visited the physiological laboratories of Cambridge University, and as stated by Sir Michael Foster\* the matter was more closely inquired into with the assistance of physiological experts. Observations were carried out on various individuals, and as stated by Professor Foster "the adoption of the habit of thorough insalivation of the food was found in a consensus of opinion to have an immediate and very striking effect upon appetite, making this more discriminating, and leading to the choice of a simple dietary, and in particular reducing the craving for flesh food. The appetite, too, is beyond all question fully satisfied with a dietary considerably less in amount than with ordinary habits is demanded." . . . "In two individuals who pushed the method to its limits it was found that complete bodily efficiency was maintained for some weeks upon a dietary which had a total energy value of less than one-half of that usually taken, and comprised little more than one-third of the proteid consumed by the average man." Finally, says Foster, "it may be doubted if continued efficiency could be maintained with such low values as these, and very prolonged observations would be necessary to establish the facts. But all subjects of the experiments who applied the principles

<sup>\*</sup> See Horace Fletcher, The A-B-Z of our own Nutrition. (1903.) New York. p. 48.

intelligently agreed in finding a very marked reduction in their needs, and experienced an increase in their sense of wellbeing and an increase in their working powers."

In the autumn of 1902 and in the early part of 1903, Mr. Fletcher spent several months with the writer, thereby giving an opportunity for studying his habits of life. For a period of thirteen days in January he was under constant observation in the writer's laboratory, when it was found that the average daily amount of proteid metabolised was 41.25 grams, his body-weight (75 kilos) remaining practically constant. Later, a more thorough series of observations was made, involving a careful analysis of the daily diet, together with analysis of the For a period of six days the daily diet averaged 44.9 grams of proteid, 38.0 grams of fat, and 253 grams of carbohydrate, the total fuel value amounting to only 1606 large calories per day. The daily intake of nitrogen averaged 7.19 grams, while the daily output through the urine was 6.30 grams and in the fæces 0.6 gram; i.e., a daily intake of 7.19 grams of nitrogen, with a total output of 6.90 grams, showing a daily gain to the body of 0.29 gram of nitrogen. and this on a diet containing less than half the proteid required by the Voit standard and having only half the fuel value of the Voit diet. Further, it was found by careful and thorough tests made at the Yale Gymnasium that Mr. Fletcher, in spite of this comparatively low ration was in prime physical condition. In the words of Dr. Anderson, the Director of the Gymnasium, "the case is unusual, and I am surprised that Mr. Fletcher can do the work of trained athletes and not give marked evidences of over-exertion. . . . Mr. Fletcher performs this work with greater ease and with fewer noticeable bad results than any man of his age and condition I have ever worked with."\* It is not our purpose here to discuss how far these results are due to insalivation, or the more thorough mastication of food. The main point for us is that we have here a striking illustration of the establishment of nitrogen

<sup>\*</sup> For a fuller account of this study, see Chittenden, Physiological Economy in Nutrition. Popular Science Monthly, June, 1903.

equilibrium on a low proteid diet and great physiological economy as shown by the low fuel value of the food consumed, coupled with remarkable physical strength and endurance.

With data such as these before us we see the possible importance of a fuller and more exact knowledge of true dietary We find here questions suggested, the answers to standards. which are of primary importance in our understanding of the nutritive processes of the body; greater ease in the maintenance of health, increased power of resistance to disease germs, duration of life increased beyond the present average, greater physiological economy and greater efficiency, increased mental and physical vigor with less expenditure of energy on the part of the body. All these questions rise before us in connection with the possibility of maintaining equilibrium on a lowered intake of food, especially nitrogenous equilibrium, with a diminished consumption of proteid or albuminous food. Is it not possible that the accepted dietary standards are altogether too high?

It is of course understood that there can be no fixed dietary standard suitable for all people, ages, and conditions of life. Dietary standards at the best are merely an approximate indication of the amounts of food needed by the body, but these needs are obviously changeable, varying with the degree of activity of the body, especially the amount of physical work performed, to say nothing of differences in body-weight, sex, etc. Further, it is doubtless true that there is what may be called a specific coefficient of nutrition characteristic of the individual, a kind of personal idiosyncrasy which exercises in some degree a modifying influence upon the character and extent of the changes going on in the body. Still, with due recognition of the general influence exerted by these various factors the main question remains, viz., how far the usually accepted standards of diet are correct; or, in other words, is there any real scientific ground for the assumption that the average individual doing an average amount of work requires any such quantity of proteid, or of total nutrients, as the ordinary dietetic standards call for? Cannot all the real physiological needs of the body be met by a greatly reduced proteid intake, with establishment of continued nitrogenous equilibrium on a far smaller amount of proteid food than the ordinary dietary standards call for, and with actual gain to the body?

Just here we may emphasize why prominence is given to the establishment of nitrogenous equilibrium, and why the proteid intake assumes a greater importance than the daily amounts of fat and carbohydrate consumed. Fats and carbohydrates when oxidized in the body are ultimately burned to simple gaseous products, viz., carbonic acid and water. Hence, these waste products are easily and quickly eliminated and cannot exercise much deleterious influence even when formed in excess. To be sure, there is waste of energy in digesting, absorbing, and oxidizing the fats and carbohydrates when they are taken in excessive amounts. troduced into the alimentary canal they must be digested, otherwise they will clog the intestine or undergo fermentation, and so cause trouble. Further, when absorbed they may be transformed into fat and deposited in the various tissues and organs of the body; a process desirable up to a certain point. but undesirable when such accumulation renders the body gross and unwieldy. With proteid foods, on the other hand, the story is quite different. These substances, when oxidized, yield a row of crystalline nitrogenous products which ultimately pass out of the body through the kidneys. Prior to their excretion, however, these products - frequently spoken of as toxins — float about through the body and may exercise more or less of a deleterious influence upon the system, or, being temporarily deposited, may exert some specific or local influence that calls for their speedy removal. Hence, the importance of restricting the production of these bodies to the minimal amount, owing to their possible physiological effect and the part they are liable to play in the causation of many Further, the elimination of excessive diseased conditions. amounts of these crystalline nitrogenous bodies through the kidneys places upon these organs an unnecessary burden which

is liable to endanger their integrity and possibly result in serious injury, to say nothing of an early impairment of function.

The present experiments were undertaken to throw light upon this broad question of a possible physiological economy in nutrition, and with special reference to the minimal proteid requirement of the healthy man under ordinary conditions of life. The writer as a student of physiology has always maintained that man is disposed to eat far more than the needs of the body require, but his active interest in this problem was aroused especially by his observations of Mr. Fletcher and the marked physiological economy the latter was able to practice, not only without detriment, but apparently with great gain to the body as regards strength, vigor, and endurance, coupled with an apparent resistance to disease. While Mr. Fletcher and Dr. Van Someren would doubtless emphasize the importance of insalivation as a means of controlling the appetite and thereby regulating the consumption of food in harmony with the real needs of the body, it is of primary importance for the physiologist and for mankind to know definitely how far it is possible to reduce the intake of food with perfect safety and without loss of that strength, mental and physical, vigor, and endurance which are characteristic of good health. Further, it is equally plain that if there is possible gain to the body from a practice of physiological economy in diet, we should know how far this can be accomplished by simple restriction in the amount of food without complicating the problem by other factors.

In planning the conduct of this series of experiments the writer has clearly recognized that, while it may be possible, as previous experiments have shown, to maintain body equilibrium and nitrogen equilibrium on a low proteid diet for a brief period, this fact does not, as Munk has previously pointed out, by any means establish the view that such a diet will prove efficient in maintaining equilibrium for a long period, or that bodily strength and vigor can be kept up and the proper resistance to disease secured. Hence, it seemed

necessary to so arrange the experiments that they should continue not for a few days or weeks merely, but through months and years. Further, it is very questionable whether the restricted diet (restricted in variety) frequently made use of for convenience in ordinary metabolism experiments is well adapted for bringing out the best results. Hence, it was decided to avoid so far as possible any monotony of diet, giving due rec ognition to the psychical influences liable to affect secretion, digestion, etc., so admirably worked out by Pawlow in his classical experiments on these subjects; influences which are unquestionably of great importance in controlling and modifying, in some measure at least, the nutritive changes in the Again, it is evident that to have experiments of this character broadly useful, they must be tried upon a large number of people and under different conditions of life, in order to avoid so far as possible the influence of personal idiosyncrasy and thereby escape misleading conclusions.

The experiments have been conducted with three distinct types or classes of individuals:

- 1st. A group of five men of varying ages, connected with the University as professors and instructors; men who while leading active lives have not engaged in very active muscular work. They were selected as representatives of the mental worker rather than the physical worker, although several of them in the performance of their daily duties had to be on their feet in the laboratory a good portion of the day.
- 2d. A detail of thirteen men, volunteers from the Hospital Corps of the United States Army and representatives of the moderate worker; men who for a period of six months took each week day a vigorous amount of systematic exercise in the gymnasium, in addition to the routine work connected with their daily life as members of the United States Hospital Corps. These men were of different nationalities, ages, and temperaments.
- 3d. A group of eight young men, students in the University, all thoroughly trained athletes, and some of them with exceptional records in athletic events.

### I. EXPERIMENTS WITH PROFESSIONAL MEN.

Before proceding with a detailed account of the experimental work, it may be well again to emphasize that what is especially desired is to ascertain how far, if any, the intake of proteid food can be diminished without detriment to the body, i. e., with maintenance of nitrogen and body equilibrium and without impairment of bodily and mental vigor. Further, if a lower proteid standard than that generally adopted can be established, it is desirable to ascertain whether it can be maintained indefinitely, or for a long period of time, without loss of strength and vigor. Obviously, it is of primary importance that we should know quite definitely what the minimal proteid requirement of the healthy man per kilo of body-weight really is, and the experimental work about to be detailed has aimed especially to determine whether it is possible to materially lower the amount of daily proteid food, without detriment to the bodily health and with maintenance of physical and mental vigor.

The writer, fully impressed with his responsibility in the conduct of an experiment of this kind, began with himself in November, 1902. At that time he weighed 65 kilos, was nearly 47 years of age, and accustomed to eating daily an amount of food approximately equal to the so-called dietary standards. Recognizing that the habits of a lifetime should not be too suddenly changed, a gradual reduction was made in the amount of proteid or albuminous food taken each day. In the writer's case, this resulted in the course of a month or two in the complete abolition of breakfast, except for a small cup of coffee. A light lunch was taken at 1.30 P. M., followed by a heavier dinner at 6.30 P. M. Occasionally, however, the heartier meal was taken at noontime, as the appetite suggested. It should be added that the total intake of food was gradually diminished, as well as the proteid constituents. no change, however, to a vegetable diet, but a simple introduction of physiological economy. Still, there was and is now a distinct tendency toward the exclusion of meat in some measure, the appetite not calling for this form of food in the same degree as formerly. At first, this change to a smaller amount of food daily was attended with some discomfort, but this soon passed away, and the writer's interest in the subject was augmented by the discovery that he was unquestionably in improved physical condition. A rheumatic trouble in the knee joint, which had persisted for a year and a half and which only partially responded to treatment, entirely disappeared (and has never recurred since). Minor troubles, such as "sick headaches" and bilious attacks, no longer appeared periodically as before. There was greater appreciation of such food as was eaten; a keener appetite and a more acute taste seemed to be developed, with a more thorough liking for simple foods. By June, 1903, the body-weight had fallen to 58 kilos.

During the summer the same simple diet was persisted in a small cup of coffee for breakfast, a fairly substantial dinner at midday and a light supper at night. Two months were spent in Maine at an inland fishing resort, and during a part of this time a guide was dispensed with and the boat rowed by the writer frequently six to ten miles in a forenoon, sometimes against head winds (without breakfast), and with much greater freedom from fatigue and muscular soreness than in previous years on a fuller dietary. The test of endurance and fitness for physical work which the writer thus carried out "on an empty stomach" tended to strengthen the opinion that it is a mistake to assume the necessity for a hearty meal because heavy work is about to be done. It is certainly far more rational from a physiological standpoint to leave the hearty meal until the day's work is accomplished. We seemingly forget that the energy of muscular contraction comes not from the food-stuffs present at the time in the stomach and intestinal tract, but rather from the absorbed material stored up in the muscles and which was digested and absorbed a day or two before. Further, it is to be remembered that the very process of digestion draws to the gastro-intestinal tract a large supply of blood, and that a large amount of energy is needed for the processes of secretion, digestion, absorption, and

peristalsis, which are of necessity incited by the presence of food in the stomach and intestine, thereby actually diminishing the amount of energy available at the place where it is most needed. Why, then, draw upon the resources of the body just at a time, or slightly prior to the time, when the work we desire to perform, either muscular or mental, calls for a copious blood supply in muscle or brain, and when all available energy is needed for the task that is to be accomplished?

We are too wont to compare the working body with a machine, the boiler, engine, etc., overlooking the fact that the animal mechanism differs from the machine in at least one important respect. When we desire to set machinery in operation we must get up steam, and so a fire is started under the boiler and steam is generated in proportion as fuel is burned. The source of the energy made use of in moving the machinery is the extraneous combustible material introduced into the fire-box, but the energy of muscular contraction, for example, comes not from the oxidizable food material in the stomach, but from the material of the muscle itself. In other words, in the animal body it is a part of the tissue framework, or material that is closely incorporated with the framework, that is burned up, and the ability to endure continued muscular strain depends upon the nutritive condition of the muscles involved, and not upon the amount of food contained in, or introduced into, the stomach. All physiologists will, I think, acknowledge the soundness of this reasoning, but how few of us apply the principle in practice. It is perfectly logical to begin the work of the day with a comparatively empty stomach, - after we have once freed ourselves from the habit of a hearty breakfast, - and in the writer's experience both mental and physical work have become the easier from this change of habit. muscle and the brain are given opportunity to repair the waste they have undergone, by the taking of food at times when the digestive processes will not draw upon the energy that in activity is needed elsewhere.

Further, it is easy to understand why on a restricted diet, especially of proteid foods, there should be a diminished sense

22

of fatigue in connection with vigorous or continued muscular work, and why at the same time there should be an increased power of endurance, with actual increase of strength. With a diminished intake of proteid food there is a decreased formation of crystalline nitrogenous waste products, such as uric acid and the purin bases, to say nothing of other bodies less fully known, which circulating through the system are undoubtedly responsible, in part at least, for what we term fatigue. We need not consider here whether the sense of fatigue is due to an action of these substances upon the muscles themselves, upon the motor nerves or their endplates, or upon the central nervous system; it is enough for the present purpose to emphasize the probable results of their presence in undue amount. Lastly, we may emphasize what is pretty clearly evident to-day, viz., that the energy of muscular contraction comes preferably from the oxidation, not of the nitrogenous or proteid constituents of the muscles, but of the non-nitrogenous components of the tissue; another reason why excess of proteid food may be advantageously avoided. Moreover, proteid food stimulates body metabolism in general, and hence undue amounts of proteid in the diet augment unnecessarily the metabolism or combustion of the non-nitrogenous material of the muscle, thereby destroying what would otherwise be preserved as a source of energy in muscular contraction, when the muscles are called upon for the performance of their daily functions.

On the writer's return to New Haven in the fall of 1903, he was surprised to find that his body-weight was practically the same as early in July. In the period between November, 1902, and July, 1903, the body had lost 8 kilos under the gradual change of diet, but from July to October, 1903, the weight had apparently remained stationary, from which it might fairly be assumed that the body had finally adjusted itself to the new conditions.

What now was the condition of the body as regards nitrogen metabolism? To answer this question the entire twenty-four hours' urine was collected practically every day, from

October 13, 1903, to June 28, 1904, representing a period of nearly nine months. This daily output through the kidneys was analyzed each day with special reference to the total nitrogen,\* as a measure of the amount of proteid material metabolized. Total volume of the urine, specific gravity, uric acid, phosphoric acid, indican, and other points were also considered, the more important results being indicated in the following tables.

Phosphoric acid was estimated by titration with a standard uranium solution, using potassium ferrocyanide as an indicator.

<sup>\*</sup> All figures for nitrogen throughout the book, whether referring to food, urine, or fæces, were obtained by exact chemical analysis, using the Kjeldahl-Gunning method.

Uric acid was determined by the method of Folin, i.e., precipitation of the urine with ammonium sulphate, etc., and titration with potassium permanganate.

At times, as will be seen from the tables, nitrogen, uric acid, etc., were not determined in each day's urine. In such cases, an aliquot part of each twenty-four hours' urine was taken and the analyses made with the mixed samples for the given period, the figures thus obtained showing the average daily composition for that period.

# 24 PHYSIOLOGICAL ECONOMY IN NUTRITION

Date.	Body-		Urine.						
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>5</sub> .			
1903	kilos	c.c.		grams	gram	grams			
Oct. 18	57.5	380	1027	5.46	0.376	1.10			
14	l <b>.</b>	550	1027	6.51					
15	1	400	1022	5.40	0.352	1.02			
16		500	1027	6.45					
17	57.4	477	1030	6.40					
18		670	1027	7.20	0.406	1.10			
19		435	1027	6.13					
20		465	1028	6.83	0.476	0.92			
21		450	1029	6.51					
22		465	1027	6.14	0.370	0.96			
27		445	1028	5.52	0.385	0.79			
28		405	1027	5.08	• • •				
Nov. 1		890	1029	5.68	0.872	0.42			
2		530	1027	6.83					
8		470	1027	5.92	0.412	0.75			
4		425	1028	5.88					
5		375	1029	4.93	0.830	0.79			
7		545	1028	6.57					
8	57.4	456	1029	5.82	0.371	0.65			
9		415	1027	5.43					
10		615	1025	6.45	0.430	1.17			
11		410	1028	4.80					
12		580	1026	5.64	0.371	1.02			
13		584	1027	5.82					
14		505	1029	6.86					
15		405	1028	5.80	0.384	0.73			
16		425	1027	5.43					
17		455	1028	5.27	0.367	0.76			
18		575	1027	6.62					
19		447	1027	5.34	0.389	0.77			
20		480	1029	6.00					
21	57.5	400	1029	5.71					
22		382	1029	5.52	0.379	0.97			
23	57.7	850	1029	5.88					
24		422	1029	6.43	0.400				
25		435	1030	5.79					
26	57.6	445	1030	6.09	0.430	1.01			
27	l	430	1030	6.17					
29	l	454	1027	5.66	0.420	0.98			
80	<b> </b>	455	1028	5.56					
L	<u> </u>		L						

D.4.	Body-		Urine.							
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>3</sub> .				
1903	kilos	c.c.		grams	gram	grams				
Dec. 1	1	420	1028	5.31	0.450	0.92				
2		465	1027	6.17						
8		430	1027	5.84	0.350	0.82				
4		865	1029	4.77						
5		430	1030	5.83						
6	1	515	1028	5.90	0.393	1.08				
7	·	400	1028	5.57						
8		390	1028	4.99	0.328	1.02				
9		405	1080	5.17						
10	57.6	370	1026	4.64	0.308	0.90				
11		827	1030	4.65	0.825	1.11				
12		890	1027	5.16	0.346	1.01				
18		429	1029	5.66						
14		360	1080	4.84	• • •					
15		295	1029	4.32	0.291	0.82				
16		445	1029	6.27						
17		890	1032	5.59	0.358	0.73				
18	1	420	1030	5.62		• • •				
19		415	1027	5.03						
20	57.5	390	1030	5.71	0.402	0.78				
21		360	1023	4.25						
22		860	1030	5.13	0.342	0.79				
23		400	1031	5.08						
24		435	1080	6.44						
25		450	1029	5.13	0.329	0.77				
26		465		5.55	• • • •					
27		470	• • •	5.53						
28		535		8.18						
29		535		7.67		• • •				
80		656		9.68						
81	57.6	490	1031	7.61	0.455	0.92				
1904										
Jan. 1	• • •	415	1030	6.41	• • •	• • •				
2		490	1081	6.56		• • •				
8	• • •	460	1030	5.91	0.819	0.79				
4	58.1	430	1030	5.72	• • •					
5		570	1028	6.86	0.402					
6		445	1028	5.68	• • •					
7		510	1028	5.91	0.367	0.99				
8	• • • •	420	1028	5.87		• • •				

	Podw			Urine.		
Date.	Body- weight.	Volume. 24 hours.	une. Sp. Gr. Nitrogen.		Uric Acid.	P <sub>2</sub> O <sub>8</sub> .
1904	kilos	0.0.		grams	gram	gram
Jan. 9		650	1027	7.29		
10		685	1024	6.82	0.414	0.99
11		410	1028	4.87		
12		450	1027	6.48	• • •	
13		410	1027	5.34	0.435	
14		532	1028	6.22	0.502	
15		530	1028	5.98	• • •	
16		515	1030	6.18		
17		537	1030	6.78	0.429	
18	57.8	395	1029	5.09	• • •	• • •
19		450	1080	5.72	0.427	• • • •
20		420	1026	4.76	• • •	• • •
21		410	1029	5.26	0.401	• • •
22		485	1029	5.41		
23		440	1081	5.07		• • •
24		485	1029	5.61	0.407	• • •
25		545	1027	6.18	• • •	• • •
26		485	1028	6.69	0.440	• • •
27		485	1028	5.64		• • •
28		490	1029	6.18	0.428	• • •
29		450	1029	5.68	• • •	• • •
80		475		5.59	0.376	• • •
81		490	••••	6.61	• • • •	• • •
Feb. 1		490	1030	6.47	• • •	• • •
2	57.5	400	1031	6.12	0.219	
8		415	1080	5.85	• • •	• • •
4	• • • •	545	1027	6.77	0.327	• • •
5		450	1030	5.64	• • •	• • •
6		485	1027	6.01		• • •
7		450	1026	5.62		• • •
8	<b>57.4</b>	415	1027	5.88	:::	• • •
9		540	1026	6.67	0.449	• • • •
10	1	410	1029	5.61	• • •	•••
11		600	1025	6.70		• • •
12	• • •	430	1029	5.57	0.437	• • •
18	1	415	1028	5.50	0.407	•••
14		480	1028	6.42	0.497	•••
15	•••	895	1080	4.95	0.004	• • •
16		500	1029	5.97	0.864	• • •
17		450	1030	5.62	• • •	• • •

Date.	Body- weight.			Urine.		
<i>Data</i> .	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>8</sub> .
1904	kilos	c.c.		grams	gram	grams
Feb. 18	57.3	430	1030	5.86	0.279	
19		450	1029	6.21		
20		455	1027	5.46		
21		500	1027	6.16	0.424	
22		445	1028	5.15		
28		455	1027	5.63		
24		420	1028	6.27		
25		560	1027	6.28	0.403	
26		630	1026	6.27	• • •	
27		570	1026	6.87		
28		515	1028	6.27	0.496	
29		450	1030	5.43		
Mar. 1		450	1027	6.02		
2		445	1029	5.15		
8		590	1028	6.30		
4		415	1029	5.40		
5	57.5	425	1027	5.48		
6		548	1025	5.92	0.870	
7	1	400	1029	4.68		
8		530	1028	5.77		
.9		560	1028	5.84		
10		560	1028	5.64		
11		495	1028	5.79		
12		515	1021	6.80		
18	1	520	1029	6.48	0.870	
14	1	600	1025	6.12		
15	1	520	1026	5.87		
16	57.5	525	1026	5.18		
17		490	1026	4.97		
18		450	1027	5.08		
19		500	1024	5.85		
20		500	1022	5.91	lì	
<b>21</b> .	. 57.4	430	1025	5.52		
22		458	1088	5.94	1	
23	57.2	400	1029	5.61	0.821	1.20
24	57.8	865	1029	4.31	daily	daily
25		420	1029	5.39	average.	average.
26	57.5	435	1027	5.85		
27		595	1026	6.38		
28		545	1027	6.00	• • •	•••

CHITTENDEN.

Date.	Body- weight.		Urine.					
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> .		
1904	kilos	c.c.		grams	gram	gram		
Mar. 29	• • •	435	1028	4.86				
80		575	1026	6.26				
31		495	1026	5.26				
Apr. 1		570	1026	6.33				
2		440	1030	6.07				
3		487	1026	6.11	0.875			
4		410	1028	5.78				
5		390	1028	5 38				
6	<b>56</b> .8	490	1028	5.56				
7	56.5	530	1027	5.69				
8		440	1029	5.41				
9		465	1019	6.05				
10	56.8	500	1029	6.00	0.382			
11		500	1028	6.18				
12	56.4	475	1029	5.55	0.366	0.870		
Daily aver. for ?		400	100		0.000	0 000		
six months.	• • •	466	1027	5.82	0.386	0.899		
13		545	1029	6.77				
14		440	1027	5.89				
15		500	1028	5.91				
16		485	1028	5.49				
17		405	1029	5.99	0.393			
18		465	1029	6.11				
19		510	1030	7.68				
20		430	1031	6.99				
21	56.6	615	1029	8.67				
22		320	1080	5.03				
23	57.1	855	1082	5.72				
24		455	1027	5.97				
25		380	1027	4.93				
26		450	1028	4.97	0.866			
27		600	1025	6.62	0.558			
28	56.9	385	1029	5.66	0.507			
29		415	1029	5.28	0.488			
30	56.9	462	1029	5.59	0.413			
May 1		486	1027	5.54	0.409			
2		405	1028	4.11	0.320			
3	57.1	505	1027	5.48				
4		456	1026	5.27				
5		380	1026	4.88				
		1						

CHITTENDEN.

Date.	Body-			Urine.		
Duig.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>8</sub> .
1904	kilos	c.c.		grams	gram	gram
May 6		580	1027	6.80		
7		470	1024	5.44		
8	57.6	460	1027	5.07		
9		460	1026	4.28	• • •	
10	57.4	498	1028	5.26		
11		415	1029	4.61		
12		530	1029	5.98		
13		415	1031	4.72	• • •	
14	57.2	405	1081	4.98	0.468	
15		500	1029	5.31		
16		505	1027	5.03		
17		650	1020	5.69		
18		550	1027	5.81		
19		560	1027	6.05		
20		615	1027	6.64		
21	56.9	380	1082	5.20	0.421	
22		475	1028	5.73		
28		378	1028	4.60		
24		388	1029	4.48		
25	1	535	1025	5.14		
26	56.9	355	1028	4.37		
27	1	435	1026	4.93	`	
28	57.5	555	1028	5.99	0.397	
29	57.7	565	1027	6.27		
30		700	1020	5.50		
31	1	500	1025	5.18		
June 1	1	630	1028	5.41		
2		510	1020	4.16		
8		530	1028	5.25		
4	57.6	890	1029	5.25		
5		400	1025	4.87		
6		480	1027	5.16		
7		480	1028	5.15		
8		410	1027	4.95		
9		420	1026	4.51		
10		395	1026	4.27		
11	57.5	510	1030	5.91		
12		580	1027	5.95		
18	57.6	485	1027	5.35		
14		470	1030	5.16	• • •	• • •

CHITTENDEN.

Date.	Body-			Urine.		
Dave.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>5</sub> .
1904	kilos	c.c.		grams	gram	grams
June 15		560	1024	4.91		
16	57.6	390	1029	5.26		
17		408	1027	5.17		
18	57.9	412	1030	5.07		
19		458	1025	5.44		
20	58.0	380	1026	4.49		l
21		480	1023	5.04		
22	57.8	580	1025	6.16		
23	57.9	585	1025	5.26	1	1
24	57.6	570	1024	5.30		
25		410	1027	4.43	} 0.897	1.08
26	57.4	400	1027	4.66	daily	daily
27	57. <b>4</b>	405	1027	4.98	average.	average.
Daily av. from } Oct. 13, 1903		468	1027	5.69	0.392	0.904
Daily av. from April 13 to June 27	•••			5.40	• • •	
June 28	57.5	595	1026	6.75	• • •	• • •

Scrutiny of the tables shows that during this period of nine months the body-weight was practically constant. The daily volume of urine was exceptionally small and fairly regular in amount, the average daily output for the nine months being 468 c.c. It is a noticeable fact that with a diminished intake of proteid food there is far less thirst, and consequently a greatly decreased demand for water or other fluids. Further, in view of the small nitrogenous waste there is no need on the part of the body for any large amount of fluid to flush out the kidneys. The writer has not had a turbid urine during the nine months' period. With heavier eating of nitrogenous foods, an abundant water supply is a necessity to prevent the kidneys from becoming clogged, thereby explaining the frequent beneficial results of the copious libations of mineral

waters, spring waters, etc., frequently called for after, or with, heavy eating. Obviously, a small volume of urine each day means so much less wear and tear of the delicate mechanism of the kidneys. Somewhat noticeable, in a general way, is the apparent relationship between the volume of the urine and the nitrogen output, in harmony with the well-known diuretic action of urea. The specific gravity of the urine shows variation only within narrow limits, the daily average for the nine months being 1027.

Uric acid is noticeably small in quantity, the average daily output for the nine months' period, based upon the determinations made, being only 0.392 gram.

Chief interest, however, centres around the figures for total nitrogen, since these figures give for each day the extent of the proteid metabolism; i.e., the amount of proteid material broken down in the body each day in connection with the wear and tear of the bodily machinery. To fully grasp the significance of these data, it should be remembered that the prevalent dietary standards are based upon the assumption that the average adult must metabolize each day at least 16 grams of nitrogen. Indeed, that is what actual analysis of the urine indicates in most cases. If now we look carefully through the figures shown in the above tables, covering a period from October 13, 1903, to June 28, 1904, it is seen that the daily nitrogen excretion is far different from 16 grams. Indeed, the figures for nitrogen are exceedingly low, and, moreover, they vary little from day to day. The average daily output of nitrogen through the urine for the entire period of nearly nine months is only 5.699 grams.

For the first six months the average daily excretion amounted to 5.82 grams of nitrogen, while from April 12 to June 28 the average daily excretion of nitrogen was 5.40 grams, thus showing a slight tendency downward. On the whole, however, there is shown a somewhat remarkable uniformity in the daily excretion. Thus, the average daily excretion for the month of November was 5.79 grams of nitrogen, for the month of March 5.66 grams, thus showing very little

difference in the output of nitrogen through the kidneys in these two periods, three months apart. In other words, the extent of proteid katabolism was essentially the same throughout the entire nine months, implying that the amount of proteid food eaten must have been fairly constant, and that the body had adapted itself to this new level of nutrition from which there was no tendency to deviate. There was no weighing out of food and no attempt to follow any specified The greatest possible variety of simple foods was indulged in, and the dictates of the appetite were followed with the single precaution that excess was avoided. In other words, it was temperance in diet, and not prohibition. it is equally true, in the writer's case at least, that the appetite itself unconsciously served as a regulator, since there was, as a rule, no necessity to hold the appetite in check to Doubtless, the writer's knowledge of the genavoid excess. eral composition of food-stuffs has had some influence in the choice of foods, and thereby aided in bringing about this somewhat remarkable uniformity in the daily output of nitrogen for such a long period of time on an unrestricted diet.

What now do the nitrogen figures show regarding the amount of proteid material metabolized each day? It will be remembered that the Voit standard calls for 118 grams of proteid or albuminous food daily, of which 105 grams should be absorbable, in order to maintain the body in a condition of nitrogen equilibrium, and in a state of physical vigor and general tone. This would mean a daily excretion through the urine of at least 16 grams of nitrogen. The daily output of nitrogen in the case under discussion, however, was 5.699 grams for a period of nearly nine months. This amount of nitrogen excreted through the urine means only 35.6 grams of proteid metabolized, or about one-third the amount called for by the Voit standard, or the standards generally adopted as expressing man's daily requirement of proteid food. But was the body in nitrogenous equilibrium on this small amount of proteid food? Naturally, this question might be answered in the affirmative, on the basis of the constancy in bodyweight for the period from October to June, but more decisive proof is needed. The question was therefore settled by a careful comparison of the income and output, in which all the food eaten was carefully weighed and analyzed, while the nitrogen of the urine and fæces was determined with equal accuracy. The first experiment of this character to be quoted is for the week commencing March 20, a period of six days.

Following are the diets made use of each day, the weights of the various food-stuffs being given in grams. Likewise is shown the nitrogen content of the several food-stuffs for each day, and also a comparison of the nitrogen intake with the output of nitrogen through the urine:

### Sunday, March 20, 1904.

- Breakfast, 7.45 A.M.—One cup coffee, i.e., coffee 137.5 grams, cream 30.5 grams, sugar 9 grams.
- Dinner, 1.30 p. m. Stewed chicken 50 grams, mashed potato 131 grams, biscuit 49 grams, butter 13 grams, chocolate pudding 106 grams, one small cup coffee, i.e., coffee 64 grams, sugar 12 grams, cheese crackers 29 grams.
- Supper, 6.30 p. m.—Lettuce sandwiches 56 grams, biscuit 35 grams, butter 6 grams, one cup tea, i. e., tea 170 grams, sugar 7 grams, sponge cake 47 grams, sliced oranges 82 grams.

Food.	Grams.	Per o	ent Nitroge	n.•	Total Nitrogen.
Coffee $.64 + 137.5 =$	201.5	×	0.042	<b>'</b> =	0.085 gram.
Cream	30.5	×	0.41	=	0.125
Sugar $12+9+7 =$	28.0	×	0.00	=	0.000
Chicken	50.0	×	4.70	=	2.350
Mashed potato	131.0	×	0.30	=	0.393
Biscuit $35 + 49 =$	84.0	×	1.49	=	1.251
Butter $.18 + 6 =$	19.0	×	0.10	=	0.019
Chocolate pudding	106.0	×	0.86	=	0.911
Cheese crackers	29.0	×	2.54	=	0.787
Lettuce sandwich	56.0	×	0.92	=	0.515
Tea	170.0	×	0.048	=	0.082
Sponge cake	47.0	×	0.98	=	0.461
Sliced orange	82.0	×	0.078	=	0.060
Total nitroger	n in food				6.989 grams.
Total nitroger	n in prin	Α.			_

Fuel value of the food . . . 1708 calories.

<sup>\*</sup> All foodstuffs were analyzed from large samples, to diminish as much as possible the errors of analysis. Nitrogen was determined by the Kjeldahl-Gunning method, the figures given being the average of closely agreeing duplicate analyses.

While nitrogen was thus determined in every sample of food by direct chemical analysis, the fuel value of the food was calculated mainly by use of the data furnished by the Bulletin issued from the U. S. Department of Agriculture, Office of Experiment Stations. No. 28.

### Monday, March 21, 1904.

Breakfast, 7.45 A.M. — Coffee 119 grams, cream 30 grams, sugar 9 grams.

Lunch, 1.30 P. M. — One shredded wheat biscuit 31 grams, cream 116 grams, wheat gems 33 grams, butter 7 grams, tea 185 grams, sugar 10 grams, cream cake 53 grams.

Dinner, 6.80 p.m. — Pea soup 114 grams, lamb chop 24 grams, boiled sweet potato 47 grams, wheat gems 76 grams, butter 13 grams, cream cake 52 grams, coffee 61 grams, sugar 10 grams, cheese crackers 16 grams.

Food.	•	Grams.	Per	. Ce	nt Nitrog	m.	Total Nitrogen.
Coffee 119 + 61	=	180	×		0.042	=	0.076 gram.
Cream 30 + 116	=	146	×		0.41	=	0.600
Sugar $9 + 10 + 10$	=	29	×		0.00	=	0.000
Shredded wheat biscuit		31	×		1.62	=	0.502
Tea		185	×		0.048	=	0.089
Wheat gems $83 + 76$	=	109	×		1.46	=	1.591
Butter 7 + 18	=	20	×		0.10	=	0.020
Cream cake $53 + 52$	=	105	×		0.97	=	1.018
Pea soup		114	×		1.00	=	1.140
Lamb chop		24	×		4.54	=	1.090
Sweet potato		47	×		0.18	=	0.085
Cheese crackers		16	×		2.54	=	0.410
Total nitroge	n i	n food					6.621 grams.
Total nitroge	en i	n urine					5.520

Fuel value of the food . . . 1713 calories.

### Tuesday, March 22, 1904.

Breakfast, 7.45 A. M. — Coffee 97 grams, cream 26 grams, sugar 9 grams.

Lunch, 1.30 P.M. — Baked potato 83 grams, fried sausage 36 grams, soda biscuit 39 grams, butter 12 grams, tea 137 grams, sugar 10 grams, cream meringue 59 grams.

Dinner, 6.30 P. M. — Chicken broth 146 grams, bread 52 grams, butter 15 grams, creamed potato 76 grams, custard 76 grams, coffee 50 grams, sugar 11 grams, cheese crackers 10 grams.

Food.		Grams.	Per cer	nt Nitrogen.	Total Nitrogen.
Coffee	.97 + 50 =	147	×	0.042 =	0.060 gram.
Cream		26	×	0.42 =	0.109
Sugar 9	9 + 10 + 11 =	80	×	0.00 =	0.000
Baked potato .		88	×	0.40 =	0.332
Fried sausage		36	×	3.06 =	1.101
Soda biscuit .		39	×	1.66 =	0.647
Butter	. 12 + 15 =	27	×	0.10 =	0.027
Tea		137	×	0.048 =	0.066
Cream meringue		59	×	0.92 =	0.548
Chicken broth .		146	×	0.78 =	1.138
Bread		<b>52</b>	×	1.66 =	0.863
Creamed potato		76	×	0.42 =	0.819
Custard		76	×	0.82 =	0.628
Cheese crackers		10	×	2.54 =	0.254
	Total nitrogen	in food			6.082 grams
	Total nitrogen	in urine			5.940

Fuel value of the food . . . 1398 calories.

## Wednesday, March 23, 1904.

Breakfast, 7.45 A. M. — Coffee 103 grams, cream 80 grams, sugar 10 grams.

Lunch, 1.80 p. m. — Creamed codfish 64 grams, potato balls 54 grams, biscuit 44 grams, butter 22 grams, tea 120 grams, sugar 10 grams, wheat griddle cakes 133 grams, maple syrup 108 grams.

Dinner, 6.30 p. m. — Creamed potato 85 grams, biscuit 53 grams, butter 15 grams, apple-celery-lettuce salad 50 grams, apple pie 127 grams, coffee 67 grams, sugar 8 grams, cheese crackers 17 grams.

Food.	Gr	ums. Per cent Nitro	gen. Total Nitrogen.
Coffee	103 + 67 = 1	70 × 0.042	= 0.071 gram.
Sugar 10 +	10 + 8 =	28 × 0.00	= 0.000
Cream		30 × 0.48	= 0.129
Potato balls		54 × 0.68	= 0.367
Creamed codfish .		64 × 1.26	= 0.806
Biscuit	44 + 53 =	97 × 1.66	= 1.610
Butter	22 + 15 =	37 × 0.10	= 0.037
Tea	1	20 × 0.048	= 0.058
Wheat griddle cake	es 1	$33 \times 1.82$	= 1.760
Maple syrup	1	08 × 0.019	= 0.021
Creamed potato .		85 × 0.53	= 0.450
Cheese crackers .		$17 \times 2.54$	= 0.431
Apple-celery salad		50 × 0.20	= 0.100
Apple pie	1	27 × 0.75	= 0.953
T	otal nitrogen in	food	6.798 grams.
T	otal nitrogen in	urine	5.610

Fuel value of the food . . . 1984 calories.

## Thursday, March 24, 1904.

Breakfast, 7.45 A. M. — Coffee 100 grams, cream 25 grams, sugar 8 grams.

Lunch, 1.80 p. m. — Shredded wheat biscuit 29 grams, cream 118 grams, wheat gems 60 grams, butter 8 grams, tea 100 grams, sugar 7 grams, apple pie 102 grams.

Dinner, 6.30 P. M. — Milk-celery soup 140 grams, bread 15 grams, butter 1 gram, lettuce sandwiches 62 grams, tea 100 grams, sugar 10 grams, lemon pie 109 grams.

Food.	Grams.	Per c	ent Nitrog	en. To	tal Nitrogen.
Coffee	100	×	0.042	=	0.042 gram.
Cream 25 + 118	= 143	×	0.43	=	0.615
Sugar $8 + 7 + 10$	= 25	×	0.00	=	0.000
Shredded wheat biscuit .	29	×	1.76	=	0.510
Wheat gems	60	×	1.17	=	0.702
Butter 8 + 1	= 9	×	0.10	=	0.009
Tea 100 + 100	= 200	×	0.048	=	0.096
Apple pie	102	×	0.75	=	0.765
Milk-celery soup	140	×	0.42	=	0.588
Bread	15	×	1.36	=	0.204
Lettuce sandwich	62	×	1.02	=	0.632
Lemon pie	109	×	0.82	=	0.894
Total nitro	gen in food				5.057 grams.
	gen in urin				4.810

Fuel value of the food . . . 1594 calories.

## Friday, March 25, 1904.

Breakfast, 7.45 a. m. — Coffee 100 grams, cream 25 grams, sugar 9 grams.

Lunch, 1.30 p. m. — Halibut with egg sauce 108 grams, mashed potato 89 grams, biscuit 48 grams, butter 10 grams, chocolate-cream cake 90 grams, tea 100 grams, sugar 9 grams.

Dinner, 6.30 p. m. — Milk-celery soup 121 grams, lettuce sandwiches 61 grams, creamed potato 65 grams, lettuce-apple-celery salad 74 grams, coffee 70 grams, sugar 10 grams.

Food. G	rams.	Per cer	t Nitroge	n. Tot	al Nitrogen.
Coffee $100 + 70 =$	170	×	0.042	=	0.071 gram.
Cream	25	×	0.40	=	0.100
Sugar $9+9+10 =$	28	×	0.00	=	0.000
Halibut, etc	108	×	3.02	=	3.262
Mashed potato	89	×	0.26	=	0.281
Biscuit	48	×	1.52	=	0.780
Butter	10	×	0.10	=	0.010
Tea	100	×	0.048	=	0.048
Chocolate-cream cake	90	×	0.99	=	0.891
Celery-milk soup	121	×	0.52	=	0.629
Lettuce sandwich	61	×	0.98	=	0.598
Lettuce-apple salad	74	×	0.21	=	0.155
Creamed potato	65	×	0.37	=	0.241
Total nitrogen in	food .				6.966 grams.
Total nitrogen in	urine				5.390

Fuel value of the food . . . 1285 calories.

#### NITROGEN BALANCE. - Chittenden.

	Nitrogen	Out	put.
	Taken in.	Nitrogen in Urine.	Weight of Faces * (dry).
March 20	6.989 grams.	5.91 grams.	3.6 grams.
21	6.621	5.52	0.0
22	6.082	5.94	12.0
23	6.793	5.61	18.5
24	5.057	4.31	23.0
25	6.966	5.39	16.9
			74.0 grams contain
			6.42% N.
	88.508	82.68 +	4.75 grams nitrogen.

38.508 grams nitrogen. 37.43 grams nitrogen.

Nitrogen balance for six days = 1.078 grams. Nitrogen balance per day = +0.179 gram.

#### Average Intake.

Calories per day . . . . . . . . . . . . . . . . . 1618.

Nitrogen per day . . . . . . . . . . . . . . . . . 6.40 grams.

<sup>\*</sup> The fæces of this period were separated by lampblack. They were dried on a water-bath after admixture with alcohol and a little sulphuric acid, nitrogen being determined by the Kjeldahl-Gunning method on samples of the dry mixture from the six-day period.

Examination of the results shown in the foregoing balance makes it quite clear that the body was essentially in nitrogenous equilibrium. Indeed, there was a slight plus balance, showing that even with the small intake of proteid food the body was storing up nitrogen at the rate of 0.16 gram per day. The average daily intake of nitrogen for the six days' period was 6.40 grams, equal to 40.0 grams of proteid or albuminous food. The average daily output of nitrogen through the urine and fæces was 6.24 grams. The average daily output of nitrogen through the urine for the six days' period was 5.44 grams, corresponding to the metabolism of 34 grams of proteid material. When these figures are contrasted with the usually accepted standards of proteid requirement for the healthy man, they are certainly somewhat impressive, especially when it is remembered that the body at that date had been in essentially this same condition for at least six months, and probably for an entire year. The Voit standard of 118 grams of proteid, with an equivalent of at least 18 grams of nitrogen and calling for the metabolism of 105 grams of proteid, or 16.5 grams of nitrogen per day, makes clear how great a physiological economy had been accomplished. other words, the consumption of proteid food was reduced to at least one-third the daily amount generally considered as representing the average requirement of the healthy man, and this with maintenance of body-weight at practically a constant point for the preceding ten months, and, so far as the writer can observe, with no loss of vigor, capacity for mental and physical work, or endurance. Indeed, the writer is disposed to maintain that he has done more work and led a more active life in every way during the period of this experiment, and with greater comfort and less fatigue than usual. His health has certainly been of the best during this period.

In this connection it may be well to call attention to the completeness of the utilization of the daily food in this six days' experiment, as shown by the small amount of refuse discharged per rectum, indicating as it does the high efficiency of the digestive processes and of the processes of absorption.

The refuse matter for the entire period of six days amounted when dry to only 74 grams, and when it is remembered how large a proportion of this refuse must of necessity be composed of the cast-off secretions from the body, it will be seen how thorough must have been the utilization of the food by the system. The loss of nitrogen to the body per day through the fæces amounted to only 0.79 gram, and this on a mixed diet containing considerable matter not especially concentrated, and on some days with noticeable amounts of food, such as salads, not particularly digestible.

Finally, emphasis should be laid upon the fact that this economy of proteid food, this establishment of nitrogen equilibrium on a low proteid intake, was accomplished without increase in the daily intake of non-nitrogenous foods. In fact, the amount of fats and carbohydrates was likewise greatly reduced, far below the minimal standard of 8000 calories as representing the potential energy or fuel value of the daily diet. Indeed, during the balance period of six days just described the average fuel value of the food per day was only a little over 1600 calories.

As the experiment continued and the record for the months of April and May was obtained, it became evident from the nitrogen results that the rate of proteid katabolism was being still more reduced. A second balance experiment was therefore tried with a view to seeing if the body was still in nitrogen equilibrium, and also to ascertain whether the fuel value of the food still showed the same low calorific power. For a period of five days, June 23 to 27, the intake of food and the entire output were carefully compared, with the results shown in the accompanying tables.

## Thursday, June 23, 1904.

Breakfast. — Coffee 123 grams, cream 50 grams, sugar 11 grams.

Lunch. — Omelette 50 grams, French fried potatoes 70 grams, bacon 10 grams, wheat gems 43 grams, butter 9 grams, strawberries 125 grams, sugar 20 grams, cream cake 59 grams.

Dinner.—Beefsteak 34 grams, peas 60 grams, creamed potato 97 grams, bread 26 grams, butter 17 grams, lettuce-orange salad 158 grams, crackers 43 grams, cream cheese 15 grams, coffee 53 grams, sugar 12 grams.

Food.	•	Grams.	Per ce	nt Nitrogen.		Total Nitrogen.
Coffee	. 123 + 53 =	176	×	0.045 =	=	0.079 gram.
Cream		50	×	0.35 =	=	0.175
Sugar 11	1 + 20 + 12 =	43	×	0.00	=	0.000
Omelette		50	×	1.32	=	0.660
French fried pote	toes	70	×	0.37	=	0.259
Bacon		10	×	3.43 =	=	0.843
Wheat gems .		43	×	1.49 :	=	0.641
Butter	. 9 + 17 =	26	×	0.18 =	=	0.034
Strawberries .		125	×	0.11 =	=	0.138
Cream cake .		<b>59</b>	×	0.98 =	=	0.578
Beefsteak		34	×	4.14 =	=	1.408
Peas		60	×	0.97	=	0.582
Creamed potato		97	×	0.34 =	=	0.880
Bread		26	×	1.23 =	=	0.820
Lettuce-orange sa	alad	153	×	0.15	=	0.230
Crackers		43	×	1.40 =	=	0.602
Cream cheese .		15	X	1.62	=	0.243
	Total nitrogen	in food .				6.622 grams.
	Total nitrogen					5.260

Fuel value of the food . . . 1863 calories.

### Friday, June 24, 1904.

Breakfast. - Coffee 96 grams, sugar 8 grams, milk 82 grams.

Lunch.—Creamed codfish 89 grams, baked potato 95 grams, butter 10 grams, hominy gems 58 grams, strawberries 86 grams, sugar 26 grams, ginger snaps 47 grams.

Dinner. — Cold tongue 14 grams, fried potato 48 grams, peas 60 grams, wheat gems 30 grams, butter 11 grams, lettuce-orange salad with mayonnaise dressing 155 grams, crackers 22 grams, cream cheese 14 grams, ginger snaps 22 grams, coffee 58 grams, sugar 10 grams.

Food.		Grams.	Per cent N	itrogen.	Total Nitrogen.
Coffee	. 96 + 58	= 154	× 0.0	)45 <b>=</b>	0.069 gram.
Sugar 8	3 + 26 + 10:	= 44	× 0.0	<b>9</b> 0 =	0.000
Milk		. 32	× 0.6	51 =	0.163
Creamed codfish		. 89	× 1.7	<b>18</b> =	1.584
Baked potato .		. 95	× 0.2	<b>29</b> =	0.276
Butter	. 10 + 11	= 21	× 0.1	13 =	0.027
Hominy gems .		. 58	× 1.5	20 =	0.696
Strawberries .		. 86	× 0.1	1 =	0.095
Ginger snaps .	.47 + 22 :	= 69	× 1.1	15 =	0.794
Cold tongue .		. 14	× 4.8	37 =	0.682
Fried potato .		. 48	× 0.8	37 =	0.178
Peas		. 60	× 0.9	)4 ==	0.564
Wheat gems .		. 30	X 1.4	15 =	0.485
Lettuce-orange sa	alad, etc	. 155	× 0.1	15 =	0.283
Crackers		. 22	× 1.4	10 =	0.308
Cream cheese .		. 14	× 1.6	32 =	0.227
	Total nitroge	n in food			6.831 grams.
	Total nitroge	n in urine			5.300

Fuel value of the food . . . . 1506 calories.

# Saturday, June 25, 1904.

Breakfast. - Coffee 101 grams, milk 36 grams, sugar 18 grams.

Lunch. — Omelette 50 grams, bacon 9 grams, French fried potato 23 grams, biscuit 29 grams, butter 8 grams, cream cheese 17 grams, iced tea 150 grams, sugar 15 grams, ginger snaps 42 grams.

Dinner. — Wheat popovers 57 grams, butter 10 grams, lettuce-orange salad with mayonnaise dressing 147 grams, cream cheese 21 grams, crackers 22 grams, cottage pudding 82 grams, coffee 48 grams, sugar 11 grams.

Food.		Grams.	Per	cent Nitrog	50a.	Total Nitrogen.
Coffee 101 +	<b>1</b> 8 =	149	×	0.045	=	0.067 gram.
Milk	. <b>.</b> .	36	×	0.46	=	0.166
Sugar $18 + 15 + 15$	ll =	89	×	0.00	=	0.000
Omelette		50	×	1.42	=	0.710
Bacon		9	×	2.66	=	0.239
French fried potato		23	×	0.57	=	0.181
Biscuit		29	×	1.85	=	0.892
Butter 8+	10 =	18	×	0.13	=	0.028
Iced tea		150	×	0.018	=	0.027
Ginger snaps		42	×	1.15	=	0.483
Cream cheese 17 +	21 =	38	×	1.62	=	0.616
Wheat popovers		57	×	1.64	=	0.985
Lettuce-orange salad		147	×	0.15	=	0.221
Crackers		22	×	1.40	=	0.808
Cottage pudding		82	×	0.76	=	0.628
Total nit	rogen	in food				4.941 grams.
Total ni	rogen	in urine				4.430

Fuel value of the food . . . . 1392 calories.

## Sunday, June 26, 1904.

Breakfast. —Coffee 122 grams, cream 31 grams, sugar 8 grams.

Dinner. — Roast lamb 50 grams, baked potato 52 grams, peas 64 grams, biscuit 32 grams, butter 12 grams, lettuce salad 43 grams, cream cheese 21 grams, toasted crackers 23 grams, blanc mange 164 grams.

Supper. — Iced tea 225 grams, sugar 29 grams, lettuce sandwich 51 grams, strawberries 130 grams, sugar 22 grams, cream 40 grams, sponge cake 31 grams.

Food.		Grams.	Per cent Nitrog	m.	Total Nitrogen.
Coffee		122	× 0.045	=	0.055 gram.
Cream	. 81 + 40 =	71	$\times$ 0.32	=	0.227
Sugar 8	3 + 29 + 22 =	59	× 0.00	=	0.000
Roast lamb .		50	× 4.28	=	2.140
Baked potato .		52	× 0.29	=.	0.151
Peas		64	× 1.04	=	0.666
Biscuit		82	× 1.35	=	0.482
Butter		12	× 0.13	=	0.016
Lettuce salad .		43	× 0.23	=	0.099
Cream cheese .		21	× 1.62	=	0.840
Toasted crackers		23	× 1.86	=	0.313
Blanc mange .		164	× 0.35	=	0.574
Iced tea		225	× 0.018	=	0.041
Lettuce sandwich	ı . <b>.</b>	51	× 0.85	=	0.484
Strawberries .		130	× 0.11	=	0.148
Sponge cake .		31	× 0.94	=	0.291
	Total nitrogen	in food .			5.922 grams.
	Total nitrogen				4.660

Fuel value of the food . . . 1533 calories.

## Monday, June 27, 1904.

- Breakfast. Coffee 112 grams, cream 22 grams, sugar 10 grams.
- Lunch.—Roast lamb 9 grams, baked potato 90 grams, wheat gems 47 grams, butter 12 grams, sugar 25 grams, iced tea 250 grams, vanilla éclair 47 grams.
- Dinner. Lamb chop 32 grams, asparagus 49 grams, butter 17 grams, creamed potato 107 grams, bread 35 grams, lettuce-orange salad with mayonnaise dressing 150 grams, cream cheese 12 grams, crackers 21 grams, coffee 63 grams, sugar 9 grams.

Food.	Grams.	Per cent Nitr	ogen.	Total Nitrogen.
Coffee $112 + 63 =$	= 175	× 0.045	=	0.079 gram.
Cream	. 22	× 0.82	=	0.070
Sugar $10 + 25 + 9 =$	= <b>44</b>	× 0.00	=	0.000
Iced tea	. 250	× 0.018	=	0.045
Baked potato	. 90	× 0.25	=	0.225
Wheat gems	. 47	× 1.65	=	0.776
Butter $12 + 17 =$	= 29	× 0.13	=	0.038
Roast lamb	. 9	× 4.28	=	0.385
Vanilla éclair	. 47	× 0.85	=	0.400
Lamb chops	. 82	× 4.57	=	1.462
Asparagus	. 49	× 0.59	=	0.289
Creamed potato	. 107	× 0.40	=	0.428
Bread	. 35	× 1.38	=	0.466
Lettuce-orange salad, etc	. 150	× 0.28	=	0.845
Cream cheese	. 12	× 1.62	=	0.194
Crackers	. 21	× 1.35	=	0.284
Total nitroger	in food			. 5.486 grams.
Total nitrogen	in urine			4.980

Fuel value of the food . . . . 1454 calories.

### NITROGEN BALANCE. — CHITTENDEN.

		Nitrogen Taken In.	Nitrogen in U	Out <sub>!</sub> Jrine.	put. Weight of <b>F</b> secon	(dry).
June	23	6.622 grams.	5.26 gr	ams.	10.6	
	24	6.331	5.30		30.7	
	25	4.941	4.43		14.2	
	26	5.922	4.66		11.9	
	27	5.486	4.98		15.2	
					82.6 gram	s contain
					6.	08% N.
		29.302	24.68	+	5.022 gra	ams nitrogen.
		29.302 grams	nitrogen.	29.6	52 grams nitro	gen.
	Nitrog	gen balance for fl	ve days	=	-0.350 gram.	
	Nitrog	gen balance per d	lay	=	-0.070 gram.	
		Av	erage Intake.			
	Calori	es per day			1549	).

Nitrogen per day . . . . . . . . . . . . . 5.860 grams.

Examination of these figures makes quite clear that the body was still in nitrogen equilibrium, or essentially so, the minus balance being so small as to have little significance. The body-weight was still stationary, and yet during this balance period the average daily intake of nitrogen was only 5.86 grams, corresponding to 36.62 grams of proteid or albuminous food. Further, the average daily fuel value of the food was only 1549 calories, a trifle less than in the preceding period. The average daily output of nitrogen through the urine for this period was 4.92 grams, corresponding to the metabolism of 30.7 grams of proteid food. Hence, the results of this period confirm those of the preceding period and make it quite clear that this subject, with a body-weight of 57.5 kilos, can be maintained in body equilibrium, and in nitrogen equilibrium, on a daily diet containing only 5.8 grams of nitrogen and with a fuel value of about 1600 calories. Under these conditions, as in the last balance period, the daily amount of nitrogen metabolized was very small, averaging only 4.92 grams. Comparison of this figure with the accepted standard of 16 grams of nitrogen makes quite clear the extent of the physiological economy which is attainable by the body, and emphasizes also the extent of the unnecessary and worse than useless labor put upon the body by the prevalent dietetic habits of the majority of mankind.

It is of course understood that the low fuel value which sufficed to keep the writer in body equilibrium would not meet the requirements of a more active life, with greater physical labor. The writer has led a very busy life during the year of this experiment, but it has been mental activity rather than physical, although doubtless he has exercised as much as the ordinary professional worker not accustomed to athletic sports. The results of the experiment, however, make it quite clear that a man of the above body-weight, even though he lead a very active life — not involving great physical labor — can maintain his body in equilibrium indefinitely with an intake of 36 to 40 grams of proteid or albuminous food, and with a total fuel value of about 1600 calories. Further, it is to be

understood that there is no special form of diet involved in the accomplishment of such a result. Scrutiny of the daily diet, tabulated in the two balance periods, will show the character of the food made use of. Personal likes and dislikes must naturally enter into the choice of any diet, and freedom of choice, freedom to follow the dictates of one's appetite, with such regulation as comes from the use of reason and intelligence, are all that is necessary to secure the desired end. Physiological economy in nutrition is easily attainable and does not involve the adoption of vegetarianism. mean, however, temperance and simplicity in diet, coupled with intelligent regulation, which, however, soon becomes a habit and eventually leads to a moderation in diet which fully satisfies all the cravings of appetite as completely as it suffices to maintain the body in equilibrium and in a general condition of health and vigor.

Taking the data recorded above, we may now calculate the nitrogen requirement of the body per kilo of body-weight. With the body-weight placed at 57 kilos and with an average daily elimination of nitrogen for nearly nine months of 5.699 grams, or practically 5.7 grams, it is evident that the nitrogen metabolized per kilo of body-weight in the present instance was exactly 0.1 gram. If we take the lower figure of 5.40 grams of nitrogen, the average daily excretion from April 13 to June 27, we find the nitrogen requirement to be 0.0947 gram per kilo of body-weight. Translating these figures into terms of proteid or albuminous matter, they mean the utilization or metabolism of 0.625 gram of proteid matter daily per kilo of body-weight, under the conditions of life, activity, and general food consumption prevailing throughout this period of nearly nine months with this particular individual.

Whether we are justified in saying that this figure represents the *minimal* proteid requirement of this particular individual is perhaps questionable, since the proteid or nitrogen requirement will of necessity vary somewhat with the amount of non-nitrogenous food consumed. Doubtless, the nitrogen metabolism could be reduced still lower by increas-

ing the intake of non-nitrogenous food, but under the above conditions of life, following a plan of living both congenial and satisfactory, one that fully sufficed to keep the body in equilibrium and with the practice of a general physiological economy, we may say that the metabolism of 0.1 gram of nitrogen per kilo of body-weight was quite sufficient to meet all the requirements of the body. Health, strength, mental and physical vigor have been maintained unimpaired, and there is a growing conviction that in many ways there is a distinct improvement in both the physical and mental condi-Greater freedom from fatigue, greater aptitude for work, greater freedom from minor ailments, have gradually become associated in the writer's mind with this lowered proteid metabolism and general condition of physiological economy. The writer, however, is fully alive to the necessity of caution in the acceptance of one's feelings as a measure of physical or mental condition, but he has been keenly watchful for any and every sign or symptom during the course of these experiments, and is now strongly of the opinion that there is much good to be gained in the adoption of dietetic habits that accord more closely with the true physiological needs of the body. If a man of 57 kilos body-weight can maintain a condition of equilibrium, with continuance of health, strength, and vigor (to say nothing of possible improvement), with a daily consumption of say 40 grams of proteid food and sufficient non-nitrogenous food to yield 2000 calories, why should he load up his system each day with three times this amount of proteid food, with enough more fat and carbohydrate to yield 8000 plus calories?

Finally, the writer in summing up his own experience is inclined to say that while he entered upon this experiment simply with a view to studying the question from a purely scientific and physiological standpoint, he has become so deeply impressed with the great gain to the body by this practice of physiological economy, and his system has become so accustomed to the new level of nutrition that there is no desire to return to the more liberal dietetic habits of former years.

Obviously, it is not wise nor safe to draw too broad deductions from a single individual, nor from a single experiment even though it extends over a long period of time; consequently, we may turn our attention to other individuals with presumably different personality and different habits of life. The writer's colleague, Dr. Lafayette B. Mendel, Professor of Physiological Chemistry in the Sheffield Scientific School, kindly volunteered to become a subject of experiment. With a body-weight of 76 kilos, 32 years of age, and of strong physique, he commenced to modify his diet about the middle of October, 1903, diminishing gradually the amount of proteid food with the results shown in the following tables, where are given, as in the preceding experiment, the amounts of nitrogen in the urine, as a measure of the quantity of proteid metabolized, uric acid, and other factors of interest in this connection.

The collection of data commenced on October 26, 1903. During some weeks the urine of each day was not analyzed by itself, but an aliquot part was taken from the 24 hours' quantity, and at the end of a week the determinations were made on the mixture, thereby giving the average daily composition for the period. With Dr. Mendel, as in the writer's case, there was no prescribing of food, but perfect freedom of choice. The appetite was satisfied each day, but with a gradual diminution of proteid food, especially of meat. Dr. Mendel appeared to accomplish the desired end best by keeping up a liberal allowance of non-nitrogenous food, and the total potential energy of the daily diet was not so greatly diminished as in the writer's case. In other words, he appeared to need more food, but succeeded without great effort in reducing the proteid intake to nearly as low a level as in the preceding experiment. For the period of three months from January 4 to April 3, 1904, the average daily excretion of nitrogen amounted to 6.46 grams, which means the metabolism of 40.37 grams of proteid or albuminous food per day for this quarter of the year.

MENDEL.

	Body-	Urine.					
Date.	weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P3O8.	
1903	kilos	c.c.		grams	gram	grams	
Oct. 26	76.2	1310	1019	10.53			
27	75.0	1650	1016	18.46	0.580	1.90	
28	74.5	915	1023	11.03			
29	74.5	825	1025	11.48	0.629	1.87	
30	74.6	1830	1018	13.17			
31	74.5	1045	1021	12.87			
Nov. 1	74.5	1030	1020	10.38	0.602	1.59	
2		1080	1017	9.59	• • •		
8	74.5	1058	1016	8.86	0.528	1.63	
4	74.5	975	1019	8.66			
5	74.5	1030	1021	8.90	0.514	1.83	
6.		1230	1015	8.11			
7	74.6	1450	1016	8.18			
8	74.0	970	1019	7.91	0.424	1.92	
9	74.0	620	1028	7.72			
· 10	74.0	543	1027	6.60	0.890	1.17	
11	74.0	1160	1016	7.03			
12	74.0	863	1024	5.87	0.422	1.52	
18	74.0	1410	1015	8.12			
14	74.0	1265	1017	8.04	0.494	1.94	
16	74.0	760	1021	6.93			
17	74.0	850	1021	7.84	0.393	1.50	
18	74.0	757	1020	6.84	0.364		
19	74.0	720	1025	7.85	0.456	1.25	
20	74.0	655	1027	7.28	0.474		
21	74.0	985	1021	7.44	0.897		
22	74.5	590	1026	7.65	0.895	1.20	
23	74.5	1100		[] [			
24	74.0	1200	1017				
25		1080		1] 1			
26	74.0	850		7.00	0.410	1.72	
27		935	1020	daily	daily	daily	
28	74.5	870	1021	average	average	average.	
29	74.5	993	1017	J I			
30	74.5	650	1023	1)			
Dec. 1	74.5	960	1018	11			
2		790	1023	7.28	0.480	1.80	
8	74.0	880	1028	11			
4		1200	1016	]]			
		<u> </u>					

# 54 PHYSIOLOGICAL ECONOMY IN NUTRITION

MENDEL.

Date.	Body-	Urine.						
Date.	weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P2O5.		
1903	kilos	c.c.		grams	gram	grams		
Dec. 5	74.0	930	1021	7.28	0 480	1.80		
6		1080	1019	daily av.	daily av.	daily av.		
7	78.5	790	1028	]				
8	73.5	965 ·	1025	11				
9	74.0	1130	1017					
10	78.0	630	1031	7.68	0.438	1.91		
11	73.0	925	1019	11 i				
12	• • •	1395	1014			1		
13	78.0	1010	1018					
14		1080	1020	7.73	• • •			
15	73.0	875	1021	7.88		• • •		
16		625	1027	6.48	]			
17		700	1027	7.60				
18		880	1022	8.89				
19		985	1020	7.68				
20		1075	1019	7.85	ii			
21		528	1032	6.37	0.259	1.10		
22		1455	1017	8.99	1			
23	• • •	920	1021	8.83				
24		725	1025	8.56	1			
25		865	1024	8.51				
26		710	1027	7.54	J			
27		910	1026	1)				
28	• • •	830	1026	li !				
29	• • •	780	1027					
30		670	1033					
81		680		7.64	0.438	1.16		
1904.								
Jan. 1	• • •	550	•••					
2		1030	1022					
8	•••	1020	1019	ן ע				
4		750	1021	5.68	1			
5	• • •	1030	1016	5.81	H			
6	78.0	815	1023	5.77				
7	72.7	980	1019	6.02	0.436	1.41		
8	73.5	1135	1017	5.72	1			
9	72.2	995	1020	6.89				
10		1175	1017	6.84	J			
11	72.0	1050	1017	6.17	0.443	1.48		
L		l						

MENDEL.

	Body-	Urine.						
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>3</sub> .		
1904	kilos	c.c.		grams	gram	grams		
Jan. 12	72.2	1230	1015	6.71	h			
18	71.8	1250	1016	7.43	,			
14	71.8	1140	1015	7.05	0.448	1.43		
15		965	1019	5.99	daily	daily		
16		860	1023	6.71	11	•		
17		1030	1018	5.81	average	average.		
18		780	1028	5.74	lı l			
19		825	1023	6.58				
20		1085	1021	6.70		·		
21		725	1030	6.96	0.465	1.48		
22		815	1023	6.99	11			
28		950	1018	6.27	H			
24	71.6	790	1025	5.98	ן' (			
25	71.8	740	1027	5.83	1			
26		600	1030	5.44	[]			
27		965	1020	6.89				
28		1045	1015	6.33	0.429	1.29		
29		895	1017	6.28				
30	70.6	660	1027	6.53				
31		905	1021	6.79	IJ			
Feb. 1	71.7	695	1025	5.13	h			
2		950	1028	6.84				
3		1210	1019	8.10	11			
4	71.5	985	1020	6.74	0.451	1.40		
5		1155	1020	6.51	] ]			
8	71.1	1035	1019	6.27	11			
7	70.8	760	1025	6.98	[]			
8	70.5	800	1022	6.29	<b> </b>			
9	70.6	1150	1028	7.52	0.448			
10	70.4	770	1022	6.75	0.818			
11	69.2	520	1031	6.71	0.458			
12	69.4	565	1038	8.24	0.390			
13	69.4	560	1030	7.88	0.420			
14	69.2	690	1027	7.99	0.447			
15	69.5	680	1027	7.50	h l			
16		905	1019	6.86				
17	:::	1055	1018	5.68	0.420			
18	l :::	1185	1015	6.11		· · · ·		
19	:::	712	1025	5.72	[] [			
	l <u></u>							

7.4	Body-			Urine.		
Date,	weight.	Volume. 24 hours.	вр. Gr.	Nitrogen.	Uric Acid.	P2O8.
1904	kilos	c.c.		grams	gram	gram
Feb. 20	70.5	1000	1021	7.38	7 0.420	
21	70.6	1235	1014	5.98	daily av.	• • • •
22		900	1018	6.16	1)	
23	70.2	840	1020	5.49	] }	
24		875	1017	5.88	] [	
25	70.5	1450	1018	8.09	0.488	
26	70.2	1485	1015	6.68		
27		1300	1013	5.93	11	
28	69.2	735	1022	5.91	IJ	
29		575	1030	6.21	1	
Mar. 1		975	1019	7.51	i I	
2	70.5	1240	1015	7.29	<b>   </b>	
8		1400	1013	6.68	0.462	
4	70.2	1375	1016	7.34	11	
5	69.9	1100	1017	7.06	11	
6	70.0	960	1020	6.51	Į)	
7	70.5	970	1017	5.53	1	
8	70.9	1220	1015	5.56	11	
9	70.9	1285	1015	5.70	11	
10		1000	1020	6.24	0.413	
11	70.8	1120	1017	5.98	1	
12		1285	1015	6.55		
13	70.4	1110	1015	5.79	J	
14	70.0	690	1024	5.92	)	
15	70.6	1240	1017	7.29	1)	
16	70.8	1450	1016	7.47	11	
17	70.2	780	1022	6.41	0.485	
18		1230	1012	6.57		
19	70.1	780	1027	6.41		
20		950	1020	6.21	IJ	
21	70.7	1005	1020	6.36	h	
22	70.9	1525	1014	6.50	11	
23	70 6	825	1023	6.39	11	
24	70.4	550	1029	6.07	0.527	
25		1070	1018	6.93		
26	70.8	1100	1017	6.40		
27	70.6	1115	1016	5.82	J .	
28	70.2	1185	1015	6.22	) 0000	
29	70.5	1370	1014	6.58	6.389	• • •

MENDEL.

	Do do			Urine.		
Date.	Body- weight,	Volume. 24 hours.	Вр. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>3</sub> .
1903	kilos	c.c.		grams	gram	grams
Mar. 30	70.3	1185	1016	6.26	ا	•
81	70.3	1325	1013	5.96		
Apr. 1		1060	1016	6.30	0.389	
2	70.6	1115	1016	6.42	daily	• • •
3	70.3	1290	1013	6.89	average	
4	70.0	845	1022	6.44	h	
5		1110	1020	6.89	li l	
6		1055	1022	7.53		
7		575	1027	6.42	0.356	1.54
8		650	1031	6.94		
9		795	1026	7.06	]	
10		1230	1020	7.01	}	
11		850	1021	5.61	15 1	
12		1005	1018	6.66		
18		695	1020	5.75		
14	69.6	910	1027	5.79	0.419	
15	70.1	1000	1018	6.42	5	• • •
16	70.9	1590	1016	6.30		
17	70.8	1250	1015	5.25	]]	
18	70.5	985	1020	5.79	5	
19	70.7	1230	1016	5.90	ii I	
20	70.5	1485	1014	5.70		
21	70.1	1125	1023	7.09	0.453	
22	70.3	1665	1013	7.09		•••
23	69.8	985	1023	6.06	[]	
24	69.7	1100	1018	6.07	IJ	
25	69.6	935	1021	5.78	5 1	
26	69.9	1000	1021	6.18	]	
27	70.1	1295	1015	6.06	H	
28	70.0	1425	1013	5.56	0.378	
29	70.2	990.	1022	6.24		•••
30	70.2	1100	1021	7.82		
May 1	70.0	1380	1014	5.96	IJ I	
2	69.8	1050	1016	5.85	Γ, Ι	
8	69.8	700	1022	5.46		
4	69.5	900	1019	6.48		
5	69.6	750	1023	6.52	0.260	• • •
6	69.7	1120	1019	7.26		
7		1010	1020	6.00	)	

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>9</sub> .
1904	kilos	c.c.		grams	gram	gram
May 8		1165	1015	6.01	0.260	
9	69.7	880	1020	5.88	1)	
10	70.0	985	1019	5.22	}}	
11	69.6	1050	1017	6.05	11	
12	69.8	950	1020	6.15	0.408	
18		1060	1020	6.62	daily	
14	69.5	1520	1015	6.47	average.	
15	69.3	1345	1014	5.65	l)	
16	69.0	1230	1015	5.09		
17	68.4	775	1019	5.11	1.	
18	69.2	660	1021	6.06	1)	
19	68.6	905	1018	7.17	] [	
20	69.4	685	1022	6.33	11	
21	69.1	1142	1018	6.78		
22	69.5	1055	1019	5.70	0.825	
23	69.6	1058	1018	5.75		
24	69.8	895	1020	6.39		
25	69.4	900	1018	6.05		
26	69.7	725	1025	6.55		
27	70.0	705	1026	7.86	Ų	
28	71.0	1115	1020	8.23	0.476	
29		1870	1016	7.88	}	• • •
30	69.7	740	1023	7.10	<b> </b> ]	
81		1185	1017	5.98		
June 1		1300	1018	6.86		
2		1420	1014	6.06		
8	69.7	1447	1015	7.03		
4		1388	1016	5.97		
5		1530	1015	6.48	1	
6		870	1023	5.53	1	
7		1010	1013	4.91		
8		815	1019	5.53		
9		865	1019	6.59		
10	69.7	1110	1015	6.86		
11		1410	1017	5.95		
12	69.1	1510	1014	6.07		
13		1100	1016	5.94	1	
14		1090	1018	5.43		
15		1380	1017	6.46		
				<u> </u>		

MENDEL.

Data Body-		Urine.							
Date.	weight.	Volume. 24 hours.	Sp. Gr. Nitrogen.		Uric Acid.	P <sub>2</sub> O <sub>8</sub> .			
1904	kilos	c.c.		grams	gram	grams			
June 16	70.0	1760	1014	6.55					
17		1755	1013	6.63					
18		1075	1026	6.06					
19	70.0	1285	1016	6.55					
20		585	1022	5.44		• • • .			
21		595	1024	6.93					
22		655	1028	7.07					
23		1230	1015	7.45	• • • •				
Daily ave Nov. 10		1001	1020	6.53	0.419	1.46			

From November 10, 1903, to June 23, 1904, a period of about seven months and a half, the average daily excretion of nitrogen through the urine was 6.53 grams. In other words, throughout this long period the average daily amount of proteid matter metabolized was 40.8 grams, only a little more than one-third the amount called for by the Voit standard. Until February, the body-weight gradually fell, but from the early part of February until the end of the experiment the bodyweight remained practically stationary at 70 kilos. Dr. Mendel, however, from the necessities of his daily work in the laboratory was compelled to a much greater degree of physical activity than the subject of the preceding experiment, and consequently required'a larger amount of non-nitrogenous food than the latter. Further, owing to his greater physical activity and the necessary variations in this daily activity, it was not so easy at first to attain equilibrium.

On February 9, a balance experiment of six days was commenced, with a careful comparison of the nitrogen intake and output. In the accompanying tables are shown all of the data. By scrutiny of these it will be seen that Dr. Mendel had adopted essentially a vegetarian diet. During this period of

six days, however, he was not in nitrogen equilibrium, neither was he strictly in body equilibrium, since there was a distinct tendency for the body to fall off in weight. In this connection it may be mentioned that there is always a tendency during a balance experiment of this character for the subject to eat less than he is ordinarily accustomed to, owing to the tediousness of weighing every particle of food consumed. Further, for the same reason, and to avoid excess of chemical work in the analysis of samples of food, he is inclined to limit his diet to a few articles and thereby unconsciously restricts his intake of food, sometimes disastrously so.

#### MENDEL.

### Tuesday, February 9, 1904.

Breakfast. — Bread 83 grams, sugar 20 grams, coffee and milk 210 grams.
 Lunch. — Consommé 150 grams, sweet potato 170 grams, bread 135 grams, tomato 106 grams, coffee and milk 210 grams, sugar 20 grams.
 Dinner. — Bread 75 grams, mashed potato 200 grams, string beans 91 grams, apple pie 282 grams, coffee and milk 210 grams, sugar 20 grams, water 100 grams.

Food.		Grams.	Per cent Nitroger	. Total Nitrogen.
Bread 8	3 + 185 + 75 =	= 243	× 1.86 =	= 3.30 grams.
Sugar 2	0 + 20 + 20 =	= 60	× 0.00 =	= 0.00
Coffee (breakfas	st)	. 210	× 0.10 =	= 0.21
Consommé .		. 150	× 0.38 =	= 0.57
Sweet potato		. 170	× 0.28 =	= 0.48
Tomato		. 106	× 0.19 =	= 0.20
Coffee (lunch)		. 210	× 0.15 =	= 0.32
Potato		. 200	× 0.36 =	= 0.72
String beans		. 91	× 0.26 =	= 0.24
Apple pie		. 282	× 0.49 =	= 1.38
Coffee (dinner) .		. 210	× 0.099 =	= 0.21
	Total nitrogen	in food .		. 7.63 grams.
•				. •

Fuel value of the food . . . . 2297 calories.

# Wednesday, February 10, 1904.

Breakfast. — Bread 37 grams, sugar 20 grams, coffee and milk 210 grams.

Lunch. — Bread 110 grams, sugar 7 grams, milk 250 grams, apple fritters
90 grams.

Dinner. — Bread 37 grams, sugar 21 grams, baked beans 100 grams, cranberry sauce 125 grams, coffee and milk 210 grams, molasses candy 54 grams.

Food.		Grams.	Per cent N	itrogen.	Total Nitrogen.
Bread 37	7 + 110 + 87	= 184	× 1.65	=	8.04 grams.
Sugar 20	0 + 7 + 21	= 48	× 0.00	=	0.00
Coffee (breakfast	t)	210	× 0.12	=	0.25
Milk		250	× 0.58	<b>=</b>	1.87
Apple fritters .		90	× 0.48	<b>i</b> =	0.40
Baked beans		100	× 1.40	=	1.40
Cranberry sauce		125	× 0.04	=	0.05
Coffee (dinner) .		210	× 0.11	. =	0.23
Candy		54	× 0.06	=	0.03
	Total nitro	ren in food			. 6.77 grama

Fuel value of the food . . . 1673 calories.

## Thursday, February 11, 1904.

Breakfast. — Bread 40 grams, sugar 20 grams, coffee and milk 210 grams.

Lunch. — Bread 95 grams, sweet potato 180 grams, sugar 7 grams, milk 250 grams, peach preserve 98 grams.

Dinner. — Bread 90 grams, mashed potato 100 grams, tomato purée 185 grams, baked beans 75 grams, lemon pie 110 grams, coffee and milk 210 grams, sugar 21 grams.

Food.	Grams.	Per cent Nitrogen.	Total Nitrogen.
Bread $40 + 95 + 90$	= 225	× 1.75 =	3.94 grams.
Sugar $20 + 7 + 21$	= 48	× 0.00 =	0.00
Coffee (breakfast)	. 210	× 0.096 =	0.20
Sweet potato	. 130	× 0.81 =	0.40
Milk	. 250	× 0.51 =	1.27

Peach preserve							93	×	0.0	•	=	:	0.08
Potato							100	×	0.36	3	=		0.86
Tomato purée							185	×	0.3	3	=	;	0.45
Baked beans.							75	×	1.30	)	=	:	0.98
Lemon pie .							110	×	0.6	l	=	:	0.67
Coffee (dinner)			•				210	×	0.18	3	=		0.27
	To	tal	ni	tro	ger	in	food						8.62 grams.
							urine						

Fuel value of the food . . . . 1828 calories.

# Friday, February 12, 1904.

Breakfast. — Bread 58 grams, sugar 21 grams, coffee and milk 210 grams.

Lunch. — Bread 120 grams, sugar 21 grams, custard 76 grams, milk 250 grams, coffee and milk 125 grams.

Dinner. — Bread 67.5 grams, sugar 21 grams, mashed potato 150 grams, lima beans 80 grams, coffee and milk 210 grams, apple dumpling 131 grams, molasses candy 27 grams.

Food.		Grams.	Per c	ent Nitrogen.	Total Nitrogen.
Bread . 58+	120 + 67.5 =	245.5	X	1.71 =	4.20 grams.
Sugar . 21 +	-21+21 =	63.0	X	0.00 =	0.00
Coffee (breakfas	st)	210.0	X	0.11 =	0.23
Custard		76.0	X	0.83 =	0.63
Milk		250.0	X	0.48 =	1.20
Coffee (lunch)		125.0	X	0.078 =	0.10
Potato		150.0	X	0.37 =	0.56
Lima beans .		80.0	×	0.90 =	0.72
Coffee (dinner)		210.0	X	0.12 =	0.25
Apple dumpling		131.0	X	0.72 =	0.94
Candy		27.0	X	0.06 =	0.00
	Total nitrogen	in food .			. 8.83 grams.
	Total nitrogen	in urine .			. 8.24

Fuel value of the food . . . . 1929 calories.

# Saturday, February 13, 1904.

Breakfast. — Bread 47.5 grams, sugar 28 grams, coffee and milk 210 grams.

Lunch.—Bread 57 grams, sugar 40 grams, sweet potato 135 grams, quince preserve 78 grams, apple turnovers 118 grams, coffee and milk 310 grams.

Dinner. — Bread 59 grams, mashed potato 175 grams, peas 80 grams, apple pie 141.5 grams, sugar 21 grams, coffee and milk 210 grams.

Food.	Grams.	Per	ent Nitrogen.	Total Nitrogen.
Bread $47.5 + 57 + 59 =$	: 164.0	×	1.64 =	2.69 grams.
Sugar $28 + 40 + 21 =$	89.0	×	0.00 =	0.00
Coffee (breakfast)	210.0	×	0.11 =	0.23
Sweet potato	. 135.0	×	0.37 =	0.50
Quince preserve	. 73.0	×	0.047 =	0.03
Apple turnovers	. 118.0	×	0.96 =	1.18
Coffee (lunch)	. 810.0	×	0.15 =	0.47
Potato	. 175.0	×	0.37 =	0.65
Peas	. 80.0	×	0.96 =	0.77
Apple pie	. 141.5	×	0.43 =	0.61
Coffee (dinner)	. 210.0	×	0.11 =	0.28
Total nitroge	n in food			. 7.31 grams.
Total nitroge	n in urine			. 7.83

Fuel value of the food . . . . 2057 calories.

# Sunday, February 14, 1904.

Breakfast.—Bread 50 grams, sugar 21 grams, banana 92.5 grams, coffee and milk 210 grams.

Lunch. — Bread 108.5 grams, sugar 28 grams, baked potato 165 grams, apple sauce 114 grams, coffee and milk 210 grams.

Dinner. — Bread 63 grams, sugar 28 grams, succotash 75 grams, mashed potato 200 grams, chocolate layer cake 80 grams, ice cream 73 grams, coffee and milk 210 grams.

Food.		Grams.	Per cent N	itrogen.	Total Nitrogen.
Bread . 50+	108.5 + 68 =	221.5	× 1.78	<b>i</b> =	3.87 grams.
Sugar . 21 +	28 + 28 =	77.0	× 0.00	) =	0.00
Banana		92.5	× 0.20	) =	0.19
Coffee (breakfas	t)	210.0	× 0.11	1 =	0.23
Baked potato .		165.0	× 0.4	l =	0.68
Apple sauce .		114.0	× 0.05	29 =	0.08
Coffee (lunch)		210.0	× 0.10	) =	0.21
Succotash		75.0	× 0.5	7 =	0.43
Mashed potato		200.0	× 0.87	<i>I</i> =	0.74
Chocolate cake		80.0	× 0.78	<b>i</b> =	0.60
Ice cream		73.0	× 0.58	3 =	0.42
Coffee (dinner)		210.0	× 0.1	= ا	0.23
	Total nitrogen	in food			. 7.68 grams
	Total nitrogen	in urine			. 7.99

Fuel value of the food . . . . 2065 calories.

### NITROGEN BALANCE. - Mendel.

	Nitrogen	Outy	at.
	Taken in. Nits	rogen in Urine.	Weight of Faces (dry).
Feb. 9	7.68 grams.	7.52 grams	• •••
10	6.77	6.75	• • •
11	8.62	6.71	•••
12	8.83	8.24	53 contain 5.84% N = 6.248 grm. N.
18	7.31	7.83	54) Contain 5.52 % N = 0.245 grin. N.
14	7.63	7.99	39.2  contain  6.72%  N = 2.634
	-		146.2 8.882 grm. N.
	46.79	45.04 +	8.882 grams nitrogen.
	46.79 grams 1	T. 58.9	2 grams nitrogen.
	Nitrogen bal	ance for six	days = -7.18 grams.
	Nitrogen bal	lance per day	= -1.19  grams.
		Averag	ge Intake.
	Calories per	day	1975.
	Nitrogen per	rday	7.88 grams.

In this balance period of six days the average daily intake of nitrogen was 7.83 grams, coupled with an average fuel value of the food per day of 1975 calories. This latter value was obviously too small for a man of Dr. Mendel's body-weight, and doing the amount of physical work he was called upon to perform. Being on one's feet in a laboratory six to eight hours a day, in addition to the ordinary activity of a vigorous man leading a strenuous life, necessitates the utilization and oxidation of considerable food material. The average daily output of nitrogen through the urine amounted to 7.50 grams, considerably above the average daily excretion for the seven months' period. Still, under these conditions there was a minus balance of 7.13 grams of nitrogen for the six days' period, indicating that the body was drawing upon its stock of proteid material to the extent of 1.19 grams of nitrogen per day. This does not necessarily mean that the body had need of that additional amount of proteid matter each day, but rather that the amount of total energy required was beyond the potential energy supplied by the food. There not being sufficient non-nitrogenous food at hand, the body was compelled to draw upon its own resources, and in so doing utilized some of its tissue proteid. This is made quite clear by the results of the second balance period shortly to be described.

It is evident, however, that while the body was not in nitrogen equilibrium for this particular period of six days, there must have been a general condition of both body and nitrogen equilibrium, otherwise the body-weight would not have remained practically stationary for so long a period as from February 7 to June 20.

Commencing May 18, a second nitrogen balance was attempted, in which, as in the preceding case, there was a careful comparison of income and output for seven days. There was as before a free choice of food, but it was essentially vegetable in character. A greater variety of foods was taken, however, and an effort was made to have the non-nitrogenous food somewhat more liberal in amount, though in as close harmony as possible with the desires of the appetite.

# Wednesday, May 18, 1904.

- Breakfast.—Banana 92 grams, bread rolls 28 grams, cream 50 grams, coffee 150 grams, sugar 21 grams.
- Lunch. Bread 66 grams, soup 150 grams, farina 154 grams, sweet potato 123 grams, beans 70 grams, syrup 50 grams, coffee 150 grams, cream 50 grams, sugar 14 grams.
- Dinner. Bread 42 grams, consommé 100 grams, spinach 100 grams, mashed potato 250 grams, apple pie 97 grams, coffee 150 grams, cream 50 grams, sugar 21 grams.

Banana
Cream $50 \times 0.46 = 0.230$
Coffee 150 × 0.068 = 0.099
8ugar
Bread 66 × 1.60 = 1.056
Soup 150 $\times$ 0.41 = 0.615
Farina 154 $\times$ 1.09 = 1.678
Sweet Potato
Beans 70 $\times$ 0.84 = 0.288
Syrup 50 $\times$ 0.024 = 0.012
Coffee 150 $\times$ 0.057 = 0.086
Cream 50 $\times$ 0.46 = 0.230
Sugar 14 $\times$ 0.00 = 0.000
Bread 42 $\times$ 1.80 = 0.758
Consommé 100 × 0.38 = 0.380
Spinach 100 $\times$ 0.58 = 0.530
Mashed potato $\dots$ 250 $\times$ 0.88 = 0.950
Apple pie 97 $\times$ 0.43 = 0.417
Coffee 150 × 0.06 = 0.090
Cream 50 $\times$ 0.46 = 0.280
Sugar 21 $\times$ 0.00 = 0.000
Total nitrogen in food 8.668 grams.
Total nitrogen in urine 6.060

Fuel value of the food . . . . 2859 calories.

## Thursday, May 19, 1904

- Breakfast. Banana 102 grams, bread rolls 50 grams, coffee 150 grams, cream 50 grams, sugar 21 grams.
- Lunch. Bread 57 grams, egg omelette 20 grams, hominy 137 grams, syrup 68 grams, potatoes 128 grams, coffee 100 grams, sugar 21 grams, cream 50 grams.
- Dinner. Tomato pureé 200 grams, bread 24 grams, fried sweet potato 100 grams, spinach 70 grams, Indian meal 100 grams, syrup 25 grams, coffee 100 grams, sugar 21 grams, cream 40 grams.

Food.							Grams.	Per	cent Nitro	gen.	Total Nitrogen.
Banana							102	×	0.23	=	0.235 grams.
Bread rolls							50	X	1.54	=	0.770
Coffee							150	×	0.06	=	0.090
Cream						•	50	X	0.47	=	0.235
Sugar							21	×	0.00	=	0.000
Bread							57	×	1.60	=	0.912
Egg Omelette .							20	×	1.58	=	0.816
Hominy							137	×	0.20	=	0.274
Syrup							68	×	0.024	=	0.016
Potatoes							128	×	0.49	=	0.627
Coffee							100	×	0.06	=	0.060
Cream							50	X	0.47	=	0.235
Sugar					•		21	×	0.00	=	0.000
Tomato purée .							200	×	0.58	=	1.060
Bread					•		24	×	1.74	=	0.418
Sweet potato .							100	×	0.88	=	0.380
Spinach							70	×	0.56	=	0.392
Indian meal .				٠.			100	X	0.20	=	0.200
Syrup							25	X	0.024	=	0.006
Coffee							100	X	0.06	=	0.060
Sugar							21	×	0 00	=	0.000
Cream							40	X	0.47	=	0.188
	•	Γot	al 1	itr	ogo	en	in food				. 6.474 grama.
	7	Γot	al 1	itr	oge	en i	in uripe				. 7.170

Fuel value of the food . . . . 2072 calories.

# Friday, May 20, 1904.

- Breakfast. Sliced orange 140 grams, coffee 100 grams, cream 80 grams, sugar 21 grams.
- Lunch.—Bread 28 grams, mashed potato 250 grams, lima beans 40 grams, coffee 100 grams, sogar 21 grams, cream 80 grams, fried hominy 115 grams, syrup 48 grams.
- Dinner.—Bread 19 grams, consommé I50 grams, string beans 140 grams, mashed potato 250 grams, rice croquette 93 grams, cranberry jam 95 grams, coffee 100 grams, sugar 21 grams, cream 30 grams, syrup 25 grams.

Food.	Grams. Per cent Nitrogen.	Total Nitrogen.
Sliced orange .	140 × 0.20 =	0.280
Coffee	100 × 0.06 =	0.060
Cream	30 × 0.44 =	0.132
Sugar	$\dots \dots 21 \times 0.00 =$	0.000
Bread	28 × 1.71 =	0.479
Mashed potato	250 × 0.30 =	0.750
Lima beans .		0.304
Coffee	100 × 0.06 =	0.060
Sugar	21 × 0.00 =	0.000
Cream	30 × 0.44 =	0.132
Fried hominy .	$.$ $.$ $.$ $.$ $.$ 115 $\times$ 0.57 $=$	0.656
Syrup	. $.$ $.$ $.$ $.$ $.$ $.$ $.$ $.$ $.$	0.012
Bread	19 × 1.97 =	0.374
Consommé	150 × 0.59 =	0.885
String beans .	140 × 0.86 =	0.504
Mashed potato	250 × 0.34 =	0.850
Rice croquettes	93 × 1.06 =	0.986
Cranberry jam	95 × 0.03 =	0.029
Coffee	100 × 0.06 =	0.060
Sugar	$.$ $.$ $.$ $.$ 21 $\times$ 0.00 $=$	0.000
Cream	30 × 0.44 =	0.132
Syrup	$ 25 \times 0.024 =$	0.006
	otal nitrogen in food	. 6.691 grams.
	otal nitrogen in urine	. 6.380

Fuel value of the food . . . . 1915 calories.

## Saturday, May 21, 1904.

- Breakfast.—Banana 153 grams, coffee 150 grams, sugar 21 grams, cream 30 grams.
- Lunch. Bread 25 grams, potato croquette 229 grams, Indian meal 109 grams, tomato 123 grams, syrup 48 grams, coffee 100 grams, sugar 14 grams, cream 20 grams.
- Dinner. Bread 31 grams, bean soup 100 grams, fried potato 200 grams, bacon 5 grams, lettuce-orange salad 47 grams, prunes 137 grams, coffee 100 grams, sugar 21 grams, cream 25 grams, banana 255 grams.

Food.		Grams.	Per cent Nitro	gen.	Total Nitrogen.
Banana		153	× 0.23	=	0.352 grams.
Coffee		150	× 0.06	=	0.090
Sugar		21	× 0.00	=	0.000
Cream		30	× 0.48	=	0.129
Bread		25	× 1.82	=	0.455
Potato croquette		229	× 0.71	=	1.626
Indian meal .		109	× 1.09	=	1.188
Tomato		123	× 0.17	=	0.209
Syrup		48	× 0.024	=	0.012
Coffee		100	× 0.06	=	0.060
Sugar		14	× 0.00	=	0.000
Cream		20	× 0.48	=	0.086
Bread		81	× 1.62	=	0.502
Bean soup		100	× 1.21	=	1.210
Fried potato .		200	× 0.60	=	1.200
Bacon		5	× 8.05	=	0.153
Lettuce-orange s	alad	47	× 0.21	=	0.099
Prunes		187	× 0.16	=`	0.219
Coffee		100	× 0.06	=	0.060
Sugar		21	× 0.00	=	0.000
Cream		25	$\times$ 0.43	=	0.108
Banana		255	× 0.23	=	0.587
	Total nitrogen	in food	<b>.</b>		8.345 grams.
	Total nitrogen				6.780

Fuel value of the food . . . . 2485 calories.

## Sunday, May 22, 1904.

- Breakfast. Banana 220 grams, orange 60 grams, coffee 100 grams, sugar 21 grams, cream 25 grams.
- Lunch. Bread 35 grams, potato 300 grams, fried rice 160 grams, syrup 63 grams, ice cream 84 grams, coffee 100 grams, sugar 14 grams.
- Dinner. Cream of celery soup 100 grams, bread 21 grams, mashed potato 250 grams, spinach 40 grams, French fried potato 100 grams, strawberry short-cake 120 grams.

Food.		Grams.	Per cent Nitr	ogen.	Total Nitrogen.
Banana		. 220	× 0.23	=	0.506 grams.
Orange		. 60	× 0.20	=	0.120
Coffee		. 100	× 0.06	=	0.060
Sugar		. 21	× 0.00	=	0.000
Cream		. 25	× 0.45	=	0.118
Bread		. 85	× 1.57	=	0.550
Potato		. 300	× 0.30	=	0.900
Fried rice		. 160	× 0.75	=	1.200
Syrup		. 63	× 0.024	=	0.015
Ice cream		. 84	× 0.53	=	0.445
Coffee		. 100	× 0.06	=	0.060
Sugar		. 14	× 0.00	=	0.000
Cream of celery so	o <b>up</b>	. 100	× 0.33	=	0.330
Bread		. 21	× 1.91	=	0.401
Mashed potato		. 250	× 0.37	=	0.925
Spinach		. 40	× 0.55	=	0.220
French fried potat	to	. 100	× 0.57	=	0.570
Strawberry short-	cake	. 120	× 0.50	=	0.600
•	Total nitroge	en in food			7.015 grams.
•	Total nitroge	en in urine			5.700

Fuel value of the food . . . . 2321 calories.

## Monday, May 23, 1904.

Breakfast. — Banana 229 grams, coffee 125 grams, sugar 21 grams, cream 25 grams.

Lunch. — Bread 58 grams, apple sauce 125 grams, scrambled egg 15 grams, consommé 75 grams, fried potato 170 grams, rice croquette 197 grams, syrup 68 grams, coffee 100 grams, sugar 21 grams, cream 30 grams.

Dinner. — Bread 72.5 grams, vegetable soup 100 grams, potato croquette 198 grams, bacon 7 grams, string beans 120 grams, water ice 77 grams, coffee 100 grams, cream 30 grams, sugar 14 grams, banana 270 grams.

Food.		Grama.	Per cent Nitrogen.	Total Nitrogen.
Banana		229 . ×	0.23 =	0.527 grams.
Coffee		. 125 ×	0.06 =	0.075
Sugar		21 ×	0.00 =	0.000
Cream		. <b>2</b> 5 ×	0.45 =	0.113
Bread		. 58 ×	1.63 =	0.945
Apple sauce		125 ×	0.02 =	0.025
Scrambled egg .	<i>.</i>	15 🗙	2.07 =	0.311
Consommé		. 75 ×	0.65 =	0.488
Fried potato		. 170 🗙	0.60 =	1.020
Rice croquette .		. 197 🗙	0.61 =	1.202
Syrup		. 68 🗙	0.024 =	0.016
Coffee		100 ×	0.0წ ==	0.060
Sugar		. 21 ×	0.00 =	0.000
Cream		. 80 🗙	0.45 =	0.135
Bread		. 72.5 ×	1.75 =	1.269
Soup		. 100 🗶	0.70 =	0.700
Potato croquette .		. 198 🗶	0.77 =	1.525
Bacon		. 7 ×	8.28 =	0.230
String beans		. 120 ×	0.22 =	0.264
Water ice		. 77 ×	= 800.0	0.005
Coffee		. 100 🗙	0.06 =	0.060
Cream		. 80 🗙	0.45 =	0.185
Sugar		. 14 🗙	0.00 =	0.000
Banana		. 270 ×	0.23 =	0 621
•	Total nitrogen	in food		9.726 grams.
•	Total nitrogen	in urine .		. 5.750

Fuel value of the food . . . . 2756 calories.

## Tuesday, May 24, 1904.

- Breakfast. Orange 100 grams, rolls 37 grams, rice 50 grams, syrup 25 grams, coffee 125 grams, sugar 21 grams, cream 50 grams.
- Lunch. Bread 77 grams, cream of celery soup 125 grams, mashed potato 270 grams, tomato sauce 50 grams, farina croquette 191 grams, syrup 78 grams, coffee 75 grams, cream 20 grams, sugar 7 grams.
- Dinner. Bread 49 grams, tomato soup 200 grams, French fried potato 200 grams, spinach 50 grams, farina croquette 276 grams, syrup 100 grams, coffee 100 grams, sugar 14 grams, cream 50 grams.

Food.		Grams.	Per cent Nitrogen.	Total Nitrogen.
Orange		. 100	× 0.20 =	0.200 grams.
Bread rolls		. 37	× 1.64 =	0.607
Rice		. 50	× 0.36 =	0.180
Syrup		. 25	× 0.024 =	0.008
Coffee		. 125	× 006 =	0.075
Sugar		. 21	× 0.00 =	0.000
Cream		. 50	× 0.45 =	0.225
Bread	<b>.</b>	. 77	× 1.66 =	1.278
Celery soup		. 125	× 0.48 =	0.600
Mashed potato		. 270	× 0.26 =	0.702
Tomato sauce .		. 50	× 0.23 =	0.115
Farina croquette		. 191	× 0.74 =	1.418
Syrup		. 78	× 0.024 =	0.019
Coffee		. 75	× 0.06 =	0.045
Cream		. 20	× 0.45 =	0.090
Sugar		. 7	× 0.00 =	0.000
Bread		. 49	× 1.82 =	0.892
Tomato soup .		. 200	× 0.19 =	0.880
French fried pota	ito	. 200	× 0.46 =	0.920
Spinach		. 60	X 0.54 =	0.270
Farina croquette		. 276	X 0.76 =	2.098
Syrup		. 100	$\times$ 0.024 =	0.024
Coffee		. 100	× 0.06 ==	0.060
Sugar		. 14	× 0.00 =	0.000
Cream		. 50	× 0.45 =	0.225
	Total nitroge	en in food		. 10.424 grams.
	Total nitrog	en in urine		. 6.890

Fuel value of the food . . . . 3229 calories.

# 74 PHYSIOLOGICAL ECONOMY IN NUTRITION

### NITROGEN BALANCE. - Mendel.

	Nitrogen	Out	out.
	Taken in. Nitro	gen in Urine.	Weight of Faces (dry).
May 18	8.668 grams.	6.06 grams.	14 grams.
19	6.474	7.17	39
20	6.691	6.33	<u>80</u>
21	8.845	6.78	88 contain $6.06\%$ N = $5.08$ grm. N.
22	7.015	5.70	••
23	9.726	5.75	88
24	10.424	6.89	<u> 57</u>
			95 contain 5.76% $N = 5.47$ grm. N.
			10.50 grm. N.
	57.843	44.18 +	10.50 grams nitrogen.
	57.843 grams N.	54.0	38 grams nitrogen.

Nitrogen balance for seven days = +2.663 grams. Nitrogen balance per day = +0.880 gram.

#### Average Intake.

Calories per day . . . . . . . . . . . . . . . . 2448.

Nitrogen per day . . . . . . . 8.192 grams

In this period of seven days the average daily intake of nitrogen was 8.192 grams, or only 0.36 gram per day more than in the first balance period, while the average fuel value of the food amounted to 2448 calories per day. Yet the average daily output of nitrogen through the urine for this period was 6.31 grams, or 1.2 grams per day less than in the first balance experiment. Further, under the conditions of this balance experiment, the body was laying up 0.380 gram of nitrogen per day, i. e., showing a plus balance of 2.66 grams of nitrogen for the seven days' period. Again, it is to be noted that the average daily amount of nitrogen metabolized, 6.31 grams, was 0.22 gram less than the average daily nitrogen excretion for the entire seven months' period, 6.53 grams. Evidently, this subject was quite able to maintain nitrogen equilibrium with a metabolism of only 6.31 grams of nitrogen per day, on a daily diet having a fuel value of about 2400 calories. Indeed, taking into account the amount of the plus nitrogen balance, it is evident that the daily food was somewhat in excess of the real requirements of the body, under the then existing conditions of body-weight and bodily activity.

Again, we would call attention to the thorough utilization of the food in this experiment, emphasizing at the same time the voluminous character of the diet, together with its largely vegetable nature. The contrast between the diet made use of by Dr. Mendel and that used by the subject of the first experiment is quite striking, since the latter employed a much more concentrated diet with an average fuel value of only 1600 calories. Yet with a total intake of 57.343 grams of nitrogen for the seven days of Dr. Mendel's balance period, 10.5 grams only passed out through the rectum, or 18.3 per cent, while in the second nitrogen balance of the first subject, with the more concentrated diet, 17.1 per cent of the total ingested nitrogen appeared in the fæces. In view of the great divergence in the character and volume of the intake, it is rather remarkable there should be so little difference in the relative utilization of the two diets.

Finally, taking the average daily excretion of nitrogen through the kidneys from November 10 to June 23, as a measure of the nitrogen metabolized daily, viz., 6.53 grams, and taking the body-weight at 70 kilos, it is plain to see that the nitrogen metabolized per kilo of body-weight throughout this experiment was 0.093 gram, closely similar to the result obtained with the first subject. In other words, both of these subjects, though widely different in body-weight, under different degrees of physical activity, and living on different forms of diet, seemingly required for the maintenance of equilibrium essentially the same amount of nitrogen per kilo of body-weight; viz., with the first subject 0.0947 gram, if we take the lower figure of the last two months, and 0.093 gram with the second subject.

Regarding the fuel value of the daily food, Dr. Mendel with a body-weight of 70 kilos, during the second balance period, apparently utilized on an average 34.9 calories per kilo of body-weight daily, while the first subject, of 57 kilos body-weight, made use of only 28 calories per kilo. The fuel value of the daily food must, however, as is well known, vary greatly with differing degrees of physical activity, from which arises the necessity for corresponding variation in the amounts of non-nitrogenous foods ingested.

Dr. Frank P. Underhill, instructor in physiological chemistry in the Sheffield Scientific School, is another subject of experiment who volunteered to study on himself the effects of a lowered proteid intake. Prior to the experiment he was in the habit of eliminating from 16 to 16.5 grams of nitrogen per day through the kidneys, representing the usual 105 grams of proteid food metabolized.

Dr. Underhill, at the beginning of the experiment, July 1, 1903, was twenty-six years of age and weighed 67.6 kilos. For a period of two weeks, from July 14 to August 2, he gradually reduced the intake of proteid food as well as the total amount of food consumed, doing this in part by diminishing the quantity eaten at breakfast, and in smaller meas-

ure at the two other meals of the day. During this period of two weeks, the urine and fæces were analyzed with the results shown in the accompanying table. Regarding the extent of proteid metabolism, it will be seen that the nitrogen content of the urine fell from 14.28 grams per day down to 5 to 6 grams per day. Having reached this level, the subject maintained it throughout the summer of 1903, occasional analyses being made to demonstrate the level of nitrogen excretion.

			Urine.		Fæces.				
Date.	Body- weight.	Volume.	Sp. Gr.	Nitrogen.	Moist.	loist. Dry.		Total Nitrogen.	
1903.		0.0.		grams.	grams.	grams.	grams.	grams.	
July 14	67.6	1800	1018	14.28	156.0	30.5	1.59	15.87	
15	67.6	1095	1020	11.72	70.0	17.0	0.80	12.52	
16	67.1	860	1021	11.72	182.5	44.5	2.12	18.84	
17	66.9	675	1022	9.39	134.0	43.0	2.05	11.44	
18	66.3	865	1021	10.45	57.5	20.0	0.95	11.40	
19	65.7	785	1021	10.34	170.0	41.0	1.95	12.29	
20	65.7	740	1024	10.84	76.0	35.0	1.66	12.00	
21	65.7	910	1017	11.61	96.0	32.0	1.52	13.18	
22	65.7	900	1018	9.50	56.0	81.0	1.75	11.25	
23	65.8	600	1017	8.65	41.0	14.0	0.79	9.35	
24	65.0	640	1018	6.65	151.0	88.0	1.86	8.51	
25	65.8	690	1012	6.01	86.0	29.0	1.68	7.64	
26	65.3	410	1028	6.65	57.0	19.0	1.06	7.71	
27	65.7	530	1017	6.75	202.0	83.0	1.86	8.61	
28	65.7	610	1013	5.49	155.0	26.0	1.46	6.95	
29	66.4	620	1017	5.96	121.0	26.0	1.46	7.42	
30	66.8	700	1016	5.80	233.0	48.0	2.71	8.51	
31	<b>65</b> .8	1265	1010	6.70	211.0	86.0	2.03	8.73	
Aug. 1	65.3	1240	1009	6.23	172.0	23.0	1.29	7.52	
2	65 3	760	1016	6.75	214.0	27.0	1.42	8.17	
11	65.8	500	1017	6.44	152.0	22.0	1.34	7.78	
12	65.3	405	1024	6.86	70.0	15.0	0.91	7.77	
13	65.3	540	1019	6.23	90.0	10.0	0.61	6.84	

In connection with the accompanying table of results, attention is called to the composition of the fæces with special reference to their content of nitrogen. The point to be emphasized is the gradual increase in the percentage content of fæcal nitrogen, associated with the changed diet and the general diminution of food intake. This is well illustrated by the following figures, giving the percentage of nitrogen in the dry fæces for the three periods of July 14-21, July 22-30, and August 11-13:

July 14-21, 4.77 per cent Nitrogen July 22-30, 5.65 " " " Aug. 11-13, 6.11 " " "

The diminished amount of fæcal discharge means naturally not only a smaller intake of food, but implies also a greater utilization of the food ingested, and as a result the increased percentage of nitrogen in the discharges shows itself because of the relatively larger preponderance of nitrogenous secretions from the intestinal tract.

The latter part of September, 1903, Dr. Underhill attempted to return to his original methods of living, but found difficulty in consuming the daily quantities of food he had formerly been in the habit of taking. From October 11 to October 25, 1903, however, he raised the consumption of proteid food to such a degree that the nitrogen excretion through the urine averaged from 10 to 12 grams per day. After this date he fell back to the lower proteid intake, and from that period to June 23, 1904, he maintained a low level of proteid metabolism without detriment to his bodily vigor, and, as he believes, with a distinct betterment.

The following tables of results extending from October 11, 1903, to June 23, 1904, show the data collected.

2.4	Body- weight.	Urine.						
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>8</sub> .		
1903.	kilos	0.0.		grams	gram	grams		
Oct. 11	65.4	1800	1015	10.87	0.611	1.72		
12		700	1022	9 07				
13		1050	1022	12.85	0.820	2.15		
14		1400	1017	12.01				
15		870	1020	11.48	0.671	1.76		
16		1165	1018	12.09	• • • •			
17	• • •	840	1020	11.24				
18		1150	1017	10.00	0.618	1.77		
19		678	1022	7.79	• • •			
20		1165	1018	10.76	0.603	1.60		
21		1460	1017	11.91	• • •	• • •		
22		950	1019	10.71	0.648	1.78		
28		1165	1017	12.82	• • •	• • •		
24	65.4	850	1025	12.49	• • •	• • •		
25		850	1022	11.07	0.645	1.76		
26		1025	1018					
27	• • • •	775	1018	11				
28	• • • •	1140	1015	8.81	0.465	1.18		
29	•••	765	1020	daily	daily	daily		
80	• • • •	860	1019	average	average	average		
31		1150	1015	ן י	Ū	J		
Nov. 1		750	1020	l)				
2	• • •	675	1022					
8	• • •	955	1014	-	0.400			
4	• • • •	1270	1012	} 7.91	0.469	1.80		
5	• • •	885	1015	H				
6	• • • •	770	1020					
7	• • •	860	1021	ן י				
8	• • • •	775	1020					
9	• • • •	890	1018					
10	• • • •	1070	1018	]]				
11	• • • •	755	1020	7.86	0.537	1.40		
12	• • • •	950	1028	[]				
18	• • •	1100	1017	11				
14	• • •	980	1015	11				
15	050	680	1020	1,				
16	65.0	700 1000	1020					
17	• • •		1015	7.82	0.467	1.49		
18 19	• • •	940 770	1018 1028	11 1				
18	•••	110	1020	<u>'</u>				

UNDERHILL.

	Rody.			Urine.		
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P2O8.
1903.	kilos	c.c.		grams	gram	grams
Nov. 20		770	1025	1)		
21		790	1020	7.82	0.467	1.49
22		770	1021	daily av.	daily av.	daily av.
23		590	1025	13		
27	• • • •	710	1024	8.55	0.476	1.67
28		1080	1017		0.210	2.01
29		760	1019	l)		
80	• • •	870	1018	1)		
Dec. 1		860	1017	11		
2		1120	1015	11 1		
3		1450	1009	7.72	0.506	1.43
4		720	1019	11		
5		720	1019	]] :		
6		510	1027	ן ין		
7	65.1	700	1018	l)		
8	• • •	650	1018	11		
9		860	1018	11		
10		975	1020	7.36	0.474	1.44
11		800	1021	11		
12		880	1018	]]		
13		750	1019	J		
14		860	1018	l)		
15		870	1019	11		
16		880	1015	11 .		
17		820		8.11	0.497	1.85
18		760	1021			
19		860	1017	]] ]		
20		520	1023	j i		
21		1200	1014	7.18		1.05
22	65.0	510	1025	15	•••	2.00
23		650		lı l		
24		670				
25		500		6.70	0.356	0.97
26		550		[]		
27		850		D		
28	65.2	930		h		
29		1200		8.70	0.626	1.21
30		750		0.70	0.020	1,21
81		1280		ן ען		

UNDERHILL.

	Body-	Urine.							
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P3O3.			
1904	kilos	c.c.		grams	gram	grams.			
Jan. 1		750		11					
2		750		8.7	0.626	1.21			
8		870		daily av.	daily av.	daily av.			
4		1230				uu, u			
5		720	1023	lì l					
6		960	1020	11					
7		760	1022	8.23	0.580	1.24			
8		1000	1017	6.20	0.000	1.22			
9		720	1025	11	1				
10	65.1	730	1021	J)					
11		700	1016						
12	• • • •	650	1020	11					
13		700	1024	11					
14	• • •	660	1025	7.74	0.618	1.20			
15		610	1025	11					
16	• • •	710	1024						
17		660	1022	J) . [					
18		770	1016						
19	64.8	700	1022	11					
20	• • •	1130	1018	11					
21		830	1020	7.80	0.705	1.14			
22	• • • •	600	1019						
28	• • • •	480	1021	11					
24		750	1019	[]					
25		680	1020	1)					
26	• • • •	800	1018	11					
27	1	800	1020			<b>.</b>			
28	1 :::	700	1020	6.85	0.500	1.07			
29	64.4	1010	1014						
30		980	1018	11					
81		820	1015	l,					
Feb. 1		990	1015	11					
2	1 :::	660	1020	11					
8	64.4	1060	1015	,,,,	0.585	1.24			
4		980	1017	7.98	U.080	1.24			
5	• • •	970	1014	11					
6		790	1019 1015						
7	1	1120	1015	8.11					
8 9	64.4	715 1225	1021	8.11 7.49	0.450				
	V								

UNDERHILL.

	Rode			Urine.		
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric <b>Acid</b> .	P <sub>3</sub> O <sub>5</sub> .
1904.	kilos	c.c.		grams	gram	grams
Feb. 10	64.4	770	1018	6.14	0.505	
11	64.5	715	1022	8.02	0.568	
12		850	1021	8.82	0.539	
13	• • • •	490	1028	7.17	0.505	
14	64.4	795	1020	7.73	0.543	
15		780	1021	h		
16		920	1023	11 1		
17		660	1021	[]	'	
18	• • •	640	1027	8.27	0.558	• • •
19		730	1023	daily	daily	
20	• • •	840	1027	average	average	
21		700	1018	ון		
22		440	1025	1)		
23	• • •	600	1023	11		
24	• • •	750	1022			
25		880	1017	7.47	0.556	• • •
26	64.0	870	1021	[ ]		
27	• • •	910	1015			
28		950	1014	ן יין		1
29	• • •	600	1021	1)		
Mar. 1	• • •	680	1021			
2	• • •	1120	1011 1021		0.500	
8	• • •	640		7.15	0.526	• • • •
4	• • •	1080	1016	11 1		
5	• • •	1220	1034		1	
6	• • • •	890	1015 1014	ן ן		
7	• • •	1000 650	1014	[] [		
8	• • •	750	1020			
9	• • • •	750 850	1020	8.18	0.682	
10	• • •	950	1018	0.10	U.002	• • • •
11 12	•••	1000	1014		1	
13	64.2	860	1010			
13 14	U-2.Z	840	1019	1,	'	
14		920	1020			
16		750	1020		'	
17		820	1020	7.88	0.540	
18		1220	1012		0.010	'''
19		920	1015	]]		
20		700	1022			
20				<u> </u>		

Date.	Body-	Urine.							
Date.	Body- weight.	Volume 24 hours.	Sp. Gт.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>5</sub> .			
1904.	kilos	c.c.		grams	gram	grams			
Mar. 21		900	1015	h i	_				
22		980	1018	11					
23		990	1014						
24		750	1020	8.04	0.733				
25		820	1017	daily	daily				
26		700	1021	average	average				
27		910	1020						
28		930	1017						
80		900	1020						
31		450		1					
Apr. 1		500		11 1					
2		420		{{					
8		600		11 1					
4		950		11					
5.		930		7.87	0.580	1.44			
6		980		11 1		daily			
7		600		11 1		average			
8		980							
9		800							
10		810		]] [					
11	65.2	940	1017	l) l	)				
12		710	1024	9.46					
18		750	1024	IJ					
14		740	1020	7.55	0.496				
15		590	1024	6.51					
16	·	900	1016	7.51					
17		1290	1009	5.11	)				
18	65.2	590	1017	5.96	ի				
19		630	1024	6.39	ļ .				
20		670	1022	7.48					
21		900	1021	8.10	0.527				
22		980	1017	7.00		- ' •			
28		850	1019	7.60					
24		640	1023	7.33	J				
25	65.0	600	1021	8.23	1				
26		610	1014	5.82					
27		600	1023	7.27	0.490				
28		970	1015	7.05	0.930	• • •			
29		795	1017 -	5.91					
80	• • •	700	1021	6.80	J				
		L	L	1					

# 84 PHYSIOLOGICAL ECONOMY IN NUTRITION

### UNDERHILL.

	Body-	Urine.							
Date.	weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Une Acid.	P <sub>3</sub> O <sub>8</sub> .			
1904	kilos	c.c.		grams	gram	grams			
May 1		795	1016	5.72	0.490				
2	65.2	860	1017	6.50	h	:			
3		1120	1016	6.92					
4		1090	1015	6.54					
5		580	1021	5.92	0.344				
6		700	1020	7.29	daily				
7		895	1017	8.97	average				
8		800	1016	7.30					
9	65.2	895	1019	6.28	lí l				
10		850	1019	6.27	[] [				
11		960	1019	7.26					
12		690	1022	6.29	0.416				
18		990	1017	7.96					
14		690	1022	7.21					
15		900	1015	6.15					
16	65.1	945	1014	5.10	K				
17		1090	1016	6.15					
18		620	1020	5.21	1				
19		1110	1015	6.53					
20		895	1016	6.12	0.364				
21		810	1019	6.95					
22		1110	1016	8.72	l i				
23	65.0	685	1020	7.07					
24		560	1021	6.78					
26		1090	1010	6.02					
26		610	1017	5.97					
27		790	1016	6.88					
28		1100	1014	7.65	0.420				
29		650	1021	6.12		•••			
30	65.0	630	1021	5.21					
81		660	1018	5.07					
June 1	65.1	920	1014	5.96	, , ,				
2	65.1	800	1013	5.81	:::	• • •			
3	65.0	950	1014	7.30	:::				
4	05.0	790	1015	6.78	[]				
5		780	1020	8.28					
6	65.1	890	1015	6.89	:::	• • •			
7	00.1	720	1017	5.87	[	• • •			
8		950	1015	5.98	l				
9		1060	1014	4.96	:::	• • •			
	L			1	L				

UNDERHILL.

	Body-	Urine.								
		Volume. 24 hours. Sp. Gr. Nitr		Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>5</sub> .				
1904	kilos	0.0.		grams	gram	grams				
June 10		940	1017	5.92						
11		1050	1019	7.62						
12		1000	1017	6.66						
18		890	1021	7.26						
14		1840	1016	6.91						
15		1190	1015	6.86						
16		720	1025	7.99						
17	65.2	800	1017	7.25						
18		820	1018	6.99						
19	• • •	920	1014	6.18						
20		890	1015	6.03						
21		900	1016	6.37						
22		760	1018	6.79						
28	• • •	630	1020	6.06		• • •				
Daily aver	-	883	1.018	7.48	0.516	1.28				

It will be seen by a study of the analytical data that Dr. Underhill had an average daily excretion of nitrogen through the kidneys from October 26, 1903, to June 23, 1904, of 7.43 grams. From October 26 to April 13, a period of nearly six months, the average daily excretion was 7.81 grams of nitrogen, while from the latter date to June 23 the average daily excretion amounted to 6.68 grams of nitrogen. Taking the body-weight at 65 kilos, the lower nitrogen figure would mean a metabolism of 0.102 gram of nitrogen per kilo of body-weight, closely akin to the figures obtained with the two preceding subjects.

An excretion of 6.68 grams of nitrogen corresponds to the metabolism of 41.75 grams of proteid matter, while the excretion of 7.43 grams of nitrogen implies the metabolism of 46.4 grams of proteid, being less than one-half the ordinarily accepted requirement for the healthy adult.

During this long period of eight months the body-weight was stationary, carrying with it the assumption that the body was in a condition of nitrogen equilibrium. It should be emphasized in connection with Dr. Underhill's case, that throughout the entire period of eight months, and more, there was perfect freedom in the choice of food. Further, it will be noticed by a study of the dietary made use of in the balance experiments that Dr. Underhill did not reduce his nitrogen intake by any exclusion of meat. He practically made use of his ordinary diet, such as he had always been accustomed to, but with a decided diminution of the amount of proteid food, accompanied by a gradual reduction in the total amount of food consumed each day. His diet, therefore, was in no sense a vegetable diet. Meat was conspicuous in his daily food, but naturally in reduced quantities.

On February 9 a nitrogen balance was attempted, in which a careful comparison of the nitrogen content of all intake and output was made for a period of six days. By a study of the results of this balance period, shown in the accompanying tables, it will be noticed that not only was there no deficiency of nitrogen, but the body was laying on nitrogen at the rate of 0.158 gram per day. Further, it will be observed that the fuel value of the food per day averaged only a little over 2000 calories. Yet this amount of food, with its comparatively low fuel value, carried with it only 8.83 grams of nitrogen per day. Upon this quantity of food the body was able to maintain itself, with a little nitrogen to spare and with sufficient fuel value in the food to supply all the energy required for muscular contraction, mental effort, and the maintenance of body temperature.

### Tuesday, February 9 1904.

Breakfast. - Bread 22 grams, tea 146 grams.

Lunch. — Macaroni 129 grams, fried sweet potato 85.5 grams, bread 59 grams, butter 15 grams, fig cake with wine sauce 115 grams, sugar 15 grams, coffee 210 grams.

Dinner. — Bread 27.9 grams, beef 48 grams, potato 207.5 grams, butter 19 grams, pie 272 grams, coffee 210 grams, sugar 10 grams.

Food.	Grams.	Per	r cent Nitrogen.	Total Nitrogen.
Bread: $22 + 59 + 27.9 =$	108.9	×	1.61 =	1.753 grams.
Tea	146.0	×	0.018 =	0.026
Macaroni	129.0	×	0.87 =	1.122
Sweet potato	85.5	×	0.28 =	0.240
Butter $15 + 19 =$	84.0	×	0.088 =	0.030
Fig cake	115.0	×	0.69 =	0.793
Sugar $15 + 10 =$	25.0	×	0.00 =	0.000
Coffee (lunch)	210.0	×	0.099 =	0.207
Beef	48.0	X	2.64 =	1.267
Potatoes	207.5	×	0.36 =	0.747
Pie	272.0	×	0.49 =	1.332
Coffee (dinner)	210.0	×	0.15 =	0.315
Total nitroger	n in food .			. 7.882 grams-
Total nitroge	n in urine			. 7.490

Fuel value of the food . . . 2232 calories.

# Wednesday, February 10, 1904.

Breakfast. — Bread 33 grams, tea 146 grams.

Lunch. — Bread 71 grams, butter 10.5 grams, apple fritters 119 grams, coffee 210 grams, sugar 21 grams.

Dinner.—Roast pork 80 grams, bread 55.7 grams, butter 27.8 grams, cranberry sauce 150 grams, coffee 210 grams, sugar 21 grams.

Food.		Grams.	Pe	r cent Nitroge	m.	Total Nitrogen.
Bread .	33 + 71 + 55.7 =	159.7	×	1.65 =	=	2.635 grams.
Tea		146.0	X	0.075 =	=	0.109
Butter .	10.5 + 27.8 =	38.3	×	0.088 =	=	0.033

Apple fritters .						119.0	×	0.4	5	=	:	0.535
Coffee (lunch) .						210.0	×	0.11	l	=	:	0.231
Sugar	2	1 +	- 21	l	=	42.0	×	0.00	)	=	:	0.000
Roast pork						80.0	×	4.80	)	=	:	8.840
Cranberry sauce						150.0	×	0.04	ŀ	=	:	0.060
Coffee (dinner) .						210.0	×	0.11	Ĺ	=	:	0.281
	To	tal	ni	tro	ogen	in food						7.674 grams.
	To	otal	ni	tro	ogen	in urine						6.140

Fuel value of the food . . . . 1694 calories.

## Thursday, February 11, 1904.

Breakfast. — Bread 88.7 grams, tea 146 grams.

Lunch. — Bread 97.5 grams, butter 31.5 grams, sweet potato 108.7 grams, spaghetti 82.5 grams, peaches 89.4 grams, coffee 210 grams, sugar 21 grams.

Dinner. — Bread 75 grams, butter 21.5 grams, roast beef 116 grams, lemon pie 188.5 grams, coffee 210 grams, sugar 21 grams.

Food.	Grams.	P	er cent Nitrogen	. Total Nitrogen.
Bread . $38.7 + 97.5 + 75 =$	211.2	×	1.75 =	3.696 grams.
Tea	146.0	×	0.084 =	0.122
Butter 31.5 + 21.5 =	58.0	×	0.088 =	0.046
Sweet potato	108.7	×	0.31 =	0.386
Spaghetti	82.5	×	0.47 =	0.387
Peaches	89.4	×	0.09 =	0.080
Coffee (lunch)	210.0	×	0.096 =	0.201
Sugar $21 + 21 =$	42.0	X	0.00 =	0.000
Roast beef	116.0	×	8.00 =	8.480
Lemon pie	188.5	×	0.61 =	1.149
Coffee (dinner)	210.0	×	0.18 =	0.273
Total nitroger	in food			. 9.770 grams.
Total nitroger	n in urine			. 8.020

Fuel value of the food . . . . 2384 calories.

### Friday, February 12, 1904.

Breakfast. — Bread 84 grams, tea 146 grams.

Lunch. — Bread 42 grams, butter 20 grams, codfish cake 72 grams, potato 100 grams, custard 100 grams, coffee 210 grams, sugar 21 grams.

Dinner. — Bread 46 grams, butter 25.5 grams, veal 53 grams, potato 75 grams, lima beans 50 grams, apple dumpling 120 grams, coffee 210 grams, sugar 21 grams.

Food.	Grams.	P	er cent Nitro	gen.	Total Nitrogen.
Bread $84 + 42 + 46 =$	122.0	X	1.71	=	2.086 grams.
Tea	146.0	X	0.045	=	0.065
Butter 20 + 25 5 =	45.5	X	0.088	=	0.040
Codfish cake	72.0	X	1.57	=	1.180
Potato	100.0	Х	0.41	=	0.410
Custard	100.0	X	0.83	=	0.880
Coffee (lunch)	210.0	X	0.078	=	0.163
Sugar $21 + 21 =$	42.0	X	0.00	==	0.000
Veal	<b>53.0</b>	X	5.51	=	2.920
Potato	75.0	X	0.87	=	0.277
Lima beans	50.0	X	0.90	=	0.450
Apple dumpling	120.0	X	0.72	=	0.860
Coffee (dinner)	210.0	Х	0.12	=	0.252
Total nitroger	n in food .				9.483 grams.
Total nitroger					8.820

Fuel value of the food . . . 1785 calories.

# Saturday, February 13, 1904.

Breakfast. — Bread 85 grams, tea 146 grams.

Lunch. — Bread 57.5 grams, butter 28 grams, sweet potato 100 grams, coffee 210 grams, sugar 21 grams, quince 105.5 grams, apple turnovers 153 grams.

Dinner. — Bread 35.5 grams, butter 11 grams, peas 70 grams, roast beef 50 grams, pie 169 grams, coffee 210 grams, sugar 21 grams.

Food.	Grams.	Per	cent Nitrog	en.	Total Nitrogen.
Bread . $35 + 57.5 + 35.5 =$	128.0	×	1.64	=	2.099 grams.
Tea :	146.0	×	0.083	=	0.048
Butter 28 + 11 =	89.0	×	0.088	=	0.034

Sweet potato			100.0	X	0.87	=	0.370
Coffee (lunch)			210.0	X	0.15	=	0.815
Sugar	21 + 2	l =	42.0	X	0.00	=	0.000
Quince			105.5	X	0.047	=	0.049
Apple turnovers			158.0	X	0.96	=	1.468
Peas			70.0	X	0.96	=	0.672
Roast beef .			50.0	X	8.22	=	1.610
Pie			169.0	X	0.43	=	0.726
Coffee (dinner)			210.0	X	0.11	=	0.231
	Total ni	trogen	in food				7.622 grams
	Total ni	trogen	in urine				7.170

Fuel value of the food . . . . 2186 calories.

# Sunday, February 14, 1904.

Breakfast. — Bread 81 grams, tea 146 grams.

Lunch. — Bread 70.5 grams, butter 29 grams, potato 123 grams, apple sauce 127.5 grams, coffee 210 grams, sugar 21 grams.

Dinner. — Bread 20.5 grams, butter 23 grams, chicken 101.5 grams, potato 80 grams, succotash 80 grams, chocolate cake 103 grams, ice cream 125.8 grams, coffee 210 grams, sugar 21 grams.

Food.	Grams.	Per o	ent Nitroger	n. Total Nitrogen.
Bread $. 31 + 70.5 + 20.5 =$	122.0	×	1.75	= 2.135 grams.
Tea	146.0	X	0.068	= 0.091
Butter $29 + 23 =$	52.0	X	0.088	= 0.045
Potato	123.0	X	0.41	= 0.504
Apple sauce	127.5	X	0.029	= 0.037
Coffee (lunch)	210.0	X	0.10	= 0.210
Sugar $21 + 21 =$	42.0	X	0.00	= 0.000
Chicken	101.5	X	5.08	= 5.150
Potato	80.0	X	0.37	= 0.296
Succotash	80.0	X	0.57	= 0.456
Chocolate cake	103.0	×	0.75	= 0.772
Ice cream	125.8	X	0.58	= 0.729
Coffee (dinner)	210.0	X	0.11	= 0.231
Total nitrogen	in food			10.656 grams.
Total nitrogen	in urine			7.730

Fuel value of the food . . . . 2231 calories.

### NITROGEN BALANCE. - Underhill.

	Nitrogen	Output.	
	•	itrogen in Urine.	Weight of Fraces (dry).
Feb. 9	7.832 grams.	7.49 grams.	9.5 grams.
10	7.67 <b>4</b>	6.14	• •
11	9.770	8.02	82.0
12	9.488	8.82	9.5
18	7.622	7.17	28.0
14	10.656	7.73	24.0
	<del></del>		108.0 grams contain
			6.52 % N.
	58.087	45.37 +	6.715 grams nitrogen
	50.005	*** FO 005	***

53.087 grams nitrogen. 52.085 grams nitrogen.

Nitrogen balance for six days = +0.952 gram. Nitrogen balance per day = +0.158 gram.

#### Average Intake.

Calories per day . . . . . . . . . . . 2068.

Nitrogen per day . . . . . . 8.83 grams.

On June 1, near the close of the experiment, a second nitrogen balance was attempted, with a view to seeing if under the existing conditions of life and activity it was possible to reduce the fuel value of the food, and at the same time diminish in still greater measure the quantity of proteid food taken. The results of this trial are seen in the accompanying tables, where for four days a careful comparison of output and intake is shown. It will be observed from these tables that the average fuel value of the food per day was reduced to 1785 calories, and that the average intake of nitrogen was restricted to 6.73 grams per day. The balance shown is a minus balance, although the deficiency per diem is not large. It is very obvious that both the nitrogen and fuel value of the food can be reduced considerably below the average maintained during the period of the first balance, but not to the degree attempted in the second balance, and secure nitrogen or body equilibrium. It will be noted in this last nitrogen balance, that the average daily output of nitrogen through the urine amounted to 6.46 grams, while the average nitrogen intake was 6.73 grams. The fuel value of the food, however, averaged only 1785 calories per day. It is quite evident, if the non-nitrogenous food had not been reduced quite so much, that a plus nitrogen balance would have been obtained. it is obvious that under the conditions of life and activity, this subject needed to metabolize only about 40 grams of proteid per day, with the total fuel value of his food equal to about 2000 calories, in order to secure both body and nitrogen equilibrium. Here, again, we have an illustration of the possibility of physiological economy which shows itself in a reduction of the daily proteid food more than 50 per cent, with a reduction of the non-nitrogenous food fully 30 per cent, and with maintenance of health, strength and vigor equal, if not superior, to that which the subject has ever experienced.

#### UNDERHILL

## Wednesday, June 1, 1904.

Breakfast. — Champagne wafer 21.5 grams, tea 146 grams, sugar 15 grams.

Dinner. — Bread 67.4 grams, butter 15.1 grams, beefsteak 55.5 grams, potato
128 grams, asparagus 103.5 grams, tea 146 grams, sugar 15 grams,

banana 88.5 grams.

Supper. — Bread 15.7 grams, butter 11.8 grams, banana and nuts 106 grams, crullers 90.2 grams, coffee 145 grams, sugar 15 grams.

Food.	Grams.	Per c	ent Nitrogen.	Total Nitrogen.
Champagne wafer	21.5	×	0.64 =	0.138 gram.
Tea	146.0	×	0.015 =	0.022
Sugar $15 + 15 + 15 =$	45.0	×	0.000 =	0.000
Bread $67.4 + 15.7 =$	83.1	×	1.46 =	1.212
Butter 15.1 + 11.8 =	26.9	×	0.10 =	0.027
Banana	83.5	×	0.28 =	0.192
Beefsteak	55.5	×	4.72 =	2.620
Potato	128.0	×	0.32 =	0.410
Asparagus	103.5	×	0.35 =	0.862
Tea	146.0	×	0.033 =	0.048
Crullers	90.2	×	1.04 =	0.938
Banana and nuts	106.0	×	0.69 ==	0.731
Coffee	145.0	×	0.060 =	0.087
Total nitrogen	in food			. 6.787 grams.
Total nitrogen	in urine			. 5.960

Fuel value of the food . . . . 1913 calories.

## Thursday, June 2, 1904.

Breakfast. — Bread 30 grams, tea 146 grams, sugar 15 grams.

Dinner. — Clam chowder 270.5 grams, bread 64 grams, butter 14.5 grams, lobster 88.5 grams, Saratoga chips 15 grams, caramel custard 79 grams, tea 146 grams, sugar 15 grams.

Supper. — Bread 35.2 grams, butter 18.8 grams, pineapple 148.7 grams, crullers 84.8 grams, tea 140 grams, sugar 15 grams.

<b>T</b> o	od.								Grams.	Per	r cent Nitre	ogen.	Total Nitrogen.
Bread									30.0	×	1.46	=	0.438 gram.
Tea.									146.0	X	0.045	=	0.066
Sugar		1	5 +	- 1	5 +	- 1	5	=	45.0	X	0.00	=	0.000

#### UNDERHILL.

Bread       64 + 35.2       =       99.2       ×       1.44       =       1.428         Butter       14.5 + 13.8       =       28.3       ×       0.10       =       0.028         Clam chowder	
Clam chowder 270.5 × 0.34 = 0.920 Lobster	
Lobster	
Saratoga chips 15.0 × 0.97 = 0.146	
Caramel custard $79.0 \times 0.77 = 0.608$	
Tea	
<b>Tea</b>	
Pineapple 148.7 $\times$ 0.054 = 0.080	
Crullers 84.8 $\times$ 1.04 = 0.882	
Total nitrogen in food 6.060	grams.
Total nitrogen in urine 5.810	0

Fuel value of the food . . . . 1921 calories.

## Friday, June 3, 1904.

Breakfast. — Bread 31 grams, tea 146 grams.

Dinner. — Fried ham 61 grams, asparagus 124.2 grams, baked potato 85 grams, bread 29 grams, butter 23.5 grams, tea 140 grams, sugar 15 grams, pineapple 74.8 grams.

Supper. — Bread 34 grams, butter 17.8 grams, pineapple 158.5 grams, tomato 132.5 grams, cup cake 121 grams, tea 140 grams, sugar 15 grams.

Food.	Grams.	Per	cent Nitrogen.	Total Nitrogen.
Bread $81 + 29 + 84 =$	94.0	×	1.44 =	1.854 grams.
Tea	146.0	×	0.039 =	0.057
Fried ham	61.0	×	4.90 =	2.989
Asparagus	124.2	×	0.50 =	0.621
Baked potato	85.0	×	0.55 =	0.468
Butter $23.5 + 17.8 =$	41.8	×	0.10 =	0.041
Tea	140.0	×	0.027 =	0.038
Sugar $15 + 15 =$	30.0	×	0.000 ==	0.000
Pineapple . $.74.8 + 158.5 =$	233.3	×	0.065 ==	0.152
Tea	140.0	×	0.054 ==	0.076
Tomato	132.5	×	0.150 =	0.199
Cup cake	121.0	×	0.85 =	1.029
Total nitrogen	in food			. 7.024 grams.
Total nitrogen	in urine			. 7.300

Fuel value of the food . . . . 2011 calories.

#### UNDERHILL.

## Saturday, June 4, 1904.

Breakfast. — Bread 32 grams, tea 140 grams.

Dinner. — Hash 138 grams, peas 151.5 grams, bread 61.4 grams, butter 16.1 grams, custard 150 grams, sugar 15 grams, tea 140 grams.

Supper. — Bread 62.5 grams, butter 17 grams, tea 140 grams, sugar 15 grams.

Food.		Grams.	Per cer	at Nitroge	n. Tot	al Nitrogen.
Bread		82.0	×	1.44	=	0.461 gram.
Tea		140.0	×	0.021	=	0.029
Hash		138.0	×	1.50	=	1.995
Peas		151.5	×	1.04	=	1.576
Bread	61.4 + 62.5 =	128.9	×	1.47	=	1.821
Butter	.16.1 + 17 =	88.1	×	0.10	=	0.038
Custard		150.0	×	0.78	=	1.170
Sugar	15+15 =	30.0	×	0.00	=	0.000
Tea		140.0	×	0.038	=	0.046
Tea		140.0	×	0.080	=	0.042
	Total nitroger Total nitroger					7.173 grams. 6.780

Fuel value of the food . . . . 1297 calories.

## NITROGEN BALANCE. - Underhill.

		rogen en in. N	ltrogen in U	Output rine. V		'æces (dry).	
June 1	6	.787 grams.	5.96 gr	rams.	10 g	rams.	
2	6	.060	5.81		10		
8	7.	.024	7.30		25		
4	7.	.178	6.78		6		
	_				51 g	rams contain	
						5.81% N.	
	27	.044	<b>25.85</b> °	+	2.96	33 grams nitrogen.	,
	27	.044 grams ni	trogen.	28.81	3 grams	— nitrogen.	
	Nitrogen be	alance for fou	r days	=	-1.76 gr	ams.	
	Nitrogen ba	dance per daj	7	=	-0. <b>44</b> 2 g	grams.	

## Average Intake.

Calories per day	•	•	•	•	•	•	•	•	1785.
Nitrogen per day									6.78 grams.

Dr. Arthur L. Dean, Instructor in Plant Physiology in the Sheffield Scientific School, twenty-five years of age, and weighing 64 kilos, likewise became a subject of study in this investigation. He is a man of strong physique, and as an undergraduate student at Harvard University trained for various athletic events. He began on the experiment October 13, 1903, and continued until April 3, 1904. From October 13th to October 27 he followed his usual dietary habits, simply reducing in some measure the amount of food consumed. During this period of fifteen days, the average excretion of nitrogen per day through the kidneys was about 12 grams. On the 28th of October he began to reduce in still greater measure the amount of proteid food eaten, and gradually diminished the extent of his proteid metabolism, although not to the same degree as the preceding subjects. He had full freedom of choice in the character and quantity of his diet, but his food was characterized by a predominance of vegetable matter, with an almost complete exclusion of meat.

For a period of nearly six months, or more exactly, from October 28 to April 3, the average daily output of nitrogen through the kidneys amounted to 8.99 grams, while the average daily output of uric acid was 0.386 gram. This daily excretion of 8.99 grams of nitrogen implies a metabolism of 56.18 grams of proteid. This, to be sure, means a reduction in proteid metabolism of about 50 per cent, as compared with the Voit and other standards, but does not show an economy equal to that practised by the preceding subjects. It is to be noted, however, in Dr. Dean's case, that the body-weight did not show at any time in the experiment a tendency to diminish. In fact, all through the experiment his body-weight was a little higher than at the beginning.

DEAN.

		Rody.			Urine.		
Dat	6. ·	Body- weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> .
190	3	kilos	c.c.		grams	gram	grams
Oct.	18	63.2	1100	1017	10.68	0.323	2.38
ł	14		1050	1020	9.83		
	15		1390	1018	14.93	0.432	1.98
l	16		1870	1017	11.26	• • •	
	17		1350	1019	11.18		
	18		1670	1017	11.62	0.490	2.51
	19	63.2	1075	1018	10.19		
ŀ	20		1260	1015	9.98	0.488	1.88
l	21	• • • •	1350	1020	13.04	• • •	• • •
Í	22		1405	1015	11.88	0.481	2.07
	23		1240	1020	11.98		
	24		1200	1028	11.16	• • •	
	25	• • •	1365	1018	11.97	0.395	2.02
ł	26		1195	1020	10.75	• • •	
l	27		1240	1021	11.90	0.477	1.97
Į.	28		1704	• • •	1		
	29	• • •	1105	• • •	9.29	0.898	1.69
i	30	64.5	1180	• • •	daily	daily	daily
	81		910		average	average	average
Nov.	1	• • •	710	• • •	ן יו		
l	2		1540		lı l		
l	8		1250		[]		
ł	4	• • •	1800		8.54	0.854	1.65
t	5		975	• • •			
Ī	6	• • • •	1150	• • •	ן ען		
•	8	64.6	1192	1025	10.86	0.498	I
ľ	9		845		lı l		
l	10		1255	• • •			
1	11		1245				
•	12		1540		8.84	0.868	1.85
ł	18		940		[] [		
į	14		1300	• • •			
•	15		1065		J		
1	16	64.5	1165	• • •	ا مرد ا		
	17		910		9.45	• • •	• • •
ł	18		840		10.59	0.444	
ł	19		1720	1012	10.82	0.381	1.95
1	20	'	640	1022	8.60	0.336	1.75

DEAN.

	Rody			Urine.		
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> .
1903	kilos	c.c.		grams	gram	grams
Nov. 21		1310	1016	9.27	0.378	h I
22		720	1026	8.42	0.415	1.75
23	64.7	900		1)		daily av.
24		1520				
25		1095	• • •	11		ł l
26		710		8.22	0.322	1.74
27		700		daily	daily	
28		1140		average	average	
29		1200	• • •	Į)		]
30		820		13	ļ	
Dec. 1		1335	• • • •	11	1	
2	64.5	940		11		
3		970		8.15	0.862	1.68
4		1240		11		
5		1190		H		
6	• • •	720		1)	,	
7		1160		13	İ	
8		960		11	1	
9	• • •	850		H		
10		935		9.12	0.342	2.11
11	• • • •	945	• • •		l	
12		1425				1
18	64.3	1065	• • • •	15	1	
14		770		9.08	l	l
15		790		B		
30 81	66.1	1230	1017	11		1
1904	00.1	1525	1020	11		1
Jan. 1	1	1010	1021	8.60	0.875	1.80
Jan. 1		1270	1021	11		1
3		1230	1020			1
3		820		lí .		
5		1425	1018	11		1
1 6	1	1100	1018	11	1	
7	1	1025	1021	8.42	0.838	1.76
8	1 :::	760	1021	0.32	0.555	1
9	:::	1450	1019	11		ļ
10	65.8	1410	1016	11	l	1
11		1030	1017	8.51	0.428	l
L						

DEAN.

	Body.			Urine.		
Date	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> .
1904	kilos	0.0.		grams	gram	grams
Jan. 12	65.0	880	1028	l)		
13		790	1025	$\mathbb{H}$		
14		1070	1020	8.51	0.428	
15		1800	1019	daily	daily	• • •
16		925	1025	11	average	
17		1100	1028	average	average	
18		850	1025	1)		
19		1120	1019	11		
20		1005	1020	11		
21		1270	1020	8.23	•••	
22		980	1020	11		
28		650		11		
24		980	1024	į į		
25	65.5	1000		h	1	
26		940		11	ł	
27		1850		11		
28		840		8.85	0.395	1.67
29		675		11	1	
30		740		[]		1
81		1160	1017	[]		
Feb. 1		700		h		
2	1	860		11		1
3		925		11		
4		590		8.22	0.381	1.19
5		835		11		1
6		935		11		l
7	1	980		IJ		
8		1035	1017	9.00		
9	64.1	800	1027	8.64	0.432	
10		940	1020	8.29	0.314	
11		720	1080	8.55	0.864	
12		860	1028	9.93	0.848	• • •
13		980	1022	8.87	0.379	
14	68.9	1285	1022	8.33	0.285	
15		910		h		
16		1210		9.80	0.801	
17		1455		8.00	0.001	
18		1000	1022	IJ		1
		1	<u> </u>			

# PHYSIOLOGICAL ECONOMY IN NUTRITION 101 DEAN.

D-4-	Body-			Urine.		
Date.	Body- weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P2O8-
1904,	kilos	o.c.		grams	gram	gram
Feb. 19		1420		1		
20		1050		9.30	0.801	
21	64.5	1365		daily av.	daily av.	
22		765		1)		
23		1070		11		1
24		1080		9.685	0.423	ŀ
25		1495		1 5.555	U. 120	
26		970				
28		750		D		
29		725		l) l		
Mar. 1	• • •	1100		[]		
2	• • •	990	1024	11		
8		1180	• • •	10.31	0.410	
4	• • •	1100		11		
5	• • •	1010				
6	• • •	970		ן (ן		
7	• • •	790	• • •	1		
8	64.4	670	• • •			
9	• • •	840				,
10	• • • •	1110	• • •	8.99	0.483	
11		1090		11		
12		755	• • •			
18	• • •	1220	• • •	ן ין		
14	• • •	745	• • •	]]		
15	• • •	1030	• • •			
16	• • •	1040	• • •		0.000	
17	• • •	1080		8.24	0.368	•••
18	• • •	1890	• • •			
19		795	• • •	]]		
20	64.0	840	• • •	ן ין		
21 22	• • •	720 970	• • •	1)		
22 28	• • •	1200	• • •	11 1		
28 24	• • •	1200 845	• • •	9.26	0.545	
24 25	• • •	1000	• • •	9.20	0.545	•••
25 26	• • •	1180	• • •			
20 27	• • •	1750	• • •			
28	• • •	820	• • •	9.59	0.409	
28	• • •	020	• • •	80.8	0.409	
		<u> </u>				L

DEAN.

Date. Body-		Urine.							
Date. Body-weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>5</sub> .				
1904	kilos	c.c.		grams	gram	grams			
Mar. 29		710		lı l					
80		1100		11 1					
31		770		9.59	0.409				
Apr. 1		1250		daily	daily				
2		885	• • •	11 - 1					
8	65.0	630	• • •	average	average	• • •			
Daily aver	-	1035	1021	8.99	0.386	1.73			

On February 9, a nitrogen balance was attempted, when for a period of six days a careful comparison of intake and output of nitrogen was made. The results show that while during this period of six days 52.999 grams of nitrogen were ingested, there was an output through the urine and fæces of 61.13 grams of nitrogen, thus indicating a minus balance for this period of 1.355 grams of nitrogen per day. The average daily intake of nitrogen was 8.83 grams. The average daily output of nitrogen through the kidneys was 8.77 grams, being 0.22 gram less than the average daily excretion through the kidneys for the six months' period. The fuel value of the food for this period averaged 2529 calories per day. The nitrogen balance, however, is so strikingly a minus balance that we are forced to conclude the above quantities of food were not quite sufficient to meet the needs of the body under the then existing condi-Still, the fact that the body-weight during the entire period of six months showed no tendency downward implies that during this longer period the body must have been essentially in nitrogen equilibrium, under conditions whereby there was a metabolism of only 56 grams of proteid per day. before stated, this means a physiological economy, as contrasted with existing standards, of about 50 per cent in proteid food. So far as was to be seen, the bodily strength and vigor of the subject, like his body-weight, were fully maintained under the restricted diet, but, for some reason, he did not apparently take as kindly to a reduction of proteid food, and did not accomplish so great a lowering in the rate of proteid metabolism.

Following are the data of the balance experiment:

#### DEAN.

## Tuesday, February 9, 1904.

Breakfast. — Coffee 210 grams, bread 38.5 grams, oatmeal 127 grams, cream 92 grams, sugar 20 grams.

Lunch. — Bread 63.5 grams, butter 12.5 grams, potato 155 grams, consommé 150 grams, stewed tomato 109 grams, coffee 210 grams, sugar 20 grams, pudding 85 grams.

Dinner. — Roast beef 17.5 grams, potato 177 grams, bread 39.1 grams, butter 12 grams, string beans 110 grams, apple pie 237.6 grams.

Food.	Grams.	Per	cent Nitrog	en.	Total Nitrogen.
Coffee	. 210.0	×	0.10	=	0.210 grams.
Bread $88.5 + 63.5 + 89.1 =$	= 141.1	×	1.36	=	1.918
Oatmeal	. 127.0	×	0.64	=	0.812
Cream	. 92.0	×	0.40	=	0.368
Butter 12.5 + 12 =	= 24.5	×	0.088	=	0.021
Potato	. 155.0	×	0.28	=	0.484
Consommé	. 150.0	×	0.88	=	0.570
Tomato	. 109.0	×	0.19	=	0.207
Sugar $20 + 20 =$	= 40.0	×	0.00	=	0.000
Pudding	. 85.0	×	0.69	=	0.586
Roast beef	. 17.5	×	2.64	=	0.462
Potato	. 177.0	×	0.36	=	0.637
String beans	. 110.0	×	0.26	=	0.286
Apple pie	. 237.6	×	0.49	=	1.164
Coffee	. 210.0	×	0.099	=	0.208
Total nitrog	en in food				7.888 grams.
Total nitrog					8.640

Fuel value of the food . . . . 2576 calories.

#### DEAN.

## Wednesday, February 10, 1904.

Breakfast.—Coffee 210 grams, bread 31.8 grams, cream 50 grams, sugar 40 grams, oatmeal 155 grams.

Lunch. — Bread 77 grams, butter 14.5 grams, apple fritters 193.5 grams, coffee 210 grams, sugar 20 grams.

Dinner. — Bread 82 grams, butter 20 grams, cranberries 150 grams, baked beans, 150 grams, coffee 210 grams, sugar 32 grams.

Food.	Grams.	Pe	r cent Nitroger	. Total Ni	itrogen.
Coffee	210.0	×	0.12 =	0.252	grams.
Bread . $.31.8 + 77 + 82 =$	190.8	×	1.65 =	8.148	_
Cream	50.0	×	0.47 =	0.285	
Sugar $40 + 20 + 32 =$	92.0	×	0.00 =	0.000	
Oatmeal	155.0	×	0.60 =	0.930	
Butter $14.5 + 20 =$	34.5	×	0.088 =	0.030	
Fritters	193.5	×	0.45 =	0.870	
Coffee	210.0	×	0.11 =	0.231	
Cranberries	150.0	×	0.04 =	0.060	
Baked beans	150.0	×	1.40 =	2.100	
Coffee	210.0	×	0.11 =	0.231	
Total nitrogen	in food .			. 8.087	grams.
Total nitroger	in urine			. 8.290	-

Fuel value of the food . . . 2145 calories.

## Thursday, February 11, 1904.

Breakfast. — Bread 49 grams, oatmeal 185.7 grams, cream 64 grams, coffee 210 grams, sugar 35 grams.

Lunch.—Bread 111.8 grams, butter 40.5 grams, sweet potato 287 grams, peach preserve 109.3 grams.

Dinner. — Tomato purée 99 grams, bread 94.5 grams, butter 21.5 grams, beans 138 grams, lemon pie 155 grams, coffee 210 grams, sugar 21 grams.

Food.	Grams.	Per cent	Nitrogen.	Total Nitrogen.
Bread $.49 + 111.8 + 94.5 =$	255.3	× 1.	75 <b>=</b>	4.467 grams.
Oatmeal	185.7	× 0.	40 =	0.742
Cream	64.0	× 0.4	19 =	0.813

#### DEAN.

Coffee								210.0	X	0.0	96	=	0.201	
Sugar			35	+	21	:	=	56.0	×	0.0	0	=	0.000	
Butter		<b>4</b> 0.	5 +	- 2	1.5	=	=	62.0	×	0.0	88	=	0.054	
Sweet potate	0							287.0	X	8.0	1	=	0.889	
Peach presen	rve							109.3	X	0.0	9	=	0.098	
Tomato pure	Śе							99.0	X	0.8	3	=	0.326	
Beans								138.0	X	1.8	0	=	1.794	
Lemon pie								155.0	×	0.6	1	=	0.945	
Coffee								210.0	X	0.1	3	=	0.278	
		7	Γot	al	nit	rog	zen	in food .					10.102	grams.
		2	Γot	al	nit	roj	zen	in urine					8.550	-

Fuel value of the food . . . . 2854 calories.

## Friday, February 12, 1904.

Breakfast. — Oatmeal 192.3 grams, cream 75.5 grams, bread 41.6 grams, coffee 210 grams, sugar 20 grams.

Lunch. - Bread 64.1 grams, butter 14 grams, fish cakes 60 grams, potato 200 grams, custard 107.8 grams.

Dinner. - Bread 60 grams, butter 15 grams, lima beans 100 grams, potato 150 grams, apple dumpling 259 grams.

Food.	Grams.	Per	pent Nitroge	n. Total Nitrogen.
Oatmeal	192.3	×	0.52 =	= 0.999 grams.
Cream	75.5	×	0.50 =	
Bread $.41.6 + 64.1 + 60 =$	165.7	×	1.71 =	2.838
Coffee	210.0	×	0.11 =	= 0.231
Sugar	20.0	×	0 00 =	= 0.000
Butter $14 + 15 =$	29.0	×	0.088 =	= 0.025
Fish cakes	60.0	×	1.57 =	= 0.942
Potato	200.0	×	0.41 =	= 0.820
Custard	107.3	×	0.88 =	= 0.890
Lima beans	100.0	×	0.90 =	= 0.900
Potato	150.0	×	0.87 =	= 0.555
Apple dumpling	259.0	×	0.72 =	= 1.864
Total nitroger	n in food			. 10.436 grams.
Total nitroger	n in urine			. 9.930

Fuel value of the food . . . . 2863 calories.

#### DEAN.

## Saturday, February 13, 1904.

Breakfast. — Oatmeal 150 grams, cream 42 grams, sugar 31 grams, bread 31 grams, coffee 210 grams.

Lunch. — Bread 36 grams, sweet potato 222 grams, butter 17 grams, quince preserve 81.5 grams, apple turnover 154.5 grams.

Dinner. — Potato 175 grams, bread 62 grams, butter 15 grams, peas 100 grams, apple pie 177 grams, coffee 210 grams, sugar 21 grams.

Food.	Grams.	Pe	r cent Nitro	gen.	Total Nitrogen.
Oatmeal	150.0	×	0.43	=	0.645 grams.
Cream	42.0	X	0.50	=	0.210
Sugar $31 + 21 =$	<b>52.0</b>	X	0.00	=	0.000
Bread $81 + 86 + 62 =$	129.0	X	1.64	=	2.115
Coffee	210.0	X	0.11	=	0.231
Sweet potato	222.0	X	0.37	=	0.821
Butter $17 + 15 =$	82.0	×	0.088	=	0.028
Quince preserve	81.5	X	0.047	=	880.0
Apple turnover	154.5	X	0.96	=	1.483
Potato	175.0	X	0.37	=	0.647
Peas	100.0	×	0.96	=	0.960
Apple pie	177.0	×	0.43	=	0.761
Coffee	210.0	X	0.11	=	0.281
Total nitroger	n in food				8.170 grams.
Total nitroge	n in urine	е.	<b>.</b>		. 8.870

Fuel value of the food . . . . 2606 calories.

#### DEAN.

## Sunday, February 14, 1904.

- Breakfast. Quaker oats 150 grams, bread 42.5 grams, coffee 210 grams, sugar 31 grams, banana 88.2 grams, cream 87.6 grams.
- Lunch. Bread 87.8 grams, butter 28.7 grams, potato 210 grams, apple sauce 116.5 grams.
- Dinner. Bread 51.5 grams, butter 12.2 grams, sugar 21 grams, potato 150 grams, succotash 100 grams, chocolate cake 122.5 grams, vanilla ice 110.7 grams.

Food.	Grams.	Per	cent Nitrogen	n. Total Nitrogen.
Quaker oats	. 150.0	×	0.46 =	= 0.690 grams.
Bread 42.5 + 87.3 + 51.5 =	= 181.3	×	1.75 =	= 3.172
Coffee	. 210.0	×	0.11 =	= 0.231
Sugar $31 + 21 =$	= 52.0	×	0.00 =	= 0.000
Banana	. 88.2	×	0.20 =	= 0.176
Cream	. 87.6	×	0.50 =	= 0.438
Butter 28.7 + 12.2 :	= 40.9	×	0.088 =	= 0.035
Potato	. 210.0	×	0.41 =	= 0.861
Apple sauce	. 116.5	X	0.029 =	= 0.038
Potato	. 150.0	×	0.37 =	= 0.555
Succotash	. 100.0	×	0.57 =	= 0.570
Chocolate cake	. 122.5	X	0.75 =	= 0.918
Vanilla ice	. 110.7	×	0.58 =	= 0.642
Total nitrog	en in food .			8.821 grams.
Total nitrog	en in urine			0.000

Fuel value of the food . . . . 2685 calories.

## NITROGEN BALANCE. - Dean.

	Nitrogen Taken in. Ni	Output. trogen in Urine. Wei	ght of Fæces (dry)
Feb. 9	7.888 grams.	8.64 grams.	• • •
10	8.087	8.29	• • •
11	10.102	8.55	18 grams.
12	10.438	9.93	18
13	8.170	8.87	80
14	8.821	8.83	58
•	<del></del>		174 grams contain
			4.90% N.
	52.999	52.61 +	8 52 grams nitrogen.
	52.999 grams n	itrogen. 61.13	grams nitrogen.
Nitro	gen balance for six	days = -	8.131 grams.
Nitro	gen balance per da	y = -	1.855 grams.

#### Average Intake.

 Mr. George M. Beers, Clerk in the Treasurer's office of the Sheffield Scientific School, became a subject of study in January, 1903. Mr. Beers was thirty-eight years of age, and had always enjoyed fairly good health, although of somewhat frail physique. His occupation has always been indoor work as accountant, etc. His body-weight was 61 kilos.

Mr. Beers came to the writer for advice as to possible ways of improving his general health, and when it was learned that he was in the habit of eating large amounts of meat, the suggestion was made to him that it might be wise to ascertain the effect of a diminished quantity of proteid food, and as a result of this advice Mr. Beers began to cut down the amount of meat consumed daily. The effect of this abstention from meat was so noticeable that voluntarily all meat was withdrawn from his diet. With this change in dietary habits there came about a loss of body-weight, which, however, was soon regained.

Commencing with May 14, 1903, the amount of nitrogen excreted from the kidneys was determined from time to time, but not each day, since it was very soon found that Mr. Beers showed great regularity in his dietary habits, and a corresponding regularity in the composition of the output. regularity was due in large measure to the fact that the subject became, for the time at least, practically a vegetarian. The beneficial effects noted in his own experience with abstention from meat led to his voluntarily excluding it from his daily diet, so that from January, 1903, to June, 1904, the subject practically tasted meat, fish, or eggs on only four occasions, namely, the 26th day of November, 1903, May 19, 20, and 21, 1904. With this limitation to a vegetable diet and with regular methods of living, the nitrogenous waste material was found to be constant within very narrow limits. Emphasis, however, should be laid upon the fact that there was no prescription of diet, but perfect freedom of choice, although this choice was limited wholly to vegetable and cereal foods.

The accompanying tables give the output of nitrogen, uric acid, etc., through the kidneys for various dates between May

14, 1903, and June 15, 1904. Scrutiny of these tables shows that the average daily output of nitrogen, so far as the data show, amounted to 8.58 grams. This indicates an average daily metabolism of 53.62 grams of proteid material.

Likewise noticeable is the perfectly steady body-weight throughout this whole period of time, while the low level of 8.5 grams of metabolized nitrogen testifies to an economy in the use of proteid food, which indicates that in this subject at least the needs of the body for proteid food could easily be met by an amount equal to about one-half that called for by the Voit and similar standard dietaries.

BEERS.

	Body-	Urine.							
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>5</sub> .			
1904	kilos	c.c.		grams	gram	grams			
May 14	61.1	940	10 <del>2</del> 0	8.23	0.830	1.82			
Oct. 8		920	1014	6.40	0.874	1.47			
18	61.1	740	1026	8.74	0.875	1.61			
14		780	1024	8.87					
15		970	1018	8.90	0.874	1.04			
16		1365	1014	9.09					
17		1295	1013	8.86					
18		1400	1013	7.56	0.847	1.57			
20	61.1	1304	1014	8.14	0.328	1.44			
21		1510	1012	7.97					
22		1450	1012	8.26	0.388	1.95			
23	• • •	1180	1013	7.86					
24		1060	1019	8.46		• • •			
25	• • •	1275	1015	8.25	0.375	1.74			
26		1390	1014	7.34					
27	61.4	1000	1020	7.50	0.344	1.38			
Nov. 21	61.4	670	1025	8.72					
22	• • •	925	1017	8.21	0.335	1.28			
23		585	1027	8.36		• • •			
24		885	1023	8.02	0.354				
25		800	1026	7.92	•••	• • •			
26		1095	1020	10.77*	0.864	1.58			
Jan. 5	61.4	1110	1018	8.99	0.824				
7		650	1024	8.15	0.278	1.29			
8		710	1020	7.88					
9	•••	910	1028	8.73	::::	• • •			
10	• • •	890	1020	8.01	0 327	1.21			
11	• • •	875	1017	7.51		• • •			
12		480	1029	6.77	:::				
13	• • •	580	1023	8.00	0.398	• • •			
14		765	1022	8.85	•••	• • •			
15		1860	1017	9.22	• • • •	• • •			
16	61.4	990	1020	8.79		• • •			
17	• • • •	835	1024	8.47	0.854	• • •			
18	• • •	780	1020	8.05		• • •			
19 Pab 0	• • •	790	1022	8.10	0.882	• • •			
Feb. 9	• • •	860	1028	8.46	0.855	• • •			

<sup>\*</sup> Turkey eaten on this day.

BEERS.

_	Rody.	· Urine.							
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>p</sub> -			
1904	kilos	C.C.		grams	gram	gram			
Feb. 10	61.4	850	1024	8.57	0.226				
11	61.4	750	1022	8.55	0.352				
12	61.4	595	1026	8.46	0.368				
18	61.4	640	1027	8.75	0.425				
14	60.8	730	1025	8.28	0.358				
Apr. 18	61.5	610	1025	7.69	1				
19	61.5	870	1020	9.87					
20	61.5	985	1019	8.87	[]				
21	61.5	795	1024	8.25	0.365	1			
22	61.5	1090	1019	9.02	daily				
28	61.5	970	1022	8.44	average				
24	61.5	810	1021	7.80		i			
25	61.5	850	1018	6.94	li l				
26	61.5	620	1024	7.11		1			
28		450	1027	8.58					
29		650	1025	9.52	0.691				
30		1260	1018	10.20					
May 1		1060	1017	8.71					
2		610	1025	8.16	l' l				
8		650	1022	8.70					
4	61.5	870	1019	8.51					
5	61.3	610	1025	8.31					
6	61.3	655	1026	8.72	h				
7	:	680	1025	8.53	[]				
8	60.8	490	1026	7.00		i			
9	61.3	550	1026	7.59	0.356	I			
10		705	1025	7.78	11 1				
11		780	1025	8.28					
12		800	1025	9.86	]]				
16		715	1022	7.29	ľ l				
17		990	1018	6.95		1			
18		715	1026	7.81					
19		845	1926	10.45*					
20	61.5	1170	1020	11.02*					
21	61.2	795	1025	10.02*					
22		835	1020	8.42					
23		695	1025	9.42					
24		660	1025	9.82					
					<u> </u>				

<sup>\*</sup> On these three days, meat, fish, and eggs were eaten.

BEERS.

	Body-			Urine.		
Date.	Body- weight.	Volume. 24 hours.	Šp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> ;
1904	kilos	c.c.		grams	gram	grams
May 25		700	1025	9.91	n l	
26		620	1026	8.59	]]	
27		665	1026	9.58	0.484	
28	61.2	960	1021	10.31	daily	
29		790	1023	9.24	average	
81	••••	930	1025	10.44		
June 1		855	1018	8.76		
2		850	1023	8.01		
3		860	1020	9.24	1	
4	61.4	1145	1021	10.17		
5		590	1024	7.47		
6		510	1029	7.53	1 1	
7		620	1027	8.26		
8		985	1020	8.45	1 1	
9		1220	1020	8.49		
10		1220	1017	8.28		
11		1710	1013	8.82		
12		925	1017	7.11		
18		920	1020	8.01		
14		1090	1015	8.64		
15	61.5	915	1022	8.68	•••	
Daily a	verage	880	1021	8.58	0.365	1.49

On February 9, a nitrogen balance was attempted covering a period of six days, in which there was an exact comparison of the nitrogen income and output. In this balance period it will be observed that the total amount of nitrogen taken in for the six days was 53.108 grams, while the output of nitrogen through the kidneys amounted to 51.07 grams. The nitrogen excreted through the fæces, however, brought the total nitrogen output up to 58.83 grams, thus making a minus balance for the six days of 5.722 grams of nitrogen. During this period the average fuel value of the food per day was 2168 calories. The average daily output of metabolized

nitrogen during the period was 8.5 grams, practically identical with the average daily excretion of nitrogen through the kidneys for the entire year, so far as determined. We have here a distinct minus balance, due either to an insufficient amount of proteid food, or an insufficient fuel value.

#### BEERS.

## Tuesday, February 9, 1904.

Breakfast. — Oatmeal 237.5 grams, butter 10 grams, sugar 85 grams, milk 60 grams, coffee 210 grams.

Lunch. — Macaroni 142 grams, cheese 10.5 grams, bread 71.5 grams, sweet potato 119.5 grams, milk 250 grams.

Dinner. — Bread 80.7 grams, butter 20 grams, mashed potato 176 grams, string beans 77.5 grams, apple pie 82 grams, milk 250 grams.

Food.	Grams.	Pe	r cent Nitr	ogen.	Total Nitrogen.
Oatmeal	237.5	X	0.64	=	1.520 grams.
Butter $10 + 20 =$	30.0	×	0.088	=	0.026
Sugar	35.0	X	0.00	=	0.000
Milk . $60 + 250 + 250 =$	560.0	X	0.52	=	2.912
Coffee	210.0	X	0.10	=	0.210
Macaroni	142.0	×	0.87	=	1.235
Cheese	10.5	X	4.45	=	0.467
Bread $71.5 + 80.7 =$	152.2	X	1.36	=	2.069
Sweet potato	119.5	X	0.28	=	0.834
Mashed potato	176.0	×	0.86	=	0.688
String beans	77.5	×	0.26	=	0.201
Apple pie	82.0	X	0.49	=	0.401
Total nitroge	n in food	١			. 10.008 grams.
Total nitroge	en in urin	ie .			. 8.460

Fuel value of the food . . . . 2094 calories.

## Wednesday, February 10, 1904.

Breakfast. — Oatmeal 299 grams, butter 19 grams, cream 71 grams, sugar 41 grams, coffee 210 grams.

Lunch.—Bread 79 grams, butter 11 grams, boiled potato 155.2 grams, milk 250 grams.

Dinner. — Bread 56 grams, butter 12 grams, baked beans 100 grams, cranberry sauce 150 grams, sugar 21 grams, coffee 210 grams.

Food.	Grams.	Per	cent Nitrogen.	Total Nitrogen.
Oatmeal	299.0	X	0.60 =	1.794 grams.
Butter $19 + 11 + 12$	= 42.0	×	0.088 =	0.036
Cream	71.0	X	0.47 =	0.833
Sugar $41 + 21$	= 62.0	×	0.00	0.000
Coffee (breakfast)	210.0	×	0.12 =	0.252
Bread $79 + 56$	= 135.0	×	1.65 =	2.227
Boiled potato	155.2	×	0.39 =	0.605
Milk	250.0	×	0.55 =	1.375
Baked beans	100.0	×	1.40 =	1.400
Cranberry sauce	150.0	×	0.04 =	0.060
Coffee (dinner)	210.0	×	0.11 =	0.231
Total nit	rogen in food .			. 8.313 grams.
Total nit	rogen in urine			. 8.570

Fuel value of the food . . . . 1728 calories.

## Thursday, February 11, 1904.

Breakfast. — Oatmeal 300 grams, cream 71 grams, butter 10 grams, sugar 41 grams, coffee 210 grams.

Lunch. — Butter 14 grams, bread 126 grams, boiled sweet potato 205 grams, milk 250 grams.

Dinner. — Bread 22 grams, butter 7.5 grams, mashed potato 100 grams, sugar 14 grams, milk 250 grams.

Food.								Grams.	P	er cent Nitr	ogen.	Total Nitrogen.
Oatmeal								300.0	X	0.40	=	1.200 grams.
Cream .								71.0	X	0.49	=	0.847
Butter .	10	+	14	+	7.	5	=	31.5	×	0.088	=	0.027

Sugar	.41 + 14 =	55.0	×	0.00	=	0.000
Coffee		210.0	×	0.096	=	0.201 grams.
Bread	126 + 22 =	148.0	×	1.75	=	2.590
Sweet potato		205.0	X	0.31	=	0.635
Milk	250 + 250 =	500.0	×	0.51	=	2.550
Mashed potato		100.0	×	0.86	=	0.360
	Total nitrogen	in food .				7.910 grams.
	Total nitrogen	in urine				8.550

Fuel value of the food . . . 1979 calories.

## Friday, February 12, 1904.

Breakfast. — Oatmeal 300 grams, butter 10 grams, cream 74 grams, sugar 41 grams, coffee 210 grams.

Lunch. — Bread 86 grams, butter 9 grams, potato 200 grams, sugar 14 grams, milk 250 grams.

Dinner. — Bread 63 grams, butter 14 grams, mashed potato 150 grams, apple dumpling 136 grams, milk 250 grams.

Food.	Grams.	P	er cent Nitr	ogen.	Total Nitrogen.
Oatmeal	. 300	×	0.52	=	1.560 grams.
Butter $.10 + 9 + 14 =$	= 33	×	0.088	=	9.029
Cream	. 74	×	0.50	=	0.870
Sugar $41 + 14 =$	= 55	×	0.00	=	0.000
Coffee	. 210	×	0.11	=	0.231
Bread $86 + 63 =$	= 149	×	1.71	=	2.547
Potato	. 200	×	0.41	=	0.820
Milk $250 + 250 =$	500	×	0.48	=	2.400
Mashed potato	. 150	×	0.37	=	0.555
Apple dumpling	. 186	×	0.72	=	0.979
Total nitroger	in food				9.491 grams.
Total nitroger					. 8.460

Fuel value of the food . . . . 2209 calories.

## Saturday, February 13, 1904.

Breakfast. — Oatmeal 300 grams, butter 12 grams, cream 70 grams, sugar 41 grams, coffee 210 grams.

Lunch. — Bread 80 grams, butter 11 grams, sweet potato 132 grams, sugar 20 grams, milk 250 grams.

Dinner.— Bread 59.5 grams, mashed potato 175 grams, butter 11 grams, sugar 20 grams, apple pie 141 grams, milk 250 grams.

Food.	Grams.	P	er cent Nit	rogen.	Total Nitrogen.
Oatmeal	300.0	X	0.43	=	1.290 grams.
Butter $12 + 11 + 11 =$	34.0	×	0.088	=	0.029
Cream	70.0	X	0.50	=	0.350
Sugar $41 + 20 + 20 =$	81.0	X	0.00	=	0.000
Coffee	210.0	×	0.11	=	0.231
Bread $80 + 59.5 =$	139.5	×	1.64	=	2.287
Sweet potato	132.0	×	0.87	=	0.488
Milk $250 + 250 =$	500.0	×	0.54	=	2.700
Mashed potato	175.0	×	0.37	=	0.647
Apple pie	141.0	×	0.48	=	0.606
Total nitroger	in food .				8.628 grams.
Total nitroger	in urine				. 8. <b>75</b> 0

Fuel value of the food . . . . 2895 calories.

## Sunday, February 14, 1904.

Breakfast.—Oatmeal 300 grams, butter 16 grams, cream 70 grams, sugar 41 grams, coffee 210 grams.

Lunch. — Baked potato 171 grams, bread 72 grams, butter 15 grams, sugar 21 grams, apple sauce 38 grams, milk 250 grams.

Dinner. — Bread 35.5 grams, butter 13.5 grams, mashed potato 180 grams, sugar 20 grams, chocolate cake 111 grams, ice cream 115 grams, milk 250 grams.

Food.	Grams.	Pe	er cent Nitroge	m.	Total Nitrogen.
Oatmeal	800.0	×	0.46 =	=	1.380 grams.
Butter . $.16 + 15 + 13.5 =$	44.5	×	0.088 =	=	0.089
Cream	70.0	×	0.50 =	=	0.850
Sugar $41 + 21 + 20 =$	82.0	×	0.00 =	=	0.000
Coffee	210.0	×	0.11 :	=	0.281
Baked potato	171.0	×	0.41 =	=	0.701
Bread $.72 + 35.5 =$	107.5	×	1.75 =	= .	1.881
Apple sauce	88.0	×	0.029 =	=	0.011
Milk $250 + 250 =$	500.0	×	0.40 =	=	2.000
Mashed potato	180.0	×	0.87 =	=	0.666
Chocolate cake	111.0	×	0.75 =	=	0.832
Ice cream	115.0	×	0.58 =	=	0.667
Total nitroger	in food .				8.758 grams.
Total nitroge	n in urine	•			8.280

Fuel value of the food . . . . 2610 calories.

#### NITROGEN BALANCE. - Beers.

	Nitrogen	Outp	at.	
		Nitrogen in Urine.	Weight of Fraces (dry).	
Feb. 9	10.008 grams.	8. <b>46</b> grams.	• • • • •	
10	8.318	8.57	44.7 grams.	
11.	7.910	8.55	19.0	
12	9.491	8.46	30.0	
13	8.628	8.75	28.0	
14	8.758	8.28	5.0	
			126.7 grams contain	
			6.18 % N.	
	53.108	51.07 +	7.76 grams nitroge	en.
	58.108 grams	nitrogen. 58.8	83 grams nitrogen.	

Nitrogen balance for six days = -5.722 grams. Nitrogen balance per day = -0.958 gram

#### Average Intake.

On May 6, a second nitrogen balance was attempted covering a period of seven days, in which, as before, there was an exact comparison of the income and output of nitrogen. In this period of seven days, as shown in the accompanying tables, the fuel value of the food was essentially the same as in the preceding period, but the amount of proteid food was increased to an average intake of 10.10 grams per day. Under these conditions there was a distinct plus balance for the seven days amounting to 2.425 grams, thus showing that with this quantity of nitrogenous food the body was laying on nitrogen to the extent of 0.346 gram per day. The average daily amount of nitrogen metabolized during this period was only 8.18 grams, being quite noticeably below the average daily amount for the year. In other words, the results of this balance period show that with a consumption of food sufficient to yield about 2200 calories per day, the body of this subject needed to metabolize only 8.25 grams of nitrogen per day to more than maintain nitrogen equilibrium. Following are the tables of results:

## Friday, May 6, 1904.

Breakfast. — Oatmeal 345 grams, butter 7 grams, sugar 30 grams, milk 100 grams, coffee 180 grams.

Dinner. — Bread 67 grams, potato 71 grams, corn 179 grams, pie 183 grams, milk 200 grams.

Supper. — Biscuit 75 grams, butter 11 grams, potato 106 grams, cake 52 grams, apricots 75 grams, milk 280 grams.

Food.		Grams.	Per	cent Nitrogen.	Total Nitrogen.
Oatmeal		845	X	0.40 =	1.880 grams,
Butter	7 + 11 =	18	X	0.11 =	0.020
Sugar		30	×	0.00 =	0.000
Milk 100 -	+200 + 230 =	530	×	0.54 =	2.862
Coffee		180	×	0.14 =	0.252
Bread		67	×	1.39 =	0.931
Potato (dinner)		71	×	0.52 =	0.369
Corn		179	×	0.44 =	0.788
Pie		138	×	0.54 =	0.718
Biscuit		75	×	1.21 =	0.908
Potato (supper)		106	×	0.86 =	0.382
Cake		52	×	0.90 =	0.468
Apricots		75	×	0.21 ==	0.158
	Total nitrogen	in food			. 9.236 grams.
	Total nitrogen	in urine			. 8.720

Fuel value of the food . . . . 2080 calories.

## Saturday, May 7, 1904.

Breakfast. — Oatmeal 882 grams, milk 100 grams, coffee 185 grams, butter 10 grams, sugar 30 grams.

Dinner. — Bread 98 grams, potato 67 grams, rice pudding 141 grams, milk 200 grams.

Supper. — Bread 67 grams, butter 13 grams, potato salad 122 grams, milk 240 grams, coffee 120 grams.

Food.	Grams.	Per	r cent Nitrogen.	Total Nitrogen.
Oatmeal	. 382	×	0.41 =	1.566 grams.
Milk $100 + 200 + 240$	= 540	X	0.54 ==	2.916
Coffee (breakfast)	. 185	×	0.18 =	0.241

Butter	10 + 13 =	28 🗙	0.11 =	0.025
Sugar		30 ×	0.00 =	0.000
Bread	. 98 + 67 =	160 ×	1.33 =	2.128
Potato		67 ×	0.56 =	0.875
Rice pudding .		141 ×	0.76 =	1.072
Potato salad .		122 ×	0.35 =	0.427
Coffee (supper)		120 ×	0.15 =	0.180
	Total nitrogen i	n food		8.930 grams.
	Total nitrogen i	n urine		8.530

Fuel value of the food . . . . 1714 calories.

## Sunday, May 8, 1904.

Breakfast. — Oatmeal 386 grams, butter 10 grams, sugar 35 grams, milk 100 grams, coffee 185 grams.

Dinner. — Rice and chicken gravy 178 grams, boiled onions 186 grams, chocolate pudding 141 grams, milk 150 grams.

Supper. — Potato salad 78 grams, bread 28 grams, chocolate cake 104 grams, milk 220 grams.

Food.	Grams.	Per	ent Nitrogen.	Total Nitrogen.
Oatmeal	. 386	×	0.85 =	1.351 grams.
Butter	. 10	×	0.11 =	0.011
Sugar	. 35	×	0.00 =	0.000
Milk . $100 + 150 + 220 =$	<b>470</b>	×	0.55 =	2.585
Coffee	. 185	×	0.11 =	0.204
Rice and chicken gravy	. 178	×	0.47 =	0.887
Onions	. 186	×	0.25	0.340
Chocolate pudding	. 141	×	1.03 =	1.452
Potato salad	. 78	×	0.35 =	0.256
Bread	. 28	×	1.33 =	0.372
Chocolate cake	. 104	×	0.95 =	0.988
Total nitrogen	in food			. 8.396 grams.
Total nitrogen	in urine			. 7.000

Fuel value of the food . . . . 1995 calories.

## Monday, May 9, 1904.

Breakfast. — Oatmeal 330 grams, butter 10 grams, sugar 35 grams, milk 100 grams, coffee 185 grams.

Dinner. — Bread 73 grams, fried potato 125 grams, boiled onions 118 grams, macaroni and cheese 128 grams, apple pie 110 grams, milk 200 grams.

Supper. — Bread 82 grams, boiled potato 180 grams, butter 12 grams, chocolate cake 114 grams, milk 245 grams.

Food.	Grams.	Per	r cent Nitrogen.	Total Nitrogen.
Oatmeal	830	×	0.44 =	1.452 grams.
Butter 12 + 10 =	22	×	0.11 =	0.024
Sugar	35	×	0.00 -=	0.000
Milk $100 + 200 + 245 =$	5 <b>4</b> 5	×	0.58 =	8.161
Coffee	185	X	0.099 =	0.183
Bread $78 + 82 =$	155	×	1.33 =	2.062
Fried potato	125	×	0.48 =	0.600
Onions	118	X	0.25 =	0.295
Macaroni and cheese	128	×	1.58 =	1.958
Apple pie	110	×	0.55 =	0.605
Boiled potato	130	X	0.80 =	0.390
Chocolote cake	114	X	0.95 =	1.083
Total nitrogen	in food			. 11.813 grams.
Total nitrogen	in urine			. 7.590

Fuel value of the food . . . . 2620 calories.

## Tuesday, May 10, 1904.

Breakfast.—Oatmeal 357 grams, butter 11 grams, sugar 85 grams, milk 100 grams, coffee 185 grams.

Dinner.—Soda biscuit 68 grams, boiled potato 160 grams, butter 20 grams, stewed tomato 108 grams, custard pie 103 grams, milk 200 grams.

Supper. — Soda biscuit 81 grams, butter 14 grams, stewed potato 97 grams, chocolate cake 66 grams, milk 200 grams.

Food.		Grams	. Pe	er cent Nitrogen.	Total Nitrogen.
Oatmeal		357	×	0.42 =	1.499 grams.
Butter		11 + 20 + 14 = 45	×	0.11 =	0.050
Sugar .	•	35	×	0.00 =	0.000

#### BEERS.

Milk 100 -	+ 200	+ (	200	)	=	500	×	0.54	=	2.700
Coffee						185	×	0.13	=	0.241
Soda biscuit .	. 6	38 +	- 81	L.	=	149	×	1.27	=	1.892
Boiled potato .						160	×	0.80	=	0.480
Stewed tomato						103	×	0.21	=	0.216
Custard pie						103	×	0.91	=	0.987
Stewed potato.						97	×	0.45	=	0.437
Chocolate cake						<b>6</b> 6	×	0.95	=	0.627
	Tota	al n	itro	oge	n i	n food				9.079 grams.
	Tota	al n	itro	oge	n i	n urine				7.780

Fuel value of the food . . . . 2190 calories.

## Wednesday, May 11, 1904.

Breakfast. — Oatmeal 394 grams, butter 10 grams, sugar 85 grams, milk 100 grams, coffee 185 grams.

Dinner. — Soup 141 grams, bread 78 grams, butter 14 grams, boiled potato 101 grams, corn 128 grams, rice pudding 116 grams, milk 200 grams.

Supper. — Biscuit 103 grams, butter 11 grams, corn 113 grams, cake 60 grams, milk 205 grams.

Food.	Grams.	Pe	er cent Nitrog	en.	Total Nitrogen.
Oatmeal	394	×	0.46	=	1.812 grams.
Butter $10 + 14 + 11 =$	85	×	0.11	=	0.039
Sugar	85	×	0.00	=	0.000
Milk $100 + 200 + 205 =$	505	×	0.54	=	2.727
Coffee	185	×	0.13	=	0.241
Soup	141	×	0.48	=	0.677
Bread	78	×	1.25	=	0.975
Boiled potato	101	×	0.31	=	0.318
Corn $113 + 128 =$	241	×	0.46	=	1.109
Rice pudding	116	×	0.63	=	0.731
Biscuit	108	×	1.42	=	1.468
Cake	60	×	0.78	=	0.468
Total nitrogen	in food				. 10.555 grams.
Total nitrogen	in urine				. 8.280

Fuel value of the food . . . . 2183 calories.

## Thursday, May 12, 1904.

Breakfast. — Oatmeal 349 grams, butter 10 grams, sugar 35 grams, milk 100 grams, coffee 185 grams.

Dinner. — Soup 137 grams, fried potato 43 grams, bread 100 grams, macaroni and cheese 122 grams, bread pudding 80 grams, milk 200 grams.

Supper. — Stewed potato 126 grams, bread 122 grams, cake 73 grams, apricots 81 grams, milk 220 grams.

Food.	Grams.	Per	cent Nitrogen.	Total Nitrogen.
Oatmeal	. 349	×	0.41 =	1.431 grams.
Butter	. 10	×	0.11 =	0.011
Sugar	. 85	×	0.00 =	0.000
Milk $100 + 200 + 220 =$	<b>520</b>	×	0.57 =	2.964
Coffee	. 185	×	0.13 =	0.241
Soup	. 137	×	0.48 =	0.658
Fried potato	. 48	×	0.76 =	0.327
Bread $100 + 122 =$	222	×	1.25 =	2.775
Macaroni and cheese	. 122	×	1.94 =	2.870
Bread pudding	. 80	×	0.82 =	0.656
Stewed potato	. 126	×	0.43 =	0.542
Cake	. 78	×	0.78 =	0.569
Apricots	. 81	×	0.23 =	0.186
Total nitrogen	in food			. 12.780 grams.
Total nitrogen	in urine			. 9.360

Fuel value of the food . . . . 2283 calories.

## NITROGEN BALANCE. - Beers.

	Nitrogen	Output.						
	Taken in.	Nitrogen in Urine.	Weight of Fraces (dry).					
May 6	9.236 grams.	8.72 grams.	17.5 grams.					
7	8.930	8.53	<b>30</b> .5					
8	8.896	7.00	26.2					
9	11.818	7.59	27.1					
10	9.079	7.78	22.1					
11	10.555	8.28	26.0					
12	12.730	9.36	24.5					
			12.5					
			186.4 grams contain					
			5.98 % N.					
	70.789	57.26 +	11.054 grams nitrogen.					
	70.739 grams ni	itrogen. 68.81	4 grams nitrogen.					
Nitro	ogen balance for seve	en days =	+2.425 grams.					
Nitro	gen balance per day	=	+0.346 gram.					
	Ave	rage Intake.						
Calo	ries per day		2152.					

Nitrogen per day . . . . . . . . . 10.10 grams.

Here we have, as in the preceding cases, marked physiological economy of non-nitrogenous as well as of nitrogenous food material. Further, taking the body-weight of the subject as 61.5 kilos, and with an average daily excretion of 8.58 grams of metabolized nitrogen, it is evident that under the existing conditions of life and activity there was need for the metabolism of only 0.139 gram of nitrogen per kilo of body-weight. Doubtless, however, still greater economy was possible.

Finally, while it hardly savors of scientific accuracy to quote simple sensations, yet it may be stated that the subject asserts a betterment of his condition, with continuance of mental and physical vigor in such a degree that he has persisted now for more than a year and a half in the maintenance of these dietetic habits which are characterized by this lowered rate of proteid metabolism. The quantity of nitrogen metabolized daily means the breaking down of approximately 50 grams of proteid, and it is very evident that this amount of proteid food, one-half the amount called for by the ordinary diet, is quite sufficient to meet all of the subject's bodily needs, even with a total fuel value considerably below 2500 calories.

#### SUMMARY

Certain general conclusions seem to be justified by the results reported. A healthy man, whose occupation is such as not to involve excessive muscular work, but whose activity is mainly mental rather than physical, though by no means excluding the latter, can live on a much smaller amount of proteid or albuminous food than is usually considered essential for life, without loss of mental or physical strength and vigor, and with maintenance of body and nitrogen equilibrium. This means that the ordinary professional man who leads an active and even strenuous life, with its burden of care and responsibility, need not clog his system and inhibit his power for work by the ingestion of any such quantities of proteid food as the ordinary dietetic standards call for.

There is no real physiological need — that is apparent — for the adoption of such dietetic habits as ordinarily prevail, or as are called for by the dietary standards set by most authorities in this branch of physiology. There is no justifiable ground for the dictum, or the assumption, that the adult man of average body-weight needs daily 118 grams of proteid food for the maintenance of health, strength and vigor, or that there is need for the metabolism of at least 16 grams of nitrogen daily. If such were the case, how could these five subjects, whose experiences have been detailed in the foregoing pages, have maintained their body-weight, established nitrogen equilibrium, pursued their ordinary vocations without loss of strength and vigor, and kept in a perpetual condition of good health, with an average daily metabolism of from 5.4 grams of nitrogen to 8.99 grams of nitrogen for periods ranging from six to eighteen months? Surely, if 16 to 18 grams of nitrogen are a daily requisite for the healthy adult, there should have been some sign of nitrogen starvation during these long periods of low proteid diet, but the sharpest scrutiny failed to find it. On the contrary, there were not wanting signs of improved conditions of the body which could not well be associated with anything but the changes in diet.

Let us briefly consider the main facts. The writer, of 57 kilos body-weight, showed for nearly nine consecutive months an average daily metabolism of 5.7 grams of nitrogen. During the last two months the daily metabolism averaged 5.4 grams of nitrogen. As body-weight and nitrogen equilibrium were both maintained under these conditions, it is certainly fair to assume that the physiological needs of the body were fully met. These figures imply a metabolism, in the first instance, of 0.1 gram of nitrogen per kilo of body-weight, while the lower figure shows a metabolism of 0.094 gram of nitrogen per kilo of body-weight. We may call this latter amount the minimal nitrogen requirement for this particular individual, under which health, strength, and vigor can be fully maintained. This lower nitrogen figure shows

that the needs of this particular individual for proteid material are met by the metabolism of 33.75 grams of proteid per day. Hence, one-third the usually accepted standard of proteid is quite sufficient for the wants of this particular person, and this too with a quantity of non-nitrogenous food far below the daily amount called for by ordinary physiological rules. A fuel value of 2000 calories per day was fully adequate to meet the ordinary wants of the body.

Dr. Mendel, with a body-weight of 70 kilos, showed for seven consecutive months an average daily metabolism of 6.53 grams of nitrogen, likewise with maintenance of health, strength, body equilibrium, and nitrogenous equilibrium. This figure implies a nitrogen metabolism of 0.093 gram per kilo of body-weight and shows that the wants of the body—in his case—can be fully met by a metabolism of 40.8 grams of proteid matter daily, and this likewise without increasing the amount of non-nitrogenous food ingested. Indeed, a total fuel value of 2500 calories per day was quite sufficient for all the needs of his body under the existing conditions.

Dr. Underhill, with a body-weight of 65 kilos, showed for six consecutive months a proteid metabolism equal to 7.81 grams of nitrogen per day, while for the last two months the daily average excretion of nitrogen was only 6.68 grams. These figures mean respectively a nitrogen metabolism of 0.120 and 0.102 gram of nitrogen per kilo of body-weight. Here, too, as in the preceding cases, this lowered rate of proteid metabolism was maintained without increasing the total fuel value of the food and with a continuance of health and strength.

Messrs. Dean and Beers, with body-weights of 64 and 61.5 kilos respectively, likewise kept up their health and strength for a long period of time with a nitrogen metabolism averaging 8.99 and 8.58 grams of nitrogen per day, i.e., with a metabolism of 0.140 and 0.139 gram of nitrogen per kilo of body-weight respectively, and this with a total fuel value in their daily food averaging not more than 2500 calories.

With this general concurrence of results, we are certainly warranted in the assertion that the professional man can safely practise a physiological economy in the use of proteid food equal to a saving of one-half to two-thirds the amount called for by existing dietary standards, and this without increasing the amount of non-nitrogenous food consumed. Indeed, the latter class of foods can likewise be diminished in amount without detriment to health or strength, where there is no call for great physical exertion. Lastly, the so-called minimal proteid requirement of the healthy man - which for this group of individuals we may place at the low level of 0.093 to 0.130 gram of nitrogen per kilo of body-weight - represents the real physiological needs of the system for nitrogen, and in so far as our present data show, anything beyond this quantity may be considered as an excess over and above what is required for the actual physiological necessities of the body. Naturally, however, there may be nothing detrimental in a slight excess of proteid beyond the daily needs. a subject, however, to be discussed later in connection with other results.

In view of the close agreement in the amount of nitrogen metabolized by these different individuals per kilo of body-weight, emphasis should be laid upon the fact that the results recorded were all obtained with perfect freedom of choice in the matter of diet, without prescription of any kind, so that the close concurrence in the final figures tends to strengthen the value of the data as pointing to a certain minimal requirement easily attainable, and fully adequate for meeting the needs of the body.

# II. EXPERIMENTS WITH VOLUNTEERS FROM THE HOSPITAL CORPS OF THE UNITED STATES ARMY.

The original Detachment from the Hospital Corps of the United States Army detailed to serve in this series of experiments arrived in New Haven September 28, 1903, under the command of Dr. Wallace DeWitt, 1st Lieutenant and assistant Surgeon of the United States Army. The detail was composed of twenty men, of whom fourteen were privates, volunteers for the experiment, the remainder being made up of non-commissioned officers, cook, cook's helper, etc. The detachment was located in a convenient house on Vanderbilt Square belonging to the Sheffield Scientific School, and there they lived during their six months' stay in New Haven under military discipline, and subject to the constant surveillance of the commanding officer and the non-commissioned officers.

In selecting the men for the experiment particular attention ws paid to securing as great a variety of types as possible, representing different nationalities, temperaments, etc. Naturally, among such a group of enlisted men brought together for the purpose in view many were found unsuited for various reasons, and were quickly exchanged for others better adapted for the successful carrying out of the experiment. Several quickly deserted, apparently not relishing the restrictions under which they were compelled to live. restriction which constituted the greatest hardship in the eyes of several of the men was the regularity of life insisted upon, and the consequent restraint placed upon their movements in the city when relieved from duty. The following Statement from Dr. DeWitt will explain the causes of removal of the men who dropped out of the experiment during the natural sifting process of the first few weeks and later.

HOSPITAL CORPS DETACHMENT U. S. ARMY, 332 TEMPLE St., NEW HAVEN, CONN. March 17, 1904.

Professor Russell H. Chittenden,

Director Sheffield Scientific School.

New Haven, Conn.

Sir, — In compliance with your verbal request concerning men of this detachment lost by transfer and desertion and the reasons therefor, I have the honor to inform you that the following men were lost by transfer, at my request, for the reasons set after their respective names:

Private Edward McDermott (October 17, 1903). Mentally and morally unsuited.

Private Paul Forkel (October 18, 1903). Physically unsuited.

Private David Acker (October 24, 1903). Physically unsuited by reason of Acute Pulmonary Tuberculosis Bilateral.

Private William C. Witzig (November 17, 1903). Physically unsuited by reason of Cardiac Irritability.

Private Philip S. Myer (December 11, 1903). Physically unsuited by reason of very high grade of Myopia both eyes.

Private first class Charles P. Davis (January 14, 1904).

Physically unsuited by reason of Acute Melancholia.

Private Barnard Bates (February 12, 1904). Morally and mentally unsuited by reason of Drunkenness.

In all these men, except Private Davis, the condition for which they were transferred was present when they reported for duty with the detachment.

In the case of Private Davis, his condition of acute melancholia in my opinion was incident to the experiment,—due to the necessary restrictions of liberty and food, assisted by a natural gloomy disposition.

The following men were lost by desertion:

Private first class Samuel R. Curtis (November 3, 1903). Private first class William Smith (November 5, 1903). Private Simon Prins (January 23, 1904). Private Edwin A. Rinard (February 3, 1904).

Of these men Private first class Smith and Private Prins were on duty in the kitchen and were at no time subject to restriction of diet and liberty. Private first class Curtis deserted before the experiment was well under way and can not be attributed to any cause arising out of the investigation. Private Rinard's desertion was in my opinion due to the restrictions of diet and liberty incident to the experiment. I would say, however, that this man was a worthless character and was discharged "without honor" from the army during a previous enlistment.

Very respectfully,
(Signed) WALLACE DEWITT,
1st Lieut. and Asst. Surgeon U. S. Army,
Commdg. Detachment.

As supplementing Dr. DeWitt's statement it may be mentioned that Rinard reported for the experiment at New Haven on December 11, 1903, and remained here until February 2, 1904. On December 11 he weighed 59.8 kilos, while on February 2, the last day he was here, his body-weight was 60 kilos. Evidently, any restriction of diet he may have suffered had not made any great impression upon his bodily condition.

There were thirteen men of the detachment who really took part in the experiment, and of these all but four were in the original detail. Of these four, two joined in October and two early in November. Of these thirteen, all but two continued to the close of the experiment, April 4, 1904.

The following statement gives the name, age, birthplace, occupation, length of service (U. S. Army), etc. of the thirteen men.

It will be noted that the men range in age from twenty-one years six months to forty-three years, and that representatives of many countries are on the list.

Regarding the duties of the men, i. e., their daily work, the following statement from Dr. DeWitt will give all needed information on this point. The character of the Gymnasium work will be referred to later.

Иапо.	Age on Oct. 1, 1903 (Years and Months).	Length of Bervice on Oct. 1, 1903. (Years and Months).	Birthplace.	Occupation Before Original Enlistment.	Remarks.
Privates first class.					
Broyles, Jonah	24-6	<b>F</b>	Campbell Co., Tenn.	Scholar	Joined Nov. 13, 1908.
Coffman, William E	22-1		Edinsburg, W. Va.	Clerk	Joined Sept. 28, 1908.
Fritz, Charles J	27-8	2	Switzerland.	Gaefftter	Joined Oct. 30, 1903.
Henderson, James D	26-1	9	Marshall, N. C.	Student	Joined Sept. 28, 1903.
Loewenthal, Maurice D	22-8	0-2	New York, N. Y.	Clerk	Joined Sept. 28, 1903.
Morris, William	21-6	8	San Juan, P. R.	Scholar	Joined Sept. 28, 1908.
Oakman, William H	9	1-7	Blackville, S. C.	Bookkeeper	Joined Sept. 28, 1903.
Sliney, William F	26-7	ĵ	Roxbury, Mass.	Painter	Joined Oct. 21, 1903.
Steltz, John J. B	23-1	8-J	Schuylkill, Pa.	Painter	Joined Sept. 28, 1903.
Zooman, Ben	8 8 8	===	London, Eng.	Hosp. Orderly	Joined Sept. 28, 1903.
Private.					
Cohn, Isaac	21-8	Į	Jerusalem, Pal.	Porter in Drug	Joined Nov. 13, 1903.
TRANSFERRED.				store	
Private first class.					
Davis, Charles P	27-0	27	Darlington Co. S. C.	Machinist	Joined Sept. 28, 1908;
Private.					leit Jan. 14, 1904.
Bates, Barnard	26-8	1-10	San Francisco, Cal.	Painter	Joined Sept. 28, 1903;
					1611 F.CO. 12, 1901.

HOSPITAL CORPS DETACHMENT U. S. ARMY, 332 TEMPLE STREET, NEW HAVEN, CONN., March 17, 1904.

Professor R. H. CHITTENDEN,

Director Sheffield Scientific School,

NEW HAVEN, CONN.

- Sir, In compliance with your verbal request I have the honor to inform you that the duties of this detachment during the tour of duty at this station have been as follows:—
- At 6.45 A. M. the men arise and their body-weights are taken immediately, after which they dress and assemble for reveille roll-call.
- 7.15. Breakfast. After this meal they are all engaged in various duties about the quarters, such as inside and outside police, kitchen police, assisting in measurement of urine and fæces and transportation of the same to the laboratory; cleansing of fæces cans and urine bottles, etc. They are occupied with these various duties until about 9 A. M.
- 9 A. M. Detachment proceeds to Gymnasium under charge of a non-commissioned officer, and by him reported to the gymnastic instructor.
  - 11 A. M. Detachment returns from Gymnasium.
  - 12 M. Dinner.
- 1 P. M. Drill, weather permitting, otherwise a walk in charge of a non-commissioned officer, or indoor instruction relative to the duties of a soldier.
  - 2 P. M. Relief from drill; walk, or instruction.
- 2.30 p.m. Instruction by a non-commissioned officer in their duties as nurses, etc.
  - 3.30 p. m. Relief from instruction.
  - 5 P. M. Supper.
  - 5.30 P. M. Retreat roll-call.
  - 10 P.M. All men in bed.

This routine is the same for every day in the week except Saturday and Sunday. On Saturday drill and instruction in the afternoon are omitted, and on Sundays the men are also free from gymnasium work.

In addition to the duties mentioned above, a special detail of two men is made every morning to assist in the weighing and serving of the food, and from time to time details are sent to the laboratory and reported to Dr. Mendel for such work connected with the investigation as he might assign them.

Very respectfully,
(Signed) WALLACE DEWITT,

1st Lieut. and Asst. Surgeon U. S. Army,

Commdg. Det. H. C.


Here we have a group of men, thirteen in number, quite different in type from the preceding group; men accustomed to living a vigorous life under varying conditions, and who naturally had great liking for the pleasures of eating. Further, they were men who had no personal interest whatever in the experiment or in the principles involved. To be sure, they had volunteered for the work, and the objects of the experiment had been fully explained to them. Like good soldiers they no doubt desired to obey orders, and they doubtless preferred to see the experiment a success rather than a failure, but they had not that interest that would lead them to undergo any great personal discomfort. This point should be kept in mind, since it has a distinct bearing upon the possibility of establishing physiological economy of diet in persons who would not willingly incommode themselves or suffer personal inconvenience.

The experiment commenced on October 4, 1903, and for a period of six months every detail bearing upon the nutrition and condition of the men under the gradually changed conditions was carefully observed. Every precaution to preserve the health and good spirits of the men was taken. Pure distilled water was sent to the quarters each morning, magazines and other periodicals were supplied through the courtesy of friends, occasional visits to the theatre were indulged in; in fact all that could be done to counterbalance any possible depressing influence from the partial restraints of the experiment was arranged for.

Regarding the details of the work; the twenty-four hours' urine was collected by each man, also the fæces for each day, and these were subjected to chemical analysis with a view to



Showing a group of the soldiers at work in the gymnasium.



following out the various changes in the amount and character of the proteid metabolism going on in the body as measured by the output of total nitrogen, uric acid, and phosphoric acid in the urine, and by the amount of nitrogen in the fæces with reference to the degree of digestion and utilization of the proteid foods ingested.

During the first two weeks the ordinary army rations were given to the men; later a gradual change was made, accompanied by a reduction in the amount of proteid food, with some reduction likewise in the total fuel value of the food. All the food placed before each man was carefully weighed, and at the close of every meal any uneaten food was weighed and the amount subtracted from the initial weights. During the balance periods, when the income and output of nitrogen were carefully compared, the food materials were weighed with greater care and large samples of each article were taken for analysis, to determine the exact content of nitrogen. Every figure for nitrogen shown in the tables was verified by at least a duplicate chemical analysis so as to avoid any possibility of error.

It is not necessary to give here any detailed description of the changes made in the character of the diet, since on pages 288 to 326 may be found the daily dietary from October 2, 1903, to April 4, 1904, - each meal of the day, - showing the amount and character of the food given the men during the six months period. It should be stated, however, that at no time were the men placed on a cereal diet or on a truly vegetable diet. The object in view was simply to study the possibilities of a general physiological economy in diet, with special reference to the minimal proteid requirement of the healthy man. To be sure, in doing this meats, owing to their high content of proteid, were very much reduced in amount and on many days no meat at all was given, but as the dietary is looked through it will be seen that the main change was from a heavy meat diet (rich in proteid) to a lighter diet, comparatively poor in proteid, with an increasing predominance of vegetable and cereal foods. Condiments are also

noticeable in the diet, together with the ordinary accessories, coffee and tea. Variety was also considered as a necessary factor, not to be overlooked, as contributing largely to the maintenance of a proper physiological condition.

From a study of the tables which follow, showing the chemical composition of the daily urine, it will be seen that during the first sixteen days, viz., from October 4 until October 20 or 21, when the men were on the ordinary army ration with opportunity to eat an abundance of meat, the daily urine frequently contained 16 to 17 grams of nitrogen, showing a metabolism of over 100 grams of proteid on such days. Sliney, indeed, averaged for four days an output of 18.19 grams of nitrogen, while Coffman, Henderson, and Zooman showed an average daily excretion of about 15 grams of nitrogen each for the sixteen days from October 4.

OAKMAN.

	Rody.			Urine.		
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P3O8.
1903	kilos	c.c.		grams	gram	grams
Oct. 4	66.7	1160	1019	16.87	0.549	2.76
5	66.7	1030	1025	12.86		
6	66.0	740	1029	11.85	0.708	1.15
7	66.0	480	1031	10.31		
8	65.4	660	1030	14.30	0.574	1.82
9	65.4	830	1029	15.94		
10	65.4	1440	1018	17.02		
11	66.1	2220	1012	16.12	0.591	2.64
12	66.7	1300	1020	18.38		
13	66.8	2140	1013	15.67	0.610	2.56
14	66.4	1290	1017	12.38		
15	66.7	1780	1017	14.95	0.653	1.93
16	66.7	1520	1017	13.68		
17	66.2	1490	1018	15.20		•••
18	66.2	2030	1014	16.44	0.646	2.66
19	65.8	1580	1017	16.78		
20	65.3	1900	1014	16.19	0.626	1.95
21	65.4	1100	1024	12.07		
22	66.0	1200	1018	11.30	0.502	1.11
23	66.4	2060	1015	11.37	• • • •	
24	67.2	1970	1015	11.88		
25	67.1	1480	1014	8.64	0.411	2.07
26	67.2	1510	1018	11.78		• • •
27	67.2	1890	1015	12.18	0.406	1.99
28	67.4	1620	1014	9.82		
29	67.6	980	1018	6.82	0.493	1.62
80	67.5	820	•1022	10.91		
81	67.0	930		• • • •		• • •
Nov. 1	67.4	1480	1015	10.20	0.460	1.49
2	67.6	1160	1018	10.87		• • •
8	67.4	1160	1020	10.44	0.489	1.53
4	67.1	720	1024	6.61		• • •
5	66.6	640	1029	8.02	0.495	1.17
6	66.5	1200	1016	9.57	• • • •	• • •
7	66.0	840	1023	8.57	::::	
8	66.0	1100	1017	8.32	0.452	1.67
9	66.0	720	1080	8.58	::::	
10	66.4	880	1022	10.19	0.515	1.80
11	66.6	1080	1018	9.72	:::	• • •
12	66.9	920	1023	8.38	0.616	1.54

OAKMAN.

Date.	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> ,
1903	kilos	e.c.		grams	gram	grams
Nov. 18	67.2	800	1025	6.48		
14	66.5	600	1026	7.38		
15	66.3	1360	1014	8.98	0.403	1.44
16	66.4	1160	1025	9.88		
17	66.4	900	1020	6.69	0.343	0.94
18	68.0	1820	1010	7.92	h	
19	<b>6</b> 5. <b>4</b>	1160	1017	5.57	0.486	1.09
20	66.0	1120	1020	8.78	daily av.	daily av.
21	66.4	1020	1020	7.89	,	duniy uv.
22	66.1	1860	1016	8.16	0.413	1.83
23	67.0	2600	1008	8.11		
24	65.9	1140	1015	6.43	0.880	1.90
25	65.9	1800	1011	7.56		
26	65.6	1200	1020	7.63	0.377	1.71
27	66.2	1300	1015	7.41		
28	65.9	1200	1012	6.70		
29	65.9	1480	1019	8.79	0.531	1.78
80	65.4	1540	1011	8.41		
Dec. 1	65.0	1080	1015	7.18	0.484	1.45
2	65.6	1440	1012	8.38		• • •
8	64.5	940	1021	8.58	0.438	1.66
4	65.0	780	1022	7.22		
5	65.4	1280	1016	8.06		• • • •
6	65.4	1880	1012	7.67	0.320	1.86
7	65.0	1600	1013	6.24		
8	64.9	1680	1011	7.86	0.304	1.77
9	65.0	1180	1013	7.74		
10	64.7	1120	1016	7.59	0.808	1.87
11	64.7	860	1021	7.58		• • • •
12	64.8	700	1026	6.34		
18	64.6	880	1022	7.87	0.404	1.47
14	64.7	1540	1017	9.83		
15	64.2	1140	1015	6.98	0.436	1.71
16	64.8	1040	1018	6.74		• • •
17	64.4	980	1018	6.94	0.868	1.05
18	64.7	1605	1015	8.57		• • •
19	64.2	1250	1013	8.77		
20	63.4	1000	1016	7.58	0.462	1.25
21	<b>6</b> 3.6	1190	1016	8.85		
22	68.0	1470	1011	8.47	0.245	1.26

OAKMAN.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> .
1908	kilos	c.c.		grams	gram	grams
Dec. 23	68.6	672	1028	5.88		
24	63.8	980	1017	6.55	0.294	0.89
25	63.5	2310	1008	6.87		
26	63.9	1860	1016	6.92		
27	63.5	1590	1015	7.54	0.880	1.11
28	63.0	1840	1018	8.89		
29	62.9	1145	1018	6.11	0.546	0.72
30	63.2	1300	1020	6.78		
81	68.5	1080	1020	5.96	0.421	1.18
1904	<b>,</b>	1				
Jan. 1	64.0	2360	1013	8.64	ı	
2	68.6	1270	1018	5.83		
3	64.0	2475	1012	7.42	0.329	1.43
4	68.6	1820	1012	5.63	daily	daily
5	63.0	1520	1013	6.66	average	average
6	63.5	1270	1016	6.71	יי	a.c.ago
7	63.0	1185	1016	6.74	0.344	1.06
8	68.5	870	1022	6.06	• • • •	
9	68.6	1640	1010	6.89		
10	68.6	1240	1015	5.95	0.409	1.42
11	63.5	1740	1012	7.31		
12	63.0	840	1020	6.00	0.439	
13	62.9	885	1021	7.33	0.490	
14	63.0	1425	1015	8.29	0.441	
15	62.8	1000	1023	7.14	0.890	
16	62.9	1525	1015	8.23	0.872	
17	62.7	1740	1017	8.14	0.400	• • • •
18	62.3	1200	1020	8.42	]	
19	62.7	990	1028	7.60		
20	62.7	985	1020	7.51	0.400	
21	62.7	1080	1021	8.28	0.423	1.32
22	62.7	1670	1010	7.01		
28 24	62.2 62.2	970	1017	6.58 7.99		
24 25	62.2	1800 1680	1013 1013	7.43	ľ.	
		1		1		
26	62.0	880	1026	6.86		
27	62.5	1250	1017	9.07	0.412	1.45
28 29	62.4 62.7	1560 1325	1016	8.61 6.28		
30	62.7	2015	1018	7.25		
30	02.8	2010	1013	1.20	J	

OAKMAN.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>8</sub> .
1904	kilos	c.c.		grams	gram	grams
Jan. 31	62.4	1730	1018	6.64	0.412	1.45
Feb. 1	62.8	910	1027	5.51	1	
2	63.0	1610	1016	7.15	[]	
3	62.5	1880	1020	7.18	]]	
4	62.7	1480	1023	7.46	0.895	1.42
5	63.0	1600	1020	6.62	daily	daily
6	63.2	1980	1012	6.53	average	average
7	63.0	1775	1015	6.39		
8	62.3	1860	1018	8.26	15	
9	62.9	2010	1013	7.36		
10	62.0	1195	1025	7.60		
11	62.9	1580	1016	8.82	0.238	
12	62.5	1900	1018	8.09	"	• • •
18	62.4	1560	1016	7.30		
14	62.5	1480	1017	7.90	11	!
15	62.5	1610	1023	8.98	6	
16	68.0	1570	1019	7.86		
17	62.8	2375	1014	9.69		
18	62.2	1060	1022	7.80	0.430	
19	62.0	910	1022	8.18	0.200	•••
20	62.3	1710	1012	8.10	[]	
20 21	62.6	1940	1012	7.83	11 1	
21	62.4	1250	1021	7.78	ľ, l	
23	62.7	1700	1012	6.53	[]	
23 24	62.4	1525	1012	8.65		
24 25	62.2	1980	1017	8.55	0.489	
26 26	62.0	1145	1013	6.77	0.200	• • •
20 27	61.8	1150	1017	6.87		
27 28	62.0	1445	1019	7.46	H	
1		1015	1020	6.88		
29	62.2 62.6	1225	1024	7.42		• • •
Mar. 1	62.5	1620	1019	7.58		• • •
				1		• • •
3	62.3	1585	1016	6.85		• • •
4	62.8	1815	1015	7.95	l	• • •
5	62.0	1565	1014	6.10		• • •
6	62.0	1700	1020	7.98		
7	62.0	1240	1016	7.44		
8	62.5	1710	1015	8.72	0.411	• • •
9	62.0	1670	1016	7.71		
10	62.5	1590	1016	7.63	ا ۲	

OAKMAN.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric acid.	P <sub>3</sub> O <sub>8</sub> .
1904	kilos	c.c.		grams	gram	grams
Mar. 11	62.0	1410	1021	8.71	h	
12	62.1	1580	1018	7.44	0.411	
13	62.1	1780	1016	8.65	daily av.	
14	62.0	1800	1019	8.11	h	
15	62.0	1820	1012	7.29	!! !	
16	62.2	1670	1017	9.12	]	•
17	62.4	1380	1020	8.20	0.468	
18	62.7	1785	1015	7.82	11 1	
19	62.5	1910	1017	7.68	il i	
20	62.7	1965	1018	6.72	IJ	
21	62.1	930	1026	5.72	h l	. 1
22	62.4	1770	1012	7.86	ll i	
23	62.0	1560	1017	7.21	11 ,	
24	62.0	1860	1015	8.15	0.429	
25	61.6	1180	1023	7.19	li l	
26	62.0	2000	1013	8.88		
27	61.9	1320	1019	7.18	IJ	
28	62.0	1025	1025	6.64	h	
29	62.4	1830	1018	8.34		
80	62.3	1500	1020	6.30	(.379	
31	62.0	1600	1021	7.10		
Apr. 1	62.0	2070	1014	6.83	IJ	
2	62.0	1250	1025	5.55		
3	62.0	2115	1009	4.57	1	
4	62.1	2110	1013	5.95		٠
Daily aver Oct. 21	rage from	1437	1017	7.48	0.405	1.89

144 PHYSIOLOGICAL ECONOMY IN NUTRITION MORRIS.

Date.	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>5</sub> .
<b>190</b> 3	kilos	c.c.		grams	gram	grams
Oct. 4	59.2	970	1028	18.74	0 563	1.46
5	59.2	1340	1018	13.43		
6	58.4	720	1026	9.16	0.546	1.42
7	58.9	685	1028	12.70		
8	58.4	400	1019	5.27	0.179	0.54
9	58.4	820	1027			
10	58.4	1260	1023	13 68	• • •	
11	58.5	1400	1020	18.48	0.782	2.82
12	58.5	1220	1021	13.40		
13	586	1580	1016	18.84	0.614	2.21
14	58.3	1070	1027	16.05	• • •	
15	58.9	850	1029	13.82	0.626	1.50
16	58.9	940	1020	15.40	• • •	
17	59.2	1500	1015	11.70		
18	59.0	1150	1026	15.73	0.796	2.19
19	58.5	1160	1028	18.86		
20	58.4	1160	1026	<b>1</b>		1
21	58.4	1050	1030	] ]		
22	58.6	910	1027	14.62	0.602	1,77
23	58.6	1100	1024	11		
24	58.8	1030	1029	daily	daily	daily
25	59.0	1080	1023	average	average	average
26	59.1	1060	1028	]		
27	58.1	1240	1021			
28	59.1	860	1025			
29	60.0	800	1026	10.30	0.638	1.87
30	59.7	880	1022			
31	59.6	640	• • •			
Nov. 1	60.0	990	1020	ĺį		[
2		750	1028			
8	59.9	900	1027	11		
4	60.0	980	1018		0.40=	
5	59.0	1180	1018	7.60	0.487	1.26
6	59.5	320	1026			
7	58.7	620	1029			
8	58.8	1220	1013	K		}
9	58.6	860	1023			
10	59.4	800	1024	7.08	0.418	1.02
11	59.0	860	1020			
12	59.1	760	1025	1		]

PHYSIOLOGICAL ECONOMY IN NUTRITION 145 morris.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>5</sub> .
1903	kilos	c.c.		grams	gram	grams
Nov. 13	59.2	740	1027			
14	59.1	1160	1014	7.03	0.418	1.02
15	59.0	560	1027	daily av.	daily av.	daily av.
16	58.6	860	1026	li l		
17	58.7	680	1020			
18	58.6	580	1022	11		
19	58.4	920	1016	5.88	0.845	0.84
20	59.0	880	1027			
21	59.5	840	1019			
22	59.0	680	1029	ון		
23	59.8	1040	1015	1)		
24	59.5	1260	1020	11	•	
<b>2</b> 5	59.3	820	1020			
26	59.3	740	1032	7.34	0.477	1.12
27	60.0	1020	1016	11		
<b>2</b> 8	<b>59.4</b>	860	1028			
29	59.4	700	1028	J		
80	59.5	880	1020	]]		
Dec. 1	59.1	1020	1019			
2	<b>5</b> 9.8	1420	1021	11 1		
8	59.2	1240	1027	9.55	0.607	1.63
4	59.5	720	1081	11		
5	59.6	800	1022	11		}
6	59.6	820	1028	li i		
7	59.4	840	1029			l l
8	59.6	540	1020	11.		
9	59.4	880	1026		0.440	4.0
10	59.7	900	1018	7.73	0.410	1.48
11	59.2	780	1025			
12	59.1	740	1028			
18	59.1	820	1022 1028	R I		
14	59.0	840				
15	58.9	1020	1018			
16	58.9	810	1025 1019	6.68	0.382	1.24
17	59.0	1020		0.05	U.88Z	1.24
18	58.6	720 785	1026 1023			
19	58.5		1023	11		
20	58.2	670		K		1
21 22	58.2 58.5	810 680	1081 1026	6.97	0.375	•••

146 PHYSIOLOGICAL ECONOMY IN NUTRITION MORRIS.

	Body-			Urine.		
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P3O8.
1903	kilos	0.0.		grams	gram	grams
Dec. 23	58.6	785	1024	1)		
24	58.6	930	1020			
25	58.8	1040	1017	6.97	0.375	
26	57.6	945	1020	daily	daily	
27	58.6	840	1023	average	average	
28	58.8	1070	1020	1)		
29	58.4	1205	1018	]]		
30	59.0	1000	1026	11		
81	59.0	935	1027	6.70	0.296	1.20
1904				••	0.200	
Jan. 1	58.5	1475	1016	11		daily
2	58.6	1250	1018	11		average
3	58.6	545	1028	li i		
4	58.7	840	1022	1)		
5	58.6	1040	1020	11		
в	58.8	680	1024	11		
7	59.0	1040	1020	6.41	0.882	1.11
8	58.4	• • •	• • •			
9	58.4	1110	1019			
10	58.6	1120	1016	ا (ا		
11	58.9	1010	1017	5.58		• • •
12	58.8	685	1019	4.19	0.405	• • •
18	58.5	800	1029	7.92	0.785	• • •
14	58.0	785	1027	7.91	0.494	• • •
15	58.0	800	1030	7.44	0.488	• • •
16	58.0	1195	1018	7.38	0.421	• • •
17	58.0	880	1020	5.28	0.804	
18	58.0	1080	1026	1)		
· 19	58.0	1075	1019			
20	58.0	920	1019	[]		
21	57.9	715	1031	7.31	0.449	1.40
22	58.0	820	1022			
23	58.0	1065	1024			
24	58.2	1370	1014	[J		
25	58.1	1490	1015	1)		
26	<b>58.2</b>	1300	1025	11		
27	58.2	1335	1019	7.18	0.369	1.33
28	58.3	1110	1014	""	0.000	2.44
29	58.4	915	1028	11		
80	58.4	1200	1020	IJ		

MORRIS.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> .
1904	kilos	c.c.		grams	gram	grams
Jan. 81	58.4	1490	1020	7.18	0.369	1.88
Feb. 1	58.4	870	1030	h		
2	58.9	680	1024	11 1		
8	58.9	1150	1029			
4	58.9	1900	1025	7.04	0.418	1.86
5	59.0	930	1080	daily	daily	daily
6	59.1	1030	1021	average	average	average
7	59.0	975	1024			
8	59.5	880	1080	h l		
9	59.4	970	1027	11 1		
10	59.0	1020	1025	il i		
11	59.0	1015	1021	7.69	0.484	
12	58.9	980	1028			
18	<b>59</b> .8	880	1029			
14	59.2	1150	1018	[]		
15	59.2	1050	1028	li l		
16	59.0	1310	1027			
17	59.4	1300	1017			
18	59.1	1400	1020	7.49	0.428	
19	58.9	930	1080			
. 20	58.7	715	1080			
21	58.7	1240	1023	J J		
22	59.3	915	1025	h l		
23	59.4	1520	1017	]		
24	58.8	940	1027			
25	59.0	1405	1022	6.30	0.471	•••
26	<b>59.4</b>	940	1022	11 1		
27	59.1	1875	1021	11 1		
28	59.1	810	1025	))		
29	59.0	1100	1026	6.40	)	
Mar. 1	59.1	1035	1026	6.64		
2	58.8	990	1025	5.40		
8	59.0	1285	1022	6.55	0.894	• • • •
4	58.9	1075	1025	4.99	1	
5	59.0	1280	1016	5.38		
6	58.8	1230	1026	7.01	J	
7	58.3	1100	1029	ו ו		
8	58.8	1200	1021	7.05	0.723	
9	59.0	1810	1021			'''
10	59.0	1280	1020	ן יין		

148 PHYSIOLOGICAL ECONOMY IN NUTRITION MORRIS.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Urio Acid.	P <sub>3</sub> O <sub>8</sub> .
1904	kilos	c.c.	`	grams	gram	grams
Mar. 11	58.8	1310	1026	1	-	
12	59.0	1350	1022	7.05	0.728	
18	59.1	1110	1025	daily av.	daily av.	
14	58.8	855	1027	1)		
15	58.9	965	1026			
16	58.8	1210	1026			
17	58.8	1410	1022	7.87	0.498	
18	59.0	1500	1020	11		
19	59.0	1290	1026	11		
20	59.0	1040	1024	}		
21	59.0	1040	1024	J)		
22	58.9	980	1028			
23	58.8	880	1030	11		
24	59.0	950	1027	6.67	0.552	
25	59.3	1210	1028			
26	59.2	1210	1024	11		
27	59.2	1210	1022	}		
28	59.1	1280	1026	6.68	1	
29	59.0	1065	1027	5.69	1	
30	59.0	1080	1028	6.06	} 0. <b>446</b>	
81	59.0	1400	1025	6.96	}	
<b>Apr.</b> 1	58.8	1940	1019	7.10	J	
_ 2	59.0	1480	1025	6.13		
8	59.0	1470	1017	4.67	• • •	•••
Daily aver Oct. 20		1017	1023	7.08	0.450	1.25

BROYLES.

	Body-			Urine.		
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>5</sub> .
1903	kilos	c.c.		grams	gram	grams
Nov. 15	59.4	2500	1008	7.85		
16	59.0	2600	1006			
17	59.8	2600	1009	1		
18	58.7	2400		li i		
19	58.0	1280	1014		0.001	
20	58.7	1800	1018	7.95	0.381	1.93
21	58.5	2240	1008	daily	daily	daily
22	59.0	2100	1011	average	average	average
23	58.0	1500	1009	1)		
24	57.7	1700	1011	11		
25	58.0	1080	1014	] [		
26	57.8	1620	1016	6.98	0.826	1.36
27	58.0	700	1026	11		
<b>2</b> 8	58.0	2100	1007	11		
29	58.0	1240	1015	}		
80	58.0	1880	1010	h		
Dec. 1	57.5	1760	1010		'	
2	57.4	1700	1009	} <b>!</b>		
8	<b>57.4</b>	1680	1011	7.40	0.333	1.68
4	57.0	1280	1018	11		
5	57.0	1420	1017			
6	57.6	2600	1009	]]		
7	57.6	600	1021	7.89	1	
8	57.5	1960	1012	10.09	i i	
9	57.5	2060	1010	8.53		
10	57.2	2280	1010	8.89	0.265	1.68
11	56.4	800	1021	6.67		
12	56.8	660	1030	6.77		
18	56.5	920	1021	8.00	J	
14	56.6	1620	1018	1)		
15	56.4	1100	1015	11		
16	56. <b>4</b>	1090	1017	11 - 40		
17	56.2	990	1019	7.48	0.819	1.28
18	56.2	590	1026			
19	56.0	750	1027	]]		
20	56.1	630	1022	J		
21	56.1	1560	1012			
22	56.0	1050	1014	6.41	0.289	0.91
28	56.5	680	1028			J. <b></b>
24	56.4	960	1020	J		

BROYLES.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P3O8.
1903	kilos	c.c.		grams	gram	grams
Dec. 25	56.6	1235	1017	li l		
26	56.5	950	1020	6.41	0.289	0.91
27	56.9	1520	1012	daily av.	daily av.	daily av.
28	56.0	1265	1019	in i	-	
29	55.8	1560	1018	11		
30	55.6	1710	1009	11		
81	56.3	1135	1016	6.70	0.297	1.17
1904				11	•	
Jan. 1	<b>56</b> .0	1110	1016	11		1
2	56.7	1470	1014	IJ į		
8	56.9				• • •	
4	57.2	1790	1010	h		i
5	58.0	1100	1013	11		
6	57.1	640	1028			
7	56.6	1180	1004	5.99	0.871	1.27
8	57.0	1190	1016	11		
9	57.6	810	1025	11		i i
10	56.8	1590	1010	IJ		
11	57.0	1820	1011	6.99		<b>.</b>
12	55.7	525	1027	5.38	0.621	
18	55.9	530	1081	6.99	0.595	
14	55.5	580	1082	7.47	0.514	
15	55.0	1300	1077	9.67	0.428	
16	56.0	1855	1016	7.65	0.886	
17	55.6	800	1020	5.28	0.291	l
18	55.4	1770	1016	h l		
19	55.0	2080	1010			
20	55.6	1285	1017			
21	55.6	1570	1013	7.80	0.364	1.29
22	56.0	2630	1007	[]		
23	55.7	1830	1015	11		
24	56.9	1470	1015	[J		
25	57.0	2140	1009	h l		
26	57.2	1670	1018	11		
27	58. <b>4</b>	2140	1015			
28	58.0	1030	1018	6.81	0.369	1.43
29	58.0	1080	1010	$\Pi$		
80	58.6	1080	1021			
81	58.8	1670	1015	IJ		
Feb. 1	58.4	770	1029	7.06	0.409	1.54

BROYLES.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>6</sub> .
1904	kilos	c.c.		grams	gram	grains
Feb. 2	58.5	1020	1020	1		
3	59.0	1800	1020	11		
4	58.6	1890	1025	7.06	0.409	1.54
5	59.0	1240	1025	11	daily	daily
6	59.6	1280	1019	daily		•
7	59.0	990	1025	average	average	average
8	58.8	1485	1017	lı l		
9	59.4	1900	1013	11		
10	59.0	1580	1025			
11	59.0	1000	1023	7.91	0.488	
12	58.6	790	1081			
18	58.7	1030	1025			
14	58.9	1260	1015	J		
15	59.0	1000	1028	lı l		
16	59.2	2110	1012			
17	59.3	1805	1017	11		
18	59.1	785	1026	7.55	0.376	
19	59.0	1260	1020		•	
20	59.0	1040	1020	11 :		
21	59.0	1775	1012	[J		
22	59.3	1290	1019	h l		
28	59.5	2010	1011			
24	59.7	650	1027			
25	59.4	2300	1011	6.18	0.423	• • •
26	60.5	1145	1019	[ ]		
27	60.0	855	1025	]		
28	60.8	670	1031			
29	60.5	1310	1017	5.74	1	
Mar. 1	60.8	1285 ·	1022	9.26		
2	60.1	1550	1014	6.97		
8	60.2	1470	1014	6.18	0.306	•••
4	60.0	1580	1015	7.68	,	
5	60.5	2060	1008	5.56		
6	60.0	1755	1018	7.69	j	
7	60.0	1230	1016	[]		
8	60.2	950	1030	[]		
9	60.5	1830	1017	9.99	0.428	
10	61.0	1620	1016			
11	60.5	1160	1029			
12	61.0	1850	1025	j .		

# 152 PHYSIOLOGICAL ECONOMY IN NUTRITION BROYLES.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P308.
1904	kilos	c.c.		grams	gram	grams
Mar. 13	61.6	1670	1015	9.99	0.428	• • •
14	62.0	1540	1017	h		
15	61.4	1160	1021	1		
16	61.4	1445	1016			
17	61.4	1610	1019	8.19	0.484	
18	61.5	1220	1018	daily	daily	
19	61.0	1145	1025	average	average	
20	61.3	1155	1017	j		
21	61.2	1230	1021	h		
22	61.3	1850	1018	11		
23	61.2	1180	1019	11		
24	61.4	1490	1016	7.07	0.580	
· 25	61.3	1620	1018			
26	61. <del>4</del>	1040	1028	11		
27	61.0	1160	1021			
28	61.0	1640	1015	6.79	1	
29	61.0	1400	1020	7.06	1	
80	61.0	2055	1014	7.27	0.359	
81	61.0	1190	1028	6.21	0.000	•••
Apr. 1	61.2	1320	1018	5.86		
2	61.0	1005	1028	5.61	J	
8	61.0	2025	1018	7.17	• • •	• • •
Daily average from Nov. 15		1396	1017	7.98	0.898	1.41

#### COFFMAN.

Date   Polymer   Volume   Sp. Gr.   Nitrogen   Uric Acid	P <sub>3</sub> O <sub>3</sub> .  grams 2.03
Oct.         4         59.1         2140         1012         17.33         0.873           5         59.1         1780         1015         15.27            6         58.7         1070         1024         12.62         0.641           7         58.6         1800         1016         16.96            8         58.6         1120         1024         14.49         0.480           9         58.4         1150         1024         10.14	2.03  2.05
5     59.1     1780     1015     15.27        6     58.7     1070     1024     12.62     0.641       7     58.6     1800     1016     16.96        8     58.6     1120     1024     14.49     0.480       9     58.4     1150     1024     10.14        10     58.6     100     100     100     100	2.03  2.05
6 58.7 1070 1024 12.62 0.641 7 58.6 1800 1016 16.96 8 58.6 1120 1024 14.49 0.480 9 58.4 1150 1024 10.14	2.05
6     58.7     1070     1024     12.62     0.641       7     58.6     1800     1016     16.96        8     58.6     1120     1024     14.49     0.480       9     58.4     1150     1024     10.14	2.05
8 58.6 1120 1024 14.49 0.480 9 58.4 1150 1024 10.14	
9 58.4 1150 1024 10.14	1.89
10 500 100 1000	
10 500 0100 1010 1000	
1 10   58.5   2180   1012   16.05	
11 59.1 1580 1014 13.55 0.474	1.75
12 59.1 980 1025 12.99	
18 59.0 1820 1014 14.85 0.613	2.50
14 59.2 1150 1025 13.94	·
15 58.9 2120 1013 16.03 0.337	2.23
16 59.0 1220 1019 14.41	
17   59.0   1680   1019   12.60	:
18 59.6 2720 1011 18.87 0.453	1.86
19 59.4 2360 1017 28.64	8.01
20 58.3 1320 1019	
21 59.1 1080 1024	
22 59.2 650 1029	
23 59.6 1640 1017 13.21 0.475	1.76
24 59.8 1820 1022 daily daily	dail <b>y</b>
25   60.0   2300   1018   average   average   4	average
26 59.8 1440 1022	
27   59.8   1280   1020	
28 60.2 1200 1017 11.40 0.524	1.91
29   60.2   1000   1017   11.40   0.524	1.91
80   59.6   820   1080	
Nov. 1 59.7 1020 1020	
2 59.4 700 1081 1	
3   60.0   880   1025	
4 59.6 560 1031	
5   59.8   540   1082   8.71   0.430	1.61
6   59.1   440   1036	
7   58.6   460   1035	
8 58.4 420 1035	
9 58.6 700 1030	
10 59.0 620 1030	
11   58.5   600   1029   8.61   0.481	1.12
12   58.7   840   1028	
13   58.7   600   1032   J	

#### COFFMAN.

Data	Body-		Urine.						
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P3O3-			
1903	kilos	c.c.		grams	gram	grams			
Nov. 14	58.7	600	1033	8.61	0.431	1.12			
15	58.9	920	1021	5 0.01	0.201	1.12			
16	58.7	720	1030	lì l					
17	58.4	720	1081	11 1					
18	59.3	720	1027						
19	58.7	980	1026	8.41	0.895	1.80			
20	59.0	1400	1022	daily	daily	daily			
21	59.2	600	1029	average	average	average			
22	58.7	800	1031	l					
28	59.0	640	1026						
24	59.5	1120	1022	[]					
25 26	59.6 59.0	1280 820	1017 1028	8.72	0.400	140			
, 20 27	59.0 59.0	740	1028	8.72	0.439	1.49			
28	59.0 59.0	360	1030	11					
29	59.2	1060	1029	11					
30	59.2	760	1032	ľ.					
Dec. 1	58.9	1140	1002	11					
2	59.5	780	1026						
8	58.5	800	1080	11.14	0.586	1.52			
4	59.5	860	1030		0.000	1.02			
5	59.5	920	1030	11					
6	59.5	760	1032	]]					
7	59.4	860	1030	h					
8	59.4	540	1030						
9	59.0	660	1034						
10	59.0	640	1083	9.95	0.400	1.62			
11	58.8	580	1034						
12	59.1	800	1032						
13	58.5	680	1030	J)					
14	57.3	560	1032	h l					
15	58.3	680	1025						
16	58.5	910	1021						
17	58.4	700	1024	7.79	0.372	1.25			
18	58.4	1110	1018						
19	57.8	600	1030						
20	57.8	550	1082	P					
21	58.0	600	1031	1					
22	57.7	470	1037	7.31	0.288	•••			
28	58.0	645	1031	J					

COFFMAN.

Date.	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> .
1903	kilos	c.c.		grams	gram	grams
Dec. 24	58.3	900	1014	h 1		
25	57.8	730	1028	7.31	0.288	1 1
26	58.0	1085	1014	11		
27	58.0	1000	1021	daily av.	daily av.	
28	57.6	1085	1016	h		
29	57.2	1145	1015	7.14		
80	57.4	1002	1023	11 .		1
81	57. <b>4</b>	1800	1016		0.246	1.20
1904		ì		7.60	0.240	daily
Jan. 1	57.6	1240	1020	11		
2	56.4	950	1018	11		average
8	57.0	1038	1021	IJ		
4	57.6	1325	1008	h		i l
5	58.2	1640	1014	[]		1
6	58.6	1090	1017			1 1
7	58.0	1090	1015	7.16	0.271	1.28
8	57.4	785	1026	11		i '
9	57.7	710	1028	11		i l
10	57. <b>4</b>	1080	1014	IJ		
11	57.0	600	1027	8.14		
12	57.0	930	1020	8.82	0.508	
13	56.9	580	1031	8.28	0.508	
14	56.7	1040	1018	8.30	0.812	
15	56.5	650	1033	7.91	0.352	
16	56.5	1130	1017	7.82	0.305	
17	56.5	800	1025	7.44	0.815	
18	56.4	1540	1012	6.19		
19	56.4	1510	<b>-1016</b>	11		
20	56.5	1220	1016			
21	56.2	505	1033	6.95	0.801	1.11
22	56.8	900	1019			
23	56.4	1825	1018	H		
24	56.2	510	1030	į j		
25	56.6	1460	1012	1)		
26	56.7	1400	1015	11		
27	56.7	1520	1018			
28	57.0	1720	1018	7.55	0.840	1.09
29	56.5	520	1085	[]		1
80	56.5	870	1028			
81	56.7	980	1024	[J		
		<u> </u>	<u>'</u>			1

#### COFFMAN.

Date.	Body-			Urine.		
	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P208.
1904	kilos	c.c.		grams	gram	grams
Feb. 1	56.2	700	1032	1		-
2	56.9	1310	1020			
3	57.2	1120	1024			
4	57.3	1260	1028	7.56	0.862	1.24
5	58.0	1970	1018	daily	daily	daily
6	57.0	810	1022	average	average	average
7	56.8	780	1080	IJ		
8	56.6	1130	1021	h l		
9	57.0	1300	1020	[]		
10	568	1140	1025			
11	57.4	1340	1020	8.65	0.420	l
12	57.2	1360	1023			
13	57.0	1020	1025	11		
14	57.0	1720	1014	IJ		
15	56.5	890	1032	)		
16	57.2	1190	1025			
17	57.4	1250	1019			
18	57.0	1680	1015	8.18	0.318	
19	56.7	1225	1025			
20	57.0	900	1025	}		
21	57.0	1590	1016	IJ I		
22	57.0	985	1028	5		
23	57.0	1465	1013			ł
24	56.7	1160	1020	11		
25	56.5	1340	1016	7.62	0.395	
26	56.8	1015	1017			
27 -	56.0	630	1032	11.	•	ļ
<b>28</b>	56.7	1205	1027	]]		
29	57.0	1530	1017	8.63	)	1
Mar. 1	57.0	1030	1022	7.48		ŀ
2	56.8	1295	1020	8.62		ļ
8	56.7	1040	1022	7.18	0.865	
4	56.0	1130	1023	7.93		
5	56.5	1540	1014	7.67		
6	56.6	1105	1024	7.95	]	
7	56.8	1190	1018	l)		
8	56.5	1350	1017	]]		1
9	56.2	880	1031	8.27	0.388	
10	56.8	1500	1016			
11	56.9	1120	1024	IJ l		1

# PHYSIOLOGICAL ECONOMY IN NUTRITION 157 COFFMAN.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P2O5.
1904	kilos	c.c.		grams	gram	grams
Mar. 12	56.6	1600	1018	8.27	0.338	
18	57.2	1230	1022	daily av.	daily av.	
14	57.0	1150	1022	li l	Ţ	
15	57.3	1580	1014			
16	57.8	1290	1028			
17	57.5	1355	1018	8.07	0.288	• • •
18	58.0	1635	1016	[]		
19	<b>56.8</b>	1320	1020	11 1		i
20	57.0	1085	1021	IJ I		
21	57.4	1030	1023	h		
22	57.7	1970	, 1013			
23	57.4	1670	1013			
24	57.0	870	1031	8.50	0.478	
25	57.0	1000	1024			
26	57.3	1320	1023	11		
27	58.0	1500	1018	])		
28	58.1	1485	1019	8.87	) l	
29	58.0	1580	1021	8.06	1	
80	57.8	1415	1019	6.88	} 0.871	• • •
81	57.8	1285	1026	7.78	1 1	
Apr. 1	57.8	1135	1028	7.32	) [	
2	57.0	1415	1022	6.45		
8	58.0	2000	1018	4.12	••••	• • •
•	Daily average from					
Nov. 2	Nov. 2		1024	8.17	0.879	1.23

SLINEY.

	Body-	Urine.						
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	PgOs.		
1904	kilos	c.c.		grams	gram	grams		
Oct. 21	61.8	1990	1021	22.68				
23	61.2	1200	1025	16.77				
24	61.2	1290	1027	17.58				
25	61.4	1700	1016	15.72	0.686	2.59		
27	62.4	1240	1024	1)				
28	62.0	840	1025	11				
29	62.2	630	1030	][	•			
30	62.2	820	1029	11.10	0.664	1.26		
81	62.0	960		daily	daily	daily		
Nov. 1	62.1	780	1030	average	average	average		
2	61.7	940	1027	h l				
8	62.4	1020	1026					
4	61.5	820	1028					
5	61.7	650	1028	10.89	0.579	1.49		
6	62.0	860	1022					
7	61.5	780	1029					
8	61.5	720	1026	IJ				
9	61.7	1180	1020	1)				
10	62.0	620	1028					
11	61.8	880	1027	H				
12	61.8	1000	1027	9.71	0.625	1.39		
18	61.6	920	1028			1		
14	61.4	640	1031					
15	61.0	920	1026	j l				
16	60.5	1000	1026	]]				
17	60.4	1080	1026	11				
18	61.1	880	1029	11	0.500			
19	60.9	940	1020	9.27	0.588	1.12		
20	61.8	1020	1015	[]				
21	60.9	640	1032	11 1				
22	60.4	800	1029	K				
23	61.1	820	1021		1			
24	60.6	700	1027	11				
25	60.6	780	1023 1031	000	0.650	1.07		
26	60.8	780		8.66	0.000	1.07		
27 28	61.7 61.3	1240 1080	1020 1021	11 1				
28 29	61.3	1000	1021	11 1				
80	60.5	820	1029	K				
Dec. 1	60.8	820 820	1026	10.12	0.677	1.88		

SLINEY.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>g</sub> ,
1903	kilos	0.0.		grams	gram	grams
Dec. 2	60.0	600	1080	h l		
8	60.9	940	1015	11		
4	60.9	1000	1024	10.12	0.677	1.88
5	59.9	1580	1016	daily	daily	daily
6	61.0	840	1029	average	average	average
7	60.4	760	1028	8.40	1	
8	60.4	920	1022	9.71		
9	61.0	1000	1020	8.94		
10	60.5	740	1025	8.66	0.671	1.66
11	59.9	760	1031	11.43	1	
12	60.0	660	1033	9.78		
13	59.9	880	1030	11.98	j	· ·
14	60.3	1120	1024	<b>I</b> )		
15	59.5	1060	1021	H		
16	59.9	710	1080	11		
17	59.9	880	1027	10.20	6.52	1.28
18	60.2	1200	1021	11		
19	60.0	1125	1015	11 1		
20	60.0	1210	1021	ן ען		
21	60.0	715	1026	h l		
22	59.5	940	1021	11 1		
23	60.0	895	1028	11 1		
24	59.9	1010	1018	8.97	0.606	• • •
25	59.8	1084	1026	11		
26	61.0	940	1028			
27	60.0	785	1023	ו יו		
28	59.8	1250	1019	l)		
29	59.7	1020	1020	11		
80	60.0	1760	1016	11 1		
81	60.0	980	1022	7.20	0.515	1.29
1904				11 1		
Jan. 1	60.0	1370	1014			
2	60.0	1152	1017	11		
8	60.4	1035	1025	ן א		
4	61.0	1210	1014	}		
5	61.0	1090	1017			
6	61.0	1400	1020	6.67	0.535	1.85
7	60.8	1140	1020			
8 9	60.0 60.0	620	1028	J		

SLINEY.

		Body-	1		Urine.		
De	te.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P308.
19	04	kilos	e.c.		grams	gram	grams
Jan.	10	60.6	1200	1020	6.67	0.535	1.35
i	11	61.8	1230	1015	7.28		
i	12	61.3	560	1027	5.94	0.953	
	18	60.7	625	1029	8.44	0.988	
l	14	60.7	600	1030	7.42	0.674	
	15	60.4	675	1032	7.89	0.693	
l	16	60.5	500	1030	7.23	0.584	1
	17	60.5	700	1030	8.15	0.691	
	18	60.3	500	1082	]}		
•	19	<b>6</b> 0.5	780	1026	11 i		
l	20	60.6	920	1021			1
ľ	21	60.4		• • •	8.13	0.308	1.22
ł	22	60.4	660	1017	daily	dail <b>y</b>	daily
l	28	60.0	920	1080	average	average	average
1	24	60.0	1320	1018	l)		
l	25	60.2	1220	1017	1)		
l	26	60.0	1260	1018	11 1		
l .	27	60.2	845	1024	11 1		
	28	60.8	560	1031	7.14	0.686	0.98
	29	60.0	1030	1024	il i		
	80	60.4	1330	1020	11 1		
ŀ	81	60.6	1125	1019	ן יו		
Feb.	1	60.7	830	1027	n I		
	2	60.8	1695	1016			
	8	61.0	1760	1015	11		
1	4	61.3	1060	1025	7.13	0.645	1.17
	5	61.2	1800	1021	11 1		•
	6	61.8	1880	1014	H		
	7	61.8	1260	1027			
	8	62.4	920	1021			
	8	62.5	1500	1022			
	10	62.6	1145	1026		0.015	
	11	62.8	710	1019	7.66	0.647	• • • •
	12	61.0	1350	1015			
	18	61.6	1080	1027			
	14	61.5	780	1029	lí		
	15	61.5	1010	1030	<b> </b>		
	16	61.7	1175	1025	8.07	0.600	
	17	62.1	1680	1017	5.5.		•••
	18	62.0	1010	1024	J		

SLINEY.

2.4	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid	P308.
1904	kilos	grams		grams	gram	grams
Feb. 19	61.6	770	1028	h	ł	
20	61.0	710	1080	8.07	0.600	
21	61.0	1240	1021	daily av.	daily av.	i
22	61.2	1450	1017	1)	1	
23	61.6	1425	1017	11	1	
24	62.0			8.86	0.746	
26	62.4			0.00	0.720	•••
27	61.4	1080	1019			
28	61.4	835	1029	IJ		
29	61.0	800	1080	9.50	]	
Mar. 1	61.4	775	1028	8.09	1	
2	61.0	760	1030	7.97	1	
8	60.8	920	1023	7.78	} 0.697	• • •
4	61.0	960	1027	7.49		
5	61.0	790	1029	7.54	ļ	
6	61.0	980	1028	8.28	J	
7	61.0					• • •
8	61.0	1480	1017	1)		
9	61.2	1960	1012	11		
10	61.0	740	1028	7.85	0.672	
11	60.8	950	1029	1.00	0.0.2	• • •
12	61.0	1870	1020	11		
13	60.8	1310	1021	IJ		
14	61.0	1460	1014	1)	1	
15	61.2	1155	1019	11		
16	61.3	1100	1025	11		
17	61.0	1465	1013	} 7.72	0.572	• • • •
18	60.9	1300	1020	11	·	
19	61.3	2270	1012		1	
20	61.6	1040	1025	μ		
21	61.2	1020	1027	1)	1	
22	61.4	1210	1019	11		
23	61.0	760	1029			
24	60.5	900	1024	8.64	0.765	•••
25	60.6	840	1029	11	ĺ	
26	60.8	890	1030	11	ļ.	
27	61.0	1000	1024	1 = 00	1.	
28	61.0	870	1026	7.09	0.502	
29	61.0	825	1080	7.87	<b> </b> '	

# 162 PHYSIOLOGICAL ECONOMY IN NUTRITION SLINEY.

	7.3.	Urine.					
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>2</sub> .	
1904	kilos	c.c.		grams	gram	grams	
Mar. 30	60.8	1080	1027	8.10	h		
31	60.6	1030	1026	7.47	0.502	• • •	
Apr. 1	60.6	1130	1021	6.78	daily av.		
2	60.0	1590	1016	6.20			
3	60.6	1860	1013	6.59		• • •	
Daily aver Nov. 1	age from	1021	1094	8.89	0.647	1.89	

STELTZ.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P2O8.
1903	kilos	c.c.		grams	gram	grams
Oct. 4	52.3	1860	1010	11.16	0.825	1.54
5	52.4	1020	1014	7.89		
в	52.0	1120	1018	8.27	0.648	1.47
7	53.1	760	1014			
8	52.0	1280	1012	9.98	0.549	1.70
9	52.0	1600	1014	8.88		
10	52.6	1900	1012	14.59		
11	52.6	1460	1010	7.97	0.368	1.55
12	52.6	1740	1013	10.07		1
13	52.9	2050	1013	12.79	0.758	2.00
14	52.9	1260	1018	8.77		
15	52.6	1540	1018	• 12.20	0.512	1.60
16	52.8	1880	1014	14.41		
17	52.5	1870	1018	14.86		
18	<b>52.4</b>	2280	1018	15.12	0.561	2.53
19	52.5	1560	1010	8.89	• • •	
20	<b>52.4</b>	1880	1013	1)		[
21	52.6	1060	1013	11		[
22	53.2	2100	1011	11.61	0.465	2.01
23	<b>53.4</b>	2320	1012	daily	daily	daily
24	52.9	1460	1017	average	average	average
25	53.4	1660	1016	)		
26	53.2	1150		• • • •	• • •	
27	<b>58.4</b>	1500	1016	j.		
28	53.2	1240	1011	[]		
29 .	53.6	1220	1015	8.65	0.493	1.44
80	58.8	1220	1015	0.00	0.200	1
81	52.9	1120		[ ]		
Nov. 1	58.0	1620	1016			
2	58.8	1640	1014	h		ĺ
8	58.5	1020	1011	11		
4	52.6	1080	1014	11		
5	53.2	1060	1015	6.81	0.364	1.82
6	58.0	1280	1014	11 1		1
7	53.1	1300	1014	11		]
8	52.9	760	1014	ן ון		
9	53.8	1060	1018	h l		Į į
10	58.2	1840	1016	7.31	0.380	1.40
11	58.0	1140	1019		0.500	1.20
12	53.4	1860	1016	ן יו		1
		l		1		1

STELTZ.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P2O8.
1903	kilos	c.c.		grams	gram	grams
Nov. 13	53.4	1800	1016	7.31	0.380	1.40
14	58.2	960	1017	daily av.	daily av.	daily av.
15	53.3					
16	58.2	1640	1016	h l		
17	53.4	1620	1015	11		1
18	53.4	1160	1017			
19	52.9	1940	1015	8.07	0.409	1.45
20	<b>53.4</b>	1800	1014			i
21	53.4	1240	1015			ľ
. 22	58.0	1180	1020	IJ		ł
23	53.4	1320	1018	h		l
24	53.4	2000	• 1013	11		<b>[</b>
25	58.5	1400	1016	11		
26	58.7	780	1026	} 6.71	0.390	1.35
27	53.5	1000	1015	11		
28	53.3	1600	1014	11		,
29	58.5	1340	1018	J		l
30	<b>58.6</b>	860	1020	h		
Dec. 1	53.9	1960	1013	11		
2	53.4	1060	1015	11		
8	52.9	940	1018	7.49	0.394	1.50
4	58.6	1580	1019	11		
5	53.6	980	1014			
6	54.2	1280	1022	ן		
7	58.7	960	1021	7.01	1	
8	53.4	620	1020	4.98		
9	53.4	1000	1016	5.88	0.400	ا مد ا
10	58.5	1420	1015	7.67	0.420	1.46
11	53.5	1360	1018	8.20		
12	58.4	1040	1024	7.92		
13	52.7	920	1022	5.57	Į J	
14	53.0	1840	1024	10.21	• • •	• • •
15	52.9	1000	1019	7.98	0.000	0.75
16	52.9	940	1016	4.79	0.226	0.75
17	58.4	820	1015	1)		
18	53.4	1330	1016	9.04	0.452	1. <b>4</b> 8
19	58.4	1395	1014	11		
20	53.1	1800	1017	ľ		1
21	53.2	1220	1016	7.42	0.387	0.80
22	53.1	1200	1018	1		

STELTZ.

Date.	Body-	Urine.						
Date.	weight.	Volume. 24 hours.	8p. Gr.	Nitrogan.	Uric Acid.	P <sub>2</sub> O <sub>5</sub> .		
1903	kilos	0.6.		grams	gram	grams		
Dec. 23	58.2	1465	1017	h				
24	53.1	1100	1021	11				
25	53.2	970	1016	7.42	0.387	0.80		
26	54.0	1350	1020	daily	daily	daily		
27	58.2	1105	1023	average	average	average		
28	53.0	1240	1020	h				
29	53.0	1180	1023	11				
30	53.4	910	1018	11	ł			
31	58. <b>4</b>	810	1018	6.44	0.356	0.92		
1904				0.22	0.550	0.52		
Jan. 1	58.8	1220	1019	11				
2	58.0	925	1024	11	İ			
8	53.6	870	1021	þ				
4	<b>54</b> .0	1055	1022	1				
5	<b>54</b> .0	1050	1020	11	ļ			
6	53.9	1195	1017	11	l			
7	53.6	980	1021	6.78	0.392	1.15		
8	53.4	1020	1021	]] .				
9	58.3	1080	1022					
10	53.6	1010	1021	IJ				
. 11	58.9	960	1020	6.05		• • •		
12	58.0	620	1021	4.61	0.487			
18	53.6	1165	1019	7.90	0.609	• • • •		
14	52.9	645	1022	4.99	0.298			
15	. 58.0	1450	1020	9.05	0.478	• • •		
16	58. <del>4</del>	1800	1018	7.58	0.405			
17	58.0	1440	1021	8.55	0.476	• • •		
18	53.0	1440	1023	l)				
19	53.0	1115	1021					
20	53.0	1180	1020	] [				
21	52.6	790	1024	6.40	0.386	1.21		
22	52.7	660	1026	[]				
28	<b>52.8</b>	1750	1018	11				
. 24	<b>52.8</b>	1440	1018	h l				
25	52.6	1200	1020	]]				
26	52.8	715	1028					
27	52.7	1625	1015	6.39	0.414	1.21		
28	52.8	1080	1016	5.55	V.111	1.01		
29	52.8	1400	1020	[ [				
80	53.0	1300	1021	ן יו				

#### STELTZ.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P <sub>1</sub> O <sub>8</sub> .
1904	kilos	6.0.		grams	gram	grams
Jan. 31	53.4	1670	1019	6.89	0.414	1.21
Feb. 1	58.0	1800	1022	la l	•	
2	53.2	770	1018	11 1		
8	53.3	1280	1022	]		
4	53.4	1580	1018	6.06	0.351	1.31
5	53.2	1400	1023	daily	daily	daily
6	53.0	1440	1021	average	average	average
7	53.4	1330	1018	ا ً ا		_
8	53.0	1500	1022	1)		
9	58.0	940	1021	11 1		
10	58.2	1400	1022	11 1	1	
11	58.4	1620	1028	} 7.71	0.523	
12	58.6	1645	1018	11 1		
18	53.4	1370	1018	11 1		
14	53.0	1200	1024	}		
15	53.2	1560	1025	1)		
16	58.0	1540	1021			
17	53.5	1610	1020	11 1		
18	53.2	1280	1024	7.85	0.423	
19	58.0	1560	1017	11		
20	53.2	1685	1016	11 !		
21	53.0	1110	1020	11		
22	53.0	1860	1016	lí l		İ
28	53.2	1470	1018	11 1		
24	53.5	1205	1019	11 !		
25	53.5	2140	1014	7.40	0.487	l l
26	58.8	1080	1016	11	7.20,	
27	53.0	1165	1020	11		
28	53.7	1860	1020	IJ İ		
29	54.0	1400	1022	7.90		١
Mar. 1	53.9	1095	1021	5.98	)	
2	58.4	1855	1020	6.58		
8	58.2	2125	1015	8.09	0.000	
4	53.0	1160	1016	4.66	0.889	• • • •
5	53.2	1610	1022	8.69		ł
l š	58.0	1220	1022	8.20	1	
Ĭ	53.0	720	1023	h		
8	53.2	1160	1021		0.455	
ğ	53.0	1280	1020	7.21	0.466	•••
10	52.5	1210	1017	11		

STELTZ.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>p</sub> .
1904	kilos	C.G.		grams	gram	grams
Mar. 11	52.6	1770	1016	h		
12	52.6	1510	1018	7.21	0.466	
18	52.3	1110	1020	daily av.	daily av.	
14	52 6	700	1028	h		
15	52.7	1215	1018			
16	53.0	1840	1018	11		
17	52.6	1685	1014	7.22	0.414	
18	52.4	1770	1015			
19	58.2	1850	1012	11		
20	52.6	910	1018	J)		
21	52.8	1210	1020	li i		
22	52.7	1680	1018	11		
23	53.0	1630	1021	11		
24	52.8	900	1020	7.70	0.550	• • •
25	52.6	1600	1018	11		
26	53.0	1330	1019	<b>[]</b>		
27	52.8	1750	1020	II I		
28	53.0	845	1019	3.60	}	
29	52.8	1490	1021	6.79	1	
30	52.6	1790	1018	7.20	0.834	• • •
81	52.6	1560	1019	7.11		
Apr. 1	52.4	1670	1021	7.82	J	
2	<b>52</b> .6	1165	1013	8.28	• • •	• • •
8	58.0	1570	1018	6.50	• • •	• • •
Daily aver Nov. 2	rage from	1271	1018	7.18	0.416	1.24

# 168 PHYSIOLOGICAL ECONOMY IN NUTRITION HENDERSON.

New   Nitrogen   Nit		Body-			Urine.		
Oct. 4         71.8         1820         1020         17.50         0.635         1.97           5         71.8         1110         1019         9.79             6         71.8         1020         1028         12.67         0.615         1.98           7         71.6         1000         1017         12.00             8         71.2         840         1023         11.29         0.379         1.85           9         71.2         910         1030         14.20             10         71.5         2220         1013         16.78             11         71.7         1280         1022         16.28         0.587         2.04           12         72.2         1860         1021         15.69             12         72.2         1860         1021         15.69             15         72.3         1700         1019         18.46         0.672         2.57           16         72.8         1740         1015         16.70	Date.	Body- weight.		Sp. Gr.	Nitrogen.	Uric Acid.	P20g.
5       71.3       1110       1019       9.79           6       71.8       1020       1028       12.67       0.515       1.98         7       71.6       1000       1017       12.00           8       71.2       910       1030       14.20           10       71.5       2220       1013       16.78           11       71.7       1280       1022       16.28       0.587       2.04         12       72.2       1360       1021       15.59           13       72.4       190       1025       12.34       0.532       1.95         14       72.4       1400       1029       16.04           15       72.3       1700       1019       18.46       0.672       2.57         16       72.8       1740       1015       16.70           17       72.5       1620       1021       17.59           19       72.8       1150       1027       17.73	1903	kilos.	0.0.		grams	gram	grams
6 71.8 1020 1028 12.67 0.615 1.98 7 71.6 1000 1017 12.00 8 71.2 840 1028 11.29 0.379 1.85 9 71.2 910 1030 14.20 10 71.5 2220 1013 16.78 11 71.7 1280 1022 16.28 0.587 2.04 12 72.2 1860 1021 15.59 13 72.4 910 1026 12.84 0.582 1.95 14 72.4 1400 1020 16.04 15 72.3 1700 1019 18.46 0.672 2.57 16 72.8 1740 1015 16.70 17 72.5 1620 1021 17.59 18 72.7 2260 1015 19.26 0.602 2.27 19 72.8 1150 1027 17.78 20 72.5 960 1020 1021 27.78 21 72.5 1060 1029 22 22 72.6 940 1017 23 72.4 1880 1017 24 72.8 1100 1029 22 72.6 940 1017 23 72.4 1120 1029 25 72.6 920 1027 25 72.6 920 1028 27 73.0 1930 1017 25 73.6 820 1026 27 73.0 1930 1017 28 73.1 1340 1020 29 73.6 820 1028 27 73.6 820 1028 27 73.0 1930 1017 28 73.1 1340 1020 29 73.6 820 1028 27 73.0 820 1028 27 73.0 820 1028 27 73.0 820 1028 27 73.0 820 1028 27 73.0 820 1028 27 73.0 820 1028 27 73.0 820 1028 27 73.0 820 1028 27 73.0 820 1028 27 73.0 820 1028 27 73.0 820 1028 27 73.0 820 1028 27 73.0 820 1028 27 73.0 8	Oct. 4	71.8	1820	1020	17.50	0.635	1.97
7 71.6 1000 1017 12.00	5	71.8	1110	1019	9.79		
8 71.2 840 1023 11.29 0.379 1.85 9 71.2 910 1030 14.20 10 71.5 2220 1013 16.78 11 71.7 1280 1022 16.28 0.587 2.04 12 72.2 1860 1021 15.59 13 72.4 910 1026 12.84 0.582 1.95 14 72.4 1400 1020 16.04 15 72.3 1700 1019 18.46 0.672 2.57 16 72.8 1740 1015 16.70 17 72.5 1620 1021 17.59 18 72.7 2260 1015 19.26 0.602 2.27 19 72.8 1150 1027 17.78 20 72.5 950 1020 1021 17.78 20 72.5 950 1020 1021 22	6	71.8	1020	1028	12.67	0.515	1.98
9 71.2 910 1030 14.20	7	71.6	1000	1017	12.00		
10 71.5 2220 1013 16.78	8	71.2	840	1028	11.29	0.379	1.85
11 71.7 1280 1022 16.28 0.587 2.04 12 72.2 1860 1021 15.59 13 72.4 910 1025 12.84 0.532 1.95 14 72.4 1400 1020 16.04 15 72.3 1700 1019 18.46 0.672 2.57 16 72.8 1740 1015 16.70 17 72.5 1620 1021 17.59 18 72.7 2260 1015 19.26 0.602 2.27 19 72.8 1150 1027 17.73 20 72.5 950 1020 21 72.5 1060 1029 22 72.6 940 1017 23 72.4 1880 1017 24 72.8 1100 1029 25 72.6 920 1027 26 72.4 1120 1023 27 78.0 1930 1017 28 78.1 1340 1020 29 73.6 820 1028 31 74.0 1040 Nov. 1 74.8 1540 1017 2 74.0 860 1028 31 74.0 1040 Nov. 1 74.8 1540 1017 2 74.0 860 1028 31 74.0 1040 Nov. 1 74.8 1540 1017 7 72.9 500 1032 8 72.5 920 1023 9 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029	. 9	71.2	910	1030	14.20		
12	10	71.5	2220	1013	16.78		
18	11	71.7	1280	1022	16.28	0.587	2.04
14       72.4       1400       1020       16.04            15       72.3       1700       1019       18.46       0.672       2.57         16       72.8       1740       1015       16.70           17       72.5       1620       1021       17.59           18       72.7       2260       1015       19.26       0.602       2.27         19       72.8       1150       1027       17.78           20       72.5       950       1020       17.78           21       72.5       1060       1029       17.78           21       72.5       1060       1029       1029       14.31       0.612       1.60         22       72.6       920       1027       1029       1024 <th>12</th> <th>72.2</th> <th>1860</th> <th>1021</th> <th>15.59</th> <th></th> <th></th>	12	72.2	1860	1021	15.59		
15 72.3 1700 1019 18.46 0.672 2.57 16 72.8 1740 1015 16.70	13	72.4	910	1025	12.84	0.532	1.95
16       72.8       1740       1015       16.70            17       72.5       1620       1021       17.59            18       72.7       2260       1015       19.26       0.602       2.27         19       72.8       1150       1027       17.78           20       72.5       950       1020       17.78            21       72.5       1060       1029       1029       17.78            22       72.6       940       1017       14.31       0.612       160       daily       average       average       average       12.10       0.557       1.91         20       72.4       1120       1023       12.10       0.557       1.91         30       73.4       960       1028       12.10       0.557       1.91         80       74.0       860       1031	14	72.4	1400	10 <del>2</del> 0	16.04		
17 72.5 1620 1021 17.59  18 72.7 2260 1015 19.26 0.602 2.27  19 72.8 1150 1027 17.78  20 72.5 950 1020  21 72.5 1060 1029  22 72.6 940 1017  23 72.4 1880 1017  24 72.8 1100 1029  25 72.6 920 1027  26 72.4 1120 1023  27 78.0 1930 1017  28 78.1 1340 1020  29 73.6 820 1028  31 74.0 1040  Nov. 1 74.8 1540 1017  2 74.0 860 1031  8 74.0 1240 1024  4 74.0 840 1029  5 73.0 1280 1012  6 73.6 1840 1017  7 72.9 500 1032  8 72.5 920 1023  9 72.4 800 1029  10 72.4 600 1030  9 98 0.563 1600	15	72.3	1700	1019	18.46	0.672	2.57
18 72.7 2260 1015 19.26 0.602 2.27 19 72.8 1150 1027 17.78 20 72.5 950 1020 21 72.5 1060 1029 22 72.6 940 1017 23 72.4 1880 1017 24 72.8 1100 1029 25 72.6 920 1027 26 72.4 1120 1023 27 78.0 1930 1017 28 78.1 1340 1020 29 73.6 820 1028 29 73.6 820 1028 31 74.0 1040 Nov. 1 74.8 1540 1017 2 74.0 860 1031 3 74.0 1240 1024 4 74.0 840 1029 5 73.0 1280 1012 6 73.6 1840 1017 7 72.9 500 1032 8 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029 10 72.4 800 1029	16	72.8	1740	1015	16.70	• • •	
19 72.8 1150 1027 17.78	17 .	72.5	1620	1021	17.59		
20		72.7	2260	1015		0.602	2.27
21 72.5 1080 1029 22 72.6 940 1017 23 72.4 1880 1017 24 72.8 1100 1029 25 72.6 920 1027 26 72.4 1120 1023 27 78.0 1930 1017 28 78.1 1340 1020 29 73.6 820 1028 30 73.4 960 1028 31 74.0 1040 Nov. 1 74.3 1540 1017 2 74.0 860 1031 3 74.0 1240 1024 4 74.0 840 1029 5 73.0 1280 1012 6 73.6 1340 1017 7 72.9 500 1032 8 72.5 920 1023 9 72.4 800 1029 10 72.4 600 1030	19	72.8	1150	1027	17.78		
22	20		1	1080	]		
23			1	1029	[]		
24 72.8 1100 1029 daily daily average 25 72.6 920 1027 27 78.0 1930 1017 28 78.1 1340 1020 29 73.6 820 1026 30 73.4 960 1028 31 74.0 1040  Nov. 1 74.3 1540 1017 2 74.0 860 1031 3 74.0 1240 1024 4 74.0 840 1029 5 73.0 1280 1012 6 73.6 1340 1017 7 7 72.9 500 1032 8 72.5 920 1023 9 72.4 800 1029 10 72.4 600 1030 1030 1030		72.6	1	1017	,,,,,		
25 72.6 920 1027 26 72.4 1120 1023 27 78.0 1930 1017 28 78.1 1340 1020 29 78.6 820 1026 30 73.4 960 1028 31 74.0 1040  Nov. 1 74.8 1540 1017 2 74.0 860 1031 8 74.0 1240 1024 4 74.0 840 1029 5 73.0 1280 1012 6 73.6 1340 1017 7 7 72.9 500 1032 8 72.5 920 1023 9 72.4 800 1029 10 72.4 600 1030 1030 1030 1030 1030 1030 1030	23	72.4	1880	1017			
26 72.4 1120 1028 27 78.0 1930 1017 28 78.1 1340 1020 29 78.6 820 1026 30 73.4 960 1028 31 74.0 1040 Nov. 1 74.8 1540 1017 2 74.0 860 1031 8 74.0 1240 1024 4 74.0 840 1029 5 73.0 1280 1012 6 73.6 1340 1017 7 72.9 500 1032 8 72.5 920 1023 9 72.4 800 1029 10 72.4 600 1030	24	72.8	1100	1029	11 -	•	
27 78.0 1930 1017 28 78.1 1340 1020 29 73.6 820 1028 30 73.4 960 1028 31 74.0 1040  Nov. 1 74.8 1540 1017 2 74.0 860 1081 8 74.0 1240 1024 4 74.0 840 1029 5 73.0 1280 1012 6 73.6 1840 1017 7 72.9 500 1032 8 72.5 920 1023 9 72.4 800 1029 10 72.4 600 1030					average	average	average
28		72.4	1120	1023	l) i		
29 73.6 820 1028 30 73.4 960 1028 31 74.0 1040 Nov. 1 74.8 1540 1017 2 74.0 860 1081 3 74.0 1240 1024 4 74.0 840 1029 5 73.0 1280 1012 6 73.6 1840 1017 7 72.9 500 1032 8 72.5 920 1023 9 72.4 800 1029 10 72.4 600 1030			1 1				
30 73.4 960 1028 31 74.0 1040 Nov. 1 74.8 1540 1017 2 74.0 860 1031 3 74.0 1240 1024 4 74.0 840 1029 5 73.0 1280 1012 6 73.6 1340 1017 7 72.9 500 1032 8 72.5 920 1023 9 72.4 800 1029 10 72.4 600 1030							
31 74.0 1040 Nov. 1 74.8 1540 1017 2 74.0 860 1031 3 74.0 1240 1024 4 74.0 840 1029 5 73.0 1280 1012 6 73.6 1340 1017 7 72.9 500 1032 8 72.5 920 1023 9 72.4 800 1029 10 72.4 600 1030					} 12.10	0.557	1.91
Nov. 1 74.8 1540 1017 2 74.0 860 1031 8 74.0 1240 1024 4 74.0 840 1029 5 73.0 1280 1012 6 73.6 1340 1017 7 72.9 500 1032 8 72.5 920 1023 9 72.4 800 1029 10 72.4 600 1030				1028	]		
2 74.0 860 1081 8 74.0 1240 1024 4 74.0 840 1029 5 73.0 1280 1012 6 73.6 1840 1017 7 72.9 500 1032 8 72.5 920 1023 9 72.4 800 1029 10 72.4 600 1030							
8     74.0     1240     1024       4     74.0     840     1029       5     73.0     1280     1012       6     73.6     1840     1017       7     72.9     500     1032       8     72.5     920     1023       9     72.4     800     1029       10     72.4     600     1030			1		! !		
4     74.0     840     1029       5     73.0     1280     1012       6     73.6     1840     1017       7     72.9     500     1032       8     72.5     920     1023       9     72.4     800     1029       10     72.4     600     1030			1		]]		
5 73.0 1280 1012 8 9.90 0.518 1.71 6 73.6 1840 1017 7 72.9 500 1032 8 72.5 920 1023 9 72.4 800 1029 10 72.4 600 1030	-						
6 73.6 1840 1017 7 72.9 500 1032 8 72.5 920 1023 9 72.4 800 1029 10 72.4 600 1030			1				
7 72.9 500 1032 8 72.5 920 1023 9 72.4 800 1029 10 72.4 600 1030					9.90	0.518	1.71
8 72.5 920 1023 3 3 4 5 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	-				[]		
9 72.4 800 1029 10 72.4 600 1030 9 9 0 563 160	· ·						
10 72.4 600 1030 998 0568 160	_		1		ا بإ		
1 998 0 688 1 160							
11   72.7   900   1022        2.00				•	9.98	0.563	1.60
		•					
12 72.8 780 1032	12	72.8	780	1032	ן יו		

#### HENDERSON.

Date.	Body-			Urine.		
Date.	Body- weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>p</sub> .
1903	kilos	c.c.		grams	gram	grams
Nov. 18	72.6	760	1032			
14	72.0	960	1021	9.98	0.568	1.60
15	72.4	1460	1017	daily av.	daily av.	daily av.
16	72.0	640	1031	n l		l
17	72.0	820	1029			
18	72.0	720	1080			
19	72.0	880	1027	} 9.38	0.478	1.81
20	72.3	1200	1028			
21	73.5	1200	1020			
22	71.5	1180	1021	lí l		
23	71.6	1040	1018			
24	71.3	1040	1028	11		
25	72.0	1020	1026	11		•
26	72.0	1200	1021	} 11.63	0.610	1.61
27	71.6	840	1080			
28	72.0	720	1028			
29	72.5	1400	1022	lí l		
80	72.4	820	1023	<b>{                                    </b>		
Dec. 1	72.0	900	1027	11		
2	72.1	1860	1017			
8	71.4	980	1029	10.69	0.586	1.58
4	71.8	1160	1025			
5	71.8	1740	1018	[ ]		
6	71.6	840	1028	l'		
7	71.0	880	1029	10.41		• • • •
8	71.2	1100	1020	12.54	• • •	• • •
9	71.8	960	1021	11.92	0.400	174
10	71.4 71.2	1220	1019	12.21	0.428	1.74
11 12	71.2 70.6	680 640	1035 1036	11.02 9.60	• • •	• • • •
12	70.6 70.6	940	1036	6.26	• • •	• • • •
18 14	70.0	1280	1018	0.20	• • •	• • • •
14	70.0	1260	1022			
16	70.0	790	1028	H		
17	70.0	980	1020	9.80	0.411	1.57
18	70.1	1060	1020	] 5.00	0.411	1.0,
19	70.0	1510	1018	]]		
20	69.8	790	1018			
20 21	69.5	650	1023	1		
22	70.0	720	1028	7.47	0.438	•••

# ,170 PHYSIOLOGICAL ECONOMY IN NUTRITION HENDERSON.

	Body-						
Date.	weight.	Volume 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P503-	
1903	kilos	c.c.		grams	gram	grams	
Dec. 23	70.0	450	1026	h l			
24	69.6	880	1024	li i			
25	69.5	1800	1019	7.47	0.488	• • •	
26	69.0	980	1024	daily	daily		
27	69.0	840	1024	) average	average		
28	69.0	920	1023	l) l			
29	<b>6</b> 8.8	1180	1020	11 1			
80	69.4	865	1024				
1904 31	70.0	1330	1026	} 7.77	0.407	1.24	
Jan. 1	68.9	890	1021				
2	69.0	947	1027				
8	69.1	1025	1030	ا را			
4	69.2	890	1028	h l			
5	69.8	925	1027	11			
6	69.3	550	1034				
7	69.0	600	1029	7.78	0.439	1.25	
8	68.8	880	1028	11 1			
9	69.0	850	1027				
10	69.0	1860	1015	ן ען			
11	68.9	610	1025	5.89			
12	<b>6</b> 8.5	715	1024	7.68	0.498		
18	68.6	835	1023	8.22	0.551	• • •	
14	68.2	1040	1020	8.24	0.461		
15	68.2	880	1025	7.76	0.440		
16	68.2	970	1023	7.56	0.575		
17	68.0	810	1023	7.87	0.441		
18	68.0	1130	1020	In 1			
19	68.0	1290	1018				
20	68.2	670	1027				
21	67.8	720	1080	7.82	0.445	1.18	
22	67.5	520	1029	11 1			
23	67.6	710	1080	11			
24	67.6	775	1081	ו יו			
25	<b>6</b> 8.0	1220	1018	h l			
26	68.0	815	1016				
27	68.3	920	1029				
28	68.1	950	1023	7.50	0.422	1.06	
29	68.2	670	1028	11 1			
80	68.4	720	1030	][ ]]			
81	<b>6</b> 8. <b>4</b>	1285	1020	PΙ			
L	L			<u>!</u>			

PHYSIOLOGICAL ECONOMY IN NUTRITION 171 HENDERSON.

	Body-					
Date.	weight.	Volume. 24 hours.	8p. Gr.	Mitrogen.	Uric Acid.	P <sub>1</sub> O <sub>5</sub> .
1904	kilos	c.c.		grams	gram	grams
Feb. 1	68.0	790	1080	h		
2	68.1	830	1031	11		
3	68.5	1835	1024	11		
4	68.5	1250	1026	8.27	0.860	1.17
5	69.0	1160	1031	daily	daily	daily
6	69.0	1150	1022	average	average	average
7	68.5	1210	1024	Į)		
. 8	68.0	985	1030	h		
9	68.0	975	1030	11		
10	68.2	990	1032	11		
11	68.6	870	1032	10.40	0.582	
12	69.0	1130	1027	11		
13	69.4	1440	1019	11		
14	69.0	715	1029	IJ		
15	69.0	940	1029	1)		
16	68.0	1070	1027	11		
17	68.1	1080	1027	11		
18	<b>6</b> 8.0	945	1029	11.80	0.510	• • •
19	68.2	1010	1029	11		
20	68.8	925	1029	11		
21	69.0	1200	1020	þ	<b>l</b> ,	
22	68.6	1165	1024	l)		-
23	68.2	1170	1022	11		
24	68.6	1035	1027	11		
25	68.4	1735	1016	7.58	0.640	• • •
26	69.0	775	1029	11	1	
27	68.6	1090	1027	11		
28	69.0	1020	1030	)		
29	69.0	1935	1016	8.36	]	
Mar. 1	68.7	840	1029	6.80	i	
2	68.4	1160	1023	8.28	0.505	
3	68.2	920	1026	7.37	0.521	• • •
4	68.0	1000	1029	8.22		
5	68.0	1645	1018	8.09	11	
6	68.0	1020	1030	8.20	'	
7	68.0	740	1030			
8	68.2	1470	1020	001	0.455	
9	68.0	1660	1019	8.21	U.400	• • • •
10 11	68.0 68.0	2040 1080	1014 1029	}		

172 PHYSIOLOGICAL ECONOMY IN NUTRITION HENDERSON.

	Rody.			Urine.		
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>2</sub> .
1904	kilos	6.6.		grams	gram	grams
Mar. 12	68.2	2450	1014	8.21	0.455	
18	68.6	2800	1014	daily av.	daily av.	
14	68.4	925	1026	h		
15	68.4	1610	1016	11		
16	68.5	1360	1019	]]		
17	68.0			8.82	0.488	
18	68.6	1975	1016	11 1		
19	69.0	2410	1015	11		
20	69.3	2480	1011	}		,
21	68.6	850	1028	h l		
22	<b>6</b> 8.7	1800	1012	11		
23	68.7	980	1028	11 1		
24	69.0	1040	1080	8.64	0.632	
25	69.2	1360	1022	11		
26	69.2	2470	1013	]]		
27	69.3	2110	1015	]]		
28	69.4	1415	1020	8.40	1	
29	69.4	1815	1019	9.04		
80	69.4	1600	1017	5.95	- 0.837	
31	69.0	1890	1018	5.42		
Apr. 1	69.0	1930	1015	6.60	J	
2	69.5	1405	1012	8.20		
3	71.0	1380	1018	7.42	•••	• • •
Daily aver Nov. 1	rage from	1102	1024	8.91	0.488	1.42

FRITZ.

Date.	Body-			Urine.		
	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> ,
1903	kilos	c.c.		grams	gram	grams
Nov. 1	76.0	1000	1024	11.34	1.14	
2	76.4	950	1022	lı l		
8	77.8	2200	1017			
4	76.0	960	1020			
5	76.5	1420	1015	8.97	0.418	1.45
6	76.4	720	1018	daily	daily	daily
7	75.8	940	1020	average	average	average
8	75.3	1580	1012	[]		
9	75.8	2240	1012	h l		
10	75.6	480	1021	11		
11	76.0	1600	1015			
12	76.2	1640	1018	8.27	0.491	1.31
18	76.2	880	1020	<b>!</b>		
14	75.8	1320	1010			
15	75.8	2000	1018	<b>[</b> ]		
16	75.6	760	1017	<b>l</b> )		
17	75.6	1520	1018			
18	76.0	1740	1013			
19	75.1	1580	1015	8.13	0.528	1.54
20	76.0	1800	1015	H		
21	75.7	1140	1017	11		
22	75.7	1440	1013	ו		
28	76.0	2060	1011	1)		
24	76.2	2360	1011			
25	75.6	2380	1011			
26	75.8	2200	1013	8.68	0.555	1.63
27	77.2	1200	1014			
28	75.5	920	1021	[]		
29	75.9	1240	1022	Ι, Ι		
30	75.4	700	1026		•	
Dec. 1	75.7	1480	1017			
2	75.9	1160	1018		0.050	1 ,,,,
8	75.5	1480	1009	8.16	0.656	1.74
4	75.6	1860	1015			
5	76.2	1480	1012			
6	76.5	1460	1018			
7	75.7	800	1024	5.81		
8	75.6	880	1023	10.61	0.746	1.97
9	76.0	1040	1010	10.01		
10	76.0	1840	1016	12.91	J	

174 PHYSIOLOGICAL ECONOMY IN NUTRITION

FRITZ.

	Body	Urine.						
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>6</sub> .		
1903	kilos	0.0.		grams	gram	grams		
Dec. 11	75.6	1240	1017	10.04	h			
12	75.4	1740	1018	8.31	0.746	1.97		
13	75.4	1400	1018	7.72	daily av.	daily av.		
14	74.5	1260	1020	1)	ł			
15	74.9	1040	1013	11				
16	75.0	1890	1019	11				
17	75.2	1220	1015	8.18	0.624	1.65		
18	75.0	1520	1016	daily				
19	75.2	1380	1017	average				
20	75.0	890	1020	Įį.	1			
21	74.8	1315	1018					
22	74.8	880	1016	6.07	• • • •	• • •		
23	74.6	1185	1022	- 40	0.504	İ		
24	74.6	1596	1006	7.42	0.584	• • • •		
25	74.6	1800	1012	[]		,		
26	75.0 75.0	1090 1520	1025 1022		İ			
27 28	75.0 74.0	1150	1022	א	1			
28 29	74.0	1250	1018	11				
30	74.4	1610	1020			ł		
30 81	74.5	1025	1024	11		ĺ		
-	14.0	1020	1024	7.27	0.592	1.41		
1904. Jan. 1	74.2	1620	1010	11				
2	73.6	1990	1017					
8	78.7	1086	1029					
4	74.0	2070	1011	15				
اَة	74.1	1820	1021	11				
6	78.9	1690	1015	11	1			
7	74.0			8.06	0.650	1.73		
8	78.4	1940	1020					
9	78.6	1560	1012	11				
10	74.0	2200	1015	IJ				
11	74.0	1800	1019	10.29				
12	73.7	810	1023	8.99	1.100			
18	74.0	680	1024	6.49	<b>Q.691</b>			
14	78.8	1230	1019	10.26	0.998			
15	73.6	1030	1024	7.97	0.730			
16	78.9	1045	1015	5.20	0.466			
17	73.9	1910	1014	9.40	0.678			
18	78.0	1060	1017	7.01	0.631	1.23		

FRITZ.

	Body-	Urine.						
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>2</sub> .		
1904	kilos	c.c.		grams	gram	grama		
Jan. 19	78.0	1510	1020	1)				
20	73.4	865	1028	<b>     </b>				
21	78. <del>4</del>	1410	1020	7.01	0.681	1.23		
22	73.0	610	1025	daily	daily	daily		
28	78.0	1630	1020	average	average	average		
24	72.4	1125	1018	)	u.o.ugo			
25	72.6	2400	1011	1)				
26	72.8	2100	1010	<b>}</b>				
27	72.6	795	1018					
28	78.0	1425	1019	8.18	0.722	1.64		
29	73.0	1770	1013					
30	73.2	1910	1015					
81	73.0	2180	1016	P				
Feb. 1	73.2	2075	1015	l)				
2	78.1	2280 2860	1012	11				
8	73.0	2200	1023	704	0.532			
4	78.4	1600	1012 1019	7.24	0.002	1.74		
5	73.0 73.8	1745	1019	H				
6	78.0	745	1019	11	i			
7	78.5	2280	1025	K				
8 9	78.2	2150	1012	11				
10	73.1	1985	1018	11				
11	73.4	2240	1010	9.02	0.699			
12	73.0	2020	1016	1	0.000			
13	73.6	1800	1016	11				
13	73.0	1355	1021					
15	78.0	1280	1024	6				
16	73.5	2440	1014	11				
17	78.4	1880	1022	[]				
18	73.6	1840	1020	8.48	0.682	١		
19	73.6	2840	1016	11		••••		
20	73.6	2480	1010	11				
21	73.4	850	1024	11				
22	73.3	1985	1019	li				
28	78.4	1585	1011	11				
24	73.5	1940	1015	7.00	0.799			
25	78.5	1520	1017	7.68	U. 199	• • • •		
26	73.8	1030	1017	11				
27	78.0	2270	1015	J				

	Pod-			Urine.		
Date.	Body- Weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P2O8.
1904	kilos	c.c.		grams	gram	grams
Feb. 28	73.0	1660	1017	7.68	0.799	
29	72.8	1165	1020	6.22	n l	
Mar. 1	73.2	1240	1020	7.44		
2	73.2	915	1022	3.96		
3	73.2	1740	1015	7.05	0.500	
4	72.8			i	daily	
5	72.4	1670	1019	8.71	average	
6	72.6	1245	1020	4.78	J - I	
7	72.2	1900	1018	11		
8	72.6	1250	1018			
9	72.0	1660	1018	]]		
10	72.6	1900	1017	7.96	0.657	
11	72.5	2000	1015	daily		
12	72.6	2130	1015	average		
13	72.4	2430	1016	ال		
14	72.8	1100	1020	i i		
15	78.0	1950	1012			
16	73.2	2010	1011			
17	72.8	1790	1018	6.94	0.511	
18	72.6	1920	1014			
19	73.2	2150	1008			
20	73.0	1855	1020	11		
21	78.0	1670	1018	lí		
22	73.2	1640	1014	11		
23	78.0	2490	1012			
24	73.0	1840	1010	69.6	0.685	
25	73.0	1340	1016	55.5	0.000	
26	72.8	2040	1018	11		
27	72.6	2320	1011	]]		
28	73.0	2020		ľ I		
29	72.8	1600	1014	5.09	h l	
80	72.9	1480	1012	7.10		
81	72.8	1960	1016	5.18	0.478	
Apr. 1	72.6	2080	1013	5.74	5.2.5	
Apr. 1	72.5	2070	1010	3.85		
8	72.6	1920	1018	6.68	• • • •	•••
Daily aver Nov. 2	rage from	1566	1016	7.84	0.649	1.58

PHYSIOLOGICAL ECONOMY IN NUTRITION 177 cohn.

	Rody-	Urine.						
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P3O8-		
1903	kilos	c.c.	_	grams	gram	grams		
Nov. 15	65.0	1140	1019	11.22				
16	65.1	840	1024	l) l		i		
17	<b>6</b> 5.6	1280	1024	]]				
18	65.5	1420	1022	11 1				
19	65.1	1500	1020	10.86	0.605	1.60		
20	<b>6</b> 5.9	1540	1021	daily	daily	daily		
21	66.2	1840	1012	average	average	average		
22	65. <b>6</b>	840	1026					
23	<b>65.5</b>	1040	1018	]				
24	65.9	1260	1021					
25	66.1	1500	1016	]]				
26	65.8	800	1027	8.85	0.542	1.26		
27	65.3	840	1022					
28	65. <b>4</b>	940	1016					
29	65.7	1200	1023	J)				
30	65.6	1480	1016	in l				
Dec. 1	64.8	740	1024	1				
2	65.8	600	1026	11				
. 3	64.8	920	1023	9.89	0.621	1.51		
4	65.0	940	1020	[]				
5	64.8	680	1029	11				
6	64.8	1460	1016	J				
7	64.9	940	1027	lı l				
8	65.0	900	1023					
9	65.0	1040	1018	11				
10	64.9	960	1021	8.60	0.392	1.32		
11	64.0	580	1083	[]				
12	64.8	860	1026					
18	64.0	660	1028	ן ון				
14	64.0	1060	1021	)				
15	63.9	880	1019					
16	63.9	470	1081					
17	68.9	760	1026	} 7.29	0.424	1.29		
18	64.3	1180	1020	11 l				
19	64.4	1760	1011					
20	64.0	1100	1017	ן ו				
21	64.4	615	1025	l) l				
22	64.0	1050	1018	7.50	0.424			
28	64.0	860	1019	''''	V. ZAFA	•••		
24	64.2	1420	1019	'				

178 PHYSIOLOGICAL ECONOMY IN NUTRITION COHN.

50-4-	Body-	Urine.						
Date.	Body- weight,	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P3O8.		
1908	kilos	c.c.		grams	gram	grams		
Dec. 25	64.2	1690	1017	1		_		
26	64.0	740	1026	7.50	0.424			
27	63.9	1370	1017	daily av.	daily av.			
28	63.7	690	1025	lı i	•			
29	63.6	1065	1024	[]				
30	64.0	960	1024	11				
81	64.0	700	1026					
1904				7.70	0.358	0.90		
Jan. 1	63.4	1630	1017			daily		
2	63.5	460	1080			average		
8	63.0	970	1031	[]				
4	68.4	1365	1012	<b>b</b>				
5	63.5	980	1021	11				
6	64.0	1175	1022	11				
7	64.1	1250	1017	8.03	0.498	1.12		
8	64.0	1500	1018	11 1				
9	63.8	920	1027	11 1				
10	63.3	1250	1019	l) l				
11	63.8	880	1022	8.08				
12	62.8	510	1026	6.49	0.510			
18	68.0	900	1024	9.10	0.679			
14	62.8	630	1026	7.38	0.528			
15	62.4	950	1025	8.04	0.497			
16	62.6	1300	1018	8.58	0.473			
17	62.4	905	1026	7.44	0.481			
18	62.0	885	1026	la l				
19	62.7	1510	1020	]]				
20	63.0	980	1023	11				
21	62.4	935	1026	8.97	0.552	1.43		
22	62.6	1610	1019					
<b>2</b> 3	62.7	970	1024					
24	62.2	1275	1020	ן ען				
25	68.0	1600	1017	h l				
26	62.3	770	1025					
27	62.2	940	1028					
28	62.4	980	1021	7.34	0.539	1.80		
29	62.0	705	1025					
80	62.9	890	1024					
81	63.4	1560	1020	μ				
Feb. 1	63.5	1040	1028	8.15	0.532	1.87		

COHN.

2.	Body-	Urine.						
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P308.		
1904	kilos	c.c.		grams	gram	grams		
Feb. 2	63.4	960	1026	h i				
8	63.0	1500	1022					
4	68.2	1480	1023	1 0.15	0.500			
5	68.6	1890	1018	8.15	0.532	1.87		
6	63.0	780	1025	daily	daily	daily		
7	68.0	920	1026	average	average	average		
8	63.3	935	1027	h				
9	64.0	1625	1016					
10	63.5	1615	1017	11				
11	64.0	1325	1020	8.00	0.522			
12	64.0	1275	1021					
13	64.0	770	1024	11				
14	68.4	940	1023	]]				
15	68.5	1300	1027	1				
16	68.1	1430	1020	11				
17	68.6	1280	1024	[]				
18	63.8	870	1025	8.59	0.510			
19	68.2	1250	1026					
20	63.1	1250	1017	11 1				
21	68.5	900	1016					
22	63.3	1345	1026	1				
23	63.5	1185	1019					
24	68.8	1560	1020					
25	63.4	1200	1015	8.45	0.683			
26	63.0	750	1029	11 1				
27	63.5	1140	1019	11 1				
28	63.4	1220	1020	]] ]				
29	63.6	1160	1023	9.74				
Mar. 1	63.5	900	1025	6.86				
2	63.5	1030	1020	7.29				
8	62.7	920	1021	6.40	0.480			
4	62.7	1225	1014	5.59				
5	62.5	1170	1026	9.55				
6	62.6	940	1025	6.77				
7	62.5	900	1026	h l				
8	63.0	1280	1020					
9	62.7	1260	1020	8.41	0.608			
10	62.5	980	1019	0.21	0.000	• • • •		
11	62.4	1150	1024	11				
12	62.4	830	1027	ן		l i		
L	<u> </u>	L	<u> </u>	1				

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>5</sub> .
1904	kilos	c.c.		grams	gram	grams
Mar. 18	62.4	1500	1020	8.41	0.608	
14	62.0	825	1025	h		
15	62.5	1200	1017	H		
16	62.5	1490	1018			
17	62.5	1145	1021	7.59	0.488	
18	63.7	975	1021	daily	daily	
19	68.0	825	1028	average	average	
20	68.5	1450	1020	J i	_	
21	63.6	1480	1018	h		
22	68.0	1100	1025	11 :		
23	62.6	1050	1023	11		
24	62.7	1050	1025	7.74	0.654	
25	62.0	900	1027	11		
26	62.4	750	1025	11		
27	62.6	1580	1016	ĮJ		
28 -	62.0	1060	1023	6.61	1	
29	62.4	1460	1020	6.48		
30	62.4	1020	1023	6.36	0.899	
81	62.6	1730	1022	7.47		
<b>A</b> pr. 1	62.4	895	1022	4.35	J	
2	62.4	1465	1014	5.11	• • • •	
3	62.6	1165	1023	9.37	•••	•••
•	erage from		4000			4.05
Nov. 2	2	1092	1022	8.05	0.519	1.28

LOEWENTHAL.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>3</sub> .
1903	kilos	c.c.		grams	gram	grams
Oct. 4	60.1	960	1025	15.78	0.606	1.78
5	60.1	780	1025	10.90		
6	60.3	820	1024	11.23	0.576	0.96
7	60.0	980	1022	14.99		
8	60.0	1080	1019	11.34	0.851	1.55
9	60.0	990	1025	18.07		• • •
10	60.0	1120	1021	12.87		
11	59.8	980	1019	10.88	0.389	1.05
12	60.4	1580	1022	17.06		
13	60.2	1380	1017	13.00	0.516	2.13
14	61.0	640	1026	7.41		
15	61.2	1220	1026	15.66	0.674	1.68
16	60.8	1060	1024	15.01		
17	60.5	1180	1021	18.63		
18	60.4	1580	1016	12.70	0.520	1.59
19	60.0	1850	1022	17.82		
20	59.8	1250	1019	1		
21	60.0	1120	1024	11		1
22	60.4	1060	1025	13.72	0.489	1.95
28	61.0	1800	1020	10.12	0.409	1
24	61.2	1320	1021	daily av.	daily av.	daily av.
25	61.4	1620	1015	Į)		
26	62.4	1160	1024	1)		
27	62.6	1520	1021	11		
28	62.6	860	1022			
29	63.4	660	1024	9.80	0.480	1.40
80	62.5	1200	1023	11 .		
81	62.2	920		11		1
Nov. 1	62.4	1050	1019	)		
2	62.6	1830	1023	[]		
8	62.0	1020	1022			1
4	61.8	920	1020			
5	62.4	580	1029	8.28	0.401	1.60
6	62.4	1400	1016	11		
7	62.0	780	1022			
8	61.9	760	1021	ĺj		
9	61.8	840	1031	[]	1	
10	60.5	440	1028	8.28	0.390	1.34
11	61.0	1060	1028		V.500	
12	61.2	880	1027	J		

LOEWENTHAL.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P2O5.
1903	kilos	o.c.		grams	gram	grams
Nov. 18	61.4	320	1028	h		
14	61.4	660	1023	8.28	0.890	1.84
15	61.2	1020	1028	J daily av.	daily av.	daily av.
16	61.4	960	1025	1)		
17	61.0	<b>6</b> 80	1028	11		
18	61.0	940	1023	11		
19	60.9	880	1024	8.62	0.349	1.80
20	61.5	1540	1018			i
21	61.7	1220	1022			·
22	61.2	1120	1022	[]		ł
28	61.1	1440	1014	]]		
24	60.2	840	1024			
25	60.8	680	1026			
26	60.8	900	1022	8.36	0.392	1.43
27	60.5	860	1025	11		
28	60.8	800	1028			
29	60.3	1040	1025	ĺ,		
30	60.8	1040	1021			
Dec. 1	60.4	1120	1017	11		
2	60.0	660	1027	7.50	0.400	1 200
8	59.9	900	1022	7.59	0.408	1.26
4	59.5	1120 400	1025 1028			İ
5 6	59.0 58.6	480	1028			
7	58.2	960	1033	9.03		
8	58.0	800	1027	8.78	1	
ŷ	58.6	700	1028	10.04		
10	60.0	1000	1028	9.06	0.879	1.49
11	60.0	800	1025	8.71	0.0.0	1.70
12	60.0	680	1032	7.51		
18	59.5	620	1030	7.74	]	
14	59.5	940	1022	J	-	
15	59.0	920	1019	11		
16	59.6	690	1027			
17	59.3	800	1020	6.90	0.408	1.14
18	59.7	1145	1020		*	
19	59.7	1110	1016			
20	59.0	840	1020			
21	59.0	775	1025	)	0.000	
22	58.4	510	1027	6.29	0.275	• • •

LOEWENTHAL.

Date.	Body-	Urine.					
Date.	weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>5</sub> .	
1903	kilos	c.c.		grams	gram	grams	
Dec. 23	59.2	700	1026	l)			
24	59.0	890	1015			1	
25	58.8	900	1027	6.29	0.275		
26	59.0	980	1015	daily	daily		
27	58.4	990	1018	average	average		
28	58.5	672	1028	h l			
29	58.4	796	1023	11 1		[	
30	59.0	870	1025				
31	58.8	920	1027	6.92	0.515	1.17	
1904		l				daily	
Jan. 1	58.7	1840	1022	)) l		average	
2	58.2	1232	1019				
8	58.6	842	1025	į į		i	
4	58.8	1030	1017	1			
5	<b>58.6</b>	1020	1026				
6	58.8	890	1024	11			
7	59.0	1350	1016	7.27	0.344	0.97	
8	57.7	900	1026	11 1			
9	58.0	785	1033				
10	57.9	760	1024	l)			
11	58.0	795	1021	6.68			
12	58.2	1110	1017	7.88	0.383	• • •	
18	58.1	1190	1015	7.64	0.879	• • •	
14	57.0	620	1027	6.21	0.415		
15	57.2	825	1028	8.18	0.409		
16	57.8	1100	1019	7.92	0.305		
17	57.5	1185	1021	7.26	0.338		
18	57.8	630	1019		• • •	• • •	
19	57.5	525	1030	h l			
20	58.0	1050	1026	[[ ]			
21	57.7	790	1027	6.84	0.321	0.99	
22	57.7	850	1030	••••	4.44	VV	
28	57.4	605	1024	[[			
24	<b>57.4</b>	800	1080	ן ען			
25	57.9	1050	1020	ו ו			
26	58.0	850	1026	[]			
27	58.2	1270	1018	6.68	0.312	1.14	
28	59.0	1255	1013				
29	58.0	615	1028	]] ]			
30	58.0	1025	1029				

#### LOEWENTHAL.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> .
1904	kilos	c.c.		grams	gram	grams
Jan 31	58.6	1120	1026	])		
Feb. 1	58.4	10.25	1029			
2	58.7	1300	1023	11		
8	58.4	1170	1024	}		
4	58. <del>4</del>	1850	1029	7.27	0.411	1.57
5	59.0	1250	1026	daily	dail <del>y</del>	daily
6	58.8	1160	1019	average	average	average
7	58.4	825	1029	1)	1	
8	58.5	830	1028			
9	58.5	1095	1025	·		
10	59.0	1140	1027	[.}		
11	<b>5</b> 9. <b>2</b>	1830	1020	7.61	0.416	
12	58.5	1020	1029			
13	59.0	1075	1025	زا		
14	59.0	1030	1024	]]		
15	58.3	1150	1027			
16	58.4	1270	1024	[]		
17	58.6	1490	1020	}		
18	59.0	1060	1027	8.00	0.336	
19	58.8	620	1026	11		
20	58.9	930	1029	Į		
21	59.1	885	1029	]]		
22	59.3	1320	1023			
28	59.4	1490	1016			
24	59.4	1195	1019	}		
25	59.4	2100	1011	6.84	0.449	• • •
26	58.4	860	1023			
27	58.4	955	1027	l)		
28	58.4	1045	1027			
29	58.2	1000	1027	7.38	1	
Mar. 1	58.6	1040	1024	7.05		
2	58.5	880	1028	7.07	0.040	
8	58.7	800	1026	7.85	0.348	• • •
4 5	58.5 58.6	965	1026	7.85		
6	58.7	730 1170	1022 1027	4.29		
7	58.3	920	1027	8.07	'	
	58.9	920 970	1026			
8 9	58.8 58.8	940	1026	6.97	0.388	
10	58.9	1600	1026	}	0.000	

#### LOEWENTHAL.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> .
1904	kilos	6.0.		grams	grams	grams
Mar. 11	59.0	1290	1020	h		
12	59.0	820	1029	6 97	0.388	
18	58.9	990	1026	daily av.	daily av.	
14	59.5	1620	1019	h	, i	
15	59.0	1250	1028	]] [		
16	59.0	1360	1018			
17	59.1	1160	1021	7.34	0.265	
18	59.0	1450	1019	41		
19	59.1	1650	1018			
20	59.0	1350	1021	]]		
21	59.0	1110	1024	1)		
22	58.8	1250	1021	11		
28	58.6	1150	1021	11		
24	58.6	760	1025	6.37	0.427	
25	59.0	1100	1025	11		
26	58.6	990	1028	11		
27	58.5	1150	1026	ן ו		
28	58.6	1175	1024	6 27	1	
29	59.0	1295	1025	7.54		
80	59.0	1570	1018	7.07	0.341	• • •
31	59.2	1120	1024	5.24		
Apr. 1	59.0	1260	1022	6.43	J	
2	59.0	1550	1020	6.33		• • • •
8	59.0	1710	1022	9.85	• • •	• • •
Daily aver Nov. 2	rage from	1007	1024	783.	0.372	1.28

ZOOMAN.

	Rode			Urine.		
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P3O8.
1903	kilos	c.c.		grams	gram	grams
Oct. 4	54.0	540	1026	8.88	0.434	0.56
5	54.0	940	1027	15.28		
6	58.6	1140	1024	16.42	0.730	2.21
7	54.5	1640	1016	17.52		
8	54.4	980	1017	10.10	0 275	1.21
9	54.4	1160	1022	15.80		
10	54.8	1080	1019	15.24		
11	54.5	1840	1014	18.65	0.450	1.78
12	54.8	1080	1022	11.78		
18	55.4	1140	1022	13.48	0.839	1.70
14	55.7	1020	1018	9.85	• • •	
15	55.2	1560	1021	20.40	0.779	2.92
· 16	56.0	1240	1024	17.33		
17	<b>55.6</b>	1070	1025	16.11	• • •	
18	55.6	1780	1017	20.86	0.786	2.26
19	55.4	1370	1020	18.57		
20	55.6	1290	1024	]]		
21	55.6	1120	1025	11		
22	55.8	1140	1023	15.88	0.683	1.87
23	55.9	1120	1025	daily	daily	daily
24	56.2	1600	1020	average	average	average
25	56.5	1620	1015	,	a recuge	go
26	56.6	1140	1024	11		
27	56.6	1770	1015			
28	56.6	960	1021			
29	57.8	1120	1019	11.70	0.583	1.56
80	56.8	1160	1019			
81	57.0	820	1010			
Nov. 1	56.9	1730	1016	l)		
2	56.5	750	1026	[]		
8	57.1	1380	1018			
4	57.1	1040	1019	1000	0.504	
5	56.6	900	1018	10.01	0.564	1.39
6	56.6	1220	1016			
7	56.2	640	1026	]]		
8	55.8	900	1020	ľ.		
9	56.1	1100	1021			
10	56.2	680	1024	8.76	0.512	1.33
11	56.4	660 1820	1016			
12	56.7	1620	1024	J		

ZOOMAN.

Date.	Body- weight.			Urine.		
Date.	weight.	Volume. 24 hours.	Bp. Gr.	Nitrogen.	Uric Acid.	P2O8.
1903	kilos	c.c.		grams	gram	grams
Nov. 18	<b>5</b> 6.7	420	1024	h		
14	56.0	920	1029	8.76	0.512	1.33
15	56.0	1540	1015	daily av.	daily av.	daily av.
16	55.7	1080	1022	h i		,
17	54.4	700	1027			
18	54.6	700	1025	[]		
19	54.8	700	1025	7.79	0.398	1.02
20	55.4	940	1021	11 1		
21	55.2	740	1025	11		
22	54.7	720	1023	J		
28	<b>54</b> .5	740	1014	h l		
24	<b>54</b> .0	860	1025	11 1		
25	54.8	620	1025	11 1		
26	54.4	980	1028	7.44	0.420	1.06
27	54.5	940	1024	]		İ
28	54.8	460	1026	11		
29	54.4	1000	1018			1
80	54.0	980	1022	lı l		
Dec. 1	54.2	1120	1020	11		
2	54.2	940	1017	11		
8	54.0	1200	1018	10.26	0.494	1.45
4	<b>54</b> .0	1080	1024	11		
5	<b>53.9</b>	640	1081	11 1		
6	54.0	960	1026	j		
7	54.1	880	1028	lı l		
8	54.6	680	1029	9.79		
9	54.6	680	1030	11 1		i
10	55.0	1220	1019	10.15	0.423	1.89
11	54.7	920	1028	]] ]		
12	55.1	780	1028	8.38		
18	54.7	1140	1020	J 8.89		
14	54.4	840	1021	7.46		
15	54.0	1200	1016	8.71		
16	54.0	860	1021	[[		
17	54.5	1140	1018	7.66	0.889	1.20
18	54.0	820	1022	11 1		
19	54.1	980	1018			
20	54.0	885	1023	ן יין		
21	54.0	660	1027	8.26	0.401	0.78
22	53.5	570	1025	ا		

ZOOMAN.

Date.	Body-			Urine.		
	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P2O8-
1903	kilos	c.c.		grams	gram	grams
Dec. 28	54.0	770	1028	b l		_
24	54.0	920	1025	11 1		
25	54.3	1820	1018	8.26	0.401	0.78
26	54.7	1075	1020	daily	daily	daily
27	54.5	1240	1018	average	average	average
28	54.1	880	1022	li I		_
29	54.0	1080	1019	[ ]		
30	54.3	775	1029	]]		
81	54.4	1010	1023	8.00	0.440	0.88
1904				lf		
Jan. 1	54.0	900	1022	11 1		
2	58.6	625	1026			
8	58.9	730	1026	]]		
4	53.0	1030	1019	li l		
5	58.4	1100	1022	11 1		
6	54.0	980	1020	11	0.879	1.15
7	54.0	860	1022	7.24		
8	54.0	960	1026			
9	53.5	695	1025	11		
10	54.0	980	1016	IJ		
11	53.5	930	1020	8.98	0.632	
12	53.8	1030	1023	11.31	0.657	
13	58.6	980	1024	11.63	0.445	
14	53.5	1150	1017	9.38	0.448	
15	53.0	700	1080	8.44	0.442	
16	53.4	920	1020	8.89	0.881	
17	58.5	1270	1019	8.99		
18	58.5	1180	1020	1)		
19	58.6	755	1026			
20	53.6	1190	1015		0.428	1.27
21	53.0	800	1030	7.78		1
22	58.2	660	1030			
23	53.4	770	1025	11		
24	58. <del>4</del>	1085	1017	ן (		
25	53.6	1250	1018	]]		1
26	540	1140	1024			
27	54.2	1170	1018	6.87	0.302	1.18
28	54.6	1120	1018	11	V.002	
29	54.5	690	1026			
30	54.5	1020	1025	ا		
<u> </u>	1	<u> </u>	L		<u> </u>	<u> </u>

ZOOMAN.

_	Body.			Urine.		
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>5</sub> .
1904	kilos	c.c.		grams	gram	grams.
Jan. 31	54.8	1320	1019	6.87	0.802	1.18
Feb. 1	54.8	1060	1026	l)		
2	54.5	1170	1019	11		
8	54.5	1350	1022			
4	<b>54</b> .3	1280	1022	8.15	0.405	1.31
5	<b>54</b> .0	1015	1025	daily	daily	daily
6	54.6	1610	1014	average	average	average
7	<b>54</b> .3	1010	1028	j		
. 8	54.6	1065	1026	h		
` 9	55.0	925	1027			
10	55.0	1195	1023	11		
11	55.0	880	1024	8.37	0.497	
12	55.0	960	1016			
13	55.2	1565	1020	11		
14	55.0	1415	1028	IJ		
15	55.0	1060	1029	1		
16	54.8	1180	1025			
17	55.0	1910	1019			
18	55.2	1260	1020	9.84	0.408	
19	55.0	970	1027			
20	54.4	670	1031	[]		
21	54.7	1070	1022	IJ		
22	54.4	760	1025	i)		
28	54.6	1225	1020	11		
24	55.2	1050	1026			
25	55.2	730	1027	7.38	0.559	• • •
26	55.4	1145	1019	11		
27	55.0	990	1021	] [		
28	54.9	985	1026	)		
29	55.0	765	1027	7.02	}	
Mar. 1	55.8	810	1024	6.66		
2	55.0	880	1027	7.26	0.400	
8	54.8	1020	1020	7.59	0.401	• • •
4	54.5	955	1025	7.51		
5	54.0 54.0	1035 880	1020	7.08		
6 7	54.7	885	1027 1023	6.81	J.	
8	55.0	970	1023			
, s	54.6	940	1025	8.24	0.428	
10	54.7	1460	1020	0.21	V.740	•••
	UX.	1400	1022	<u> </u>		

# 190 PHYSIOLOGICAL ECONOMY IN NUTRITION ZOOMAN.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> .
1904	kilos	0.0.		grams	gram	grams
Mar. 11	55.0	1340	1017	li l		
12	54.7	1180	1023	8.24	0.428	
18	55.0	1040	1021	daily av.	daily av.	
14	55.0	1270	1016	1		
15	54.8	900	1028	11		
16	55.0	1145	1019			
17	55.0	1155	1018	7.90	0.455	
18	53.0	1480	1016	''''		
19	55.2	1355	1023	11		
20	54.6	1000	1023			
21	54.7	750	1026	lí l		
22	55.0	1270	1018	11		
23	55.0	1090	1016	11		
24	55.0	1080	1025	7.57	0.586	
25	54.8	1080	1028			
26	55.0	980	1021	11		
27	55.2	1110	1020	[J		
28	55.2	1270	1019	7.47		
29	55.2	1160	1024	7.80		
30	55.0	1140	1018	5.40		
31	55.3	1340	1024	8.04		
Apr. 1	55.0	1480	1019	8.44		
2	55.1	1300	1023	7.10		
8	55.0	1445	1018	8.15		
Daily ave Nov. 2	rage from	1008	1022	8.25	0.457	1.19

BATES.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>5</sub> .
1903.	kilos	c.c.		grams	gram	grams
Oct. 4	72.7	870	1033	14.56	0.651	2.18
5	72.7	1120	1022	18.13		
6	72.4	760	1027	9.44	0.489	1.38
7	72.8	920	1022	12.86	• • •	
8	72.0	630	1029	11.11	0.415	1.29
9	72.0	1030	1025			
10	72.0	1230	1022	14.46		
11	72.3	740	1024	10.79	0.397	1.44
12	72.8	860	1024	10.54		• • •
18	72.1	1040	1025	14.23	0.769	2.46
14	72.8	1010	1027	13.70		
15	72.4	1800	1021	14.43	0.510	1.88
16	72.4	1040	1025	15.85	• • •	• • •
17	72.4	1440	1020	16.83	• • •	• • •
18	72.2	1420	1021	18.66	0.657	2.51
19	71.8	780	1026	13.14	• • •	
20	71.8	1060	1027	lı l		
21	72.0	1020	1025	11 1		
22	72.5	1080	1024	14.21	0.517	1.26
28	72.4	1200	1024	daily	daily	daily
24	72.4	1100	1025	average	average	average
25	72.4	1030	1024	ا عقد ا		
26	72.4	1060			• • •	• • •
27	72.6	1020	1022			
28	72.4	720	1027	11		
29	72.4	400	1032	9.20	0.409	1.27
80	72.8	720	1027			
81	72.8	720	• • •			
Nov. 1	72.8	820	1027	ין		
2	72.0	960	1026	n		
8	72.4	920	1025			
4	71.9	700	1025	000	0.40	1.58
5	71.6	900	1012	9.01	0.487	1.05
6	71.5	700	1029	]]		
7	70.5	600	1028 1031	11		
8	70.0	680	1031	ן ין		
9	70.8	660		1)		
10	70.6	660 560	1028 1030	7.97	0.460	1.20
11	70.6	740	1030			
12	71.0	/190	1027	<b> </b>		

BATES.

Date.	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Urio Acid.	P <sub>2</sub> O <sub>2</sub> .
1903.	kilos	c.c.		grams	gram	grams
Nov. 13	71.0	660	1027	h		
14	70.6	480	1032	7.97	0.460	1.20
15	70.4	800	1026	daily av.	daily av.	daily av.
16	70.8	860	1025	h		
17	70.0	860	1028			
18	69.8	540	1025		•	
19	69.4	700	1026	6.94	0.374	1.04
20	70.8	1080	1023			
21	70.2	640	1025			
22	69.8	720	1027	IJ		
23	69.1	720	1025	li l		
24	68.7	600	1028			
25	68.7	620	1028	11		
26	69.0	840	1081	8.04	0.894	1.56
27	68.6	820	1026	]		
28	67.9	800	1022	11		
29	69.4	700	1027	[]		
80	69.0	780	1025	h		
Dec. 1	68.5	760	1026	11		
2	68.7	640	1027			
8	68.1	940	1027	8.24	0.893	1.87
, 4	69.0	540	1023			
5	68.0	1360	1011			
6	69.0	880	1027	Į) į		
7	67.9	960	1026	10.14	)	
8	68.2	700	1026	7.85	1	
9	68.0	800	1025	10.45		
10	68.7	700	1027	8.74	0.392	1.57
11	67.9	780	1028	9.27		
12	68.0	820	1029	8.85		
13	67.6	860	1028	9.44	,	]
14	67.8	700	1029	7.36		
15	67.4	540	1028	6.51		
16	67.4	640	1032	1		
17	67.6	680	1027	11	0.007	
18	68.8	1000	1023	7.61	0.867	1.36
19	68.3	800	1028	11		
20	67.0	660	1027	ון		
21	67.0	595	1027	6.77	0045	
22	66.2	560	1028	], "	0.345	

BATES.

	Body-			Urine.		
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>8</sub> .
1903	kilos	c.c.		grams	gram	grams
Dec. 28	66.5	660	1027	h		i ' i
24	66.4	610	1021			1
25	65.6	1057	1021	6.77	0.345	
26	66.0	700	1029	daily av.	daily av.	ł į
27	66.2	755	1026	Į)		
28	65.8	740	1024	1		
29	65.5	710	1026			
80	66.0	900	1025			i
81	65.7	710	1028	6.53	0.269	0.80
1904		1		1 0.00	0.200	daily av.
Jan. 1	65.7	690	1027			Lany av.
2	65.8	1015	1010	11		i I
8	65.0	850	1027	[]		
4	65.0	730	1024	1)		1
5	65.1	835	1022	11 1		
6	65.6	1045	1027	11		i I
7	66.2	900	1024	8.54	0.405	1.18
8	65.4	1080	1025			
9	66.0	840	1028			
10	66.0	965	1025	]]		
11	65.8	700	1026	6.80		
12	65.5	740	1025	7.46	0.365	
18	65.2	630	1027	7.03	0.899	
14	64.5	550	1028	7.18	0.857	
15	64.6	680	1022	8.04	0.434	
16	65.0	620	1029	7.66	0.470	
17	65.0	925	1024	7.38	0.347	
18	65.0	590	1082	l)		
19	65.0	650	1028			1
20	64.8	575	1029			
. 21	64.4	735	1028	6.69	0.356	0.80
22	64.8	640	1029			]
23	64.6	760	1026			
24	64.8	870	1023	J		
25	64.0	740	1027	h l		
26	64.5	975	1024	11		( )
27	68.8	910	1024	11	1	
28	64.0	600	1030	8.54	0.859	1.27
29	68.8	1300	1017	11		1
80	64.0	1185	1024	ן י		1

#### BATES.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>2</sub> .
1904	kilos	c.c.		grams	gram	grams
Jan. 31	64.0	1460	1018	8.54	0.859	1.27
Feb. 1	64.0	910	1027	h		
2	63.9	1250	1021	11		
8	63.5	1080	1026	11		
4	68.5	970	1024	9.19	0.862	1.07
5	63.6	1250	1024	daily	daily	daily
6	64.0	1260	1018	average	average	average
7	64.0	785	1026			
8	64.6	1235	1022	9.94		
9	64.3	1460	1018	8.95	• • •	• • • •
10	64.2					
11	64.3					
12	64.8		• • •			
Daily aver Oct. 27	rage from	805	1025	8.08	0.387	1.23

DAVIS.

	Body- weight,			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P2O8-
1903	kilos	c.c.		grams	gram	grams
Oct. 4	59.8	2030	1014	15.15	0.672	1.99
5	59.8	1160	1020	10.99		
6	59.0	850	1024	9.88	0.503	1.26
7	59.1	1860	1012	18.96		
8	59.3	960	1021	9.38	0.898	1.47
9	58.8	900	1027	11.56		
10	58.3	1660	1017	17.28		
11	58.7	1460	1019	16.02	0.724	2.27
12	58.5	1060	1022	11.82		
18	59.4	1920	1014	10.59	0.689	2.58
14	59.4	1460	1022	18.14		
15	58.9	1320	1021	13.88	0.498	1.53
16	58.9	1900	1015	17.10		
17	59.0	1410	1019	14.55		
18	59.5	1650	1017	15.25	0.647	1.92
19	59.1	1120	1022	13.37		
20	58.2	1150	1020	h		
21	58.2	980	1021			·
22	59.1	1820	1022	11.56	0.480	1.50
28	59.4	1820	1016	daily	daily	daily
24	58.6	1160	1022	average	average	average
25	59.2	1800	1018	Average	average	m.c.ago
26	59.0	860	1024	lı l		
27	59.2	1240	1024	11		
28	59.4	1440	1014	11		
29	59.5	820	1022	} 9.20	0.474	1.85
30	59.4	1000	1017			
81	59.2	920		11		
Nov. 1	59.4	1120	1015	) .		1
2	58.7	840	1026	1		
8	59.1	1290	1018			
4	59.0	680	1025			
5	58.5	305	1027	9.07	0.474	1.48
6	58.4	700	1023			
7	58.7	780	1022			
8	58.5	1720	1011	ון		
9	58.2	1820	1015	h		
10	58.9	920	1020	8.38	0.422	1.88
11	59.8	1640	1014		V.222	1,00
12	58.9	780	1024	,		

DAVIS.

	Rody			Urine.		
Date.	Body- weight.	Volume. 24 bours.	Sp. Gr.	Nitrogen.	Uric Acid.	P309.
1904	kilos	c.c.		grams	gram	grams
Nov. 18	59.0	940	1025	l)		
14	59.0	1180	1016	8.38	0.422	1.38
15	59.0	1900	1018	daily av.	daily av.	daily av.
16	58.8	1080	1022	lı l		
17	59.0	1080	1018			
18	58.4	1500	1017	11		
19	58.2	1260	1018	9.60	0.398	1.50
20	58.1	1220	1025	11		
21	58.7	1120	1022			
22	58.7	1100	1021	j j		
23	58.6	1140	1017	1		
24	58.4	1860	1017	11		
25	58.8	1700	1010			
26	58.2	1120	1023	9.00	0.447	1.52
27	59.4	1620	1017			
28	58.0	700	1027			
29	58.7	1660	1025	]]		
30	58.5	800	1025	l)		
Dec. 1	58.0	700	1020			
2	<b>57.4</b>	680	1028	11 1		
3	57.0	640	1028	11.41	0.417	2.12
4	57.0	940	1032			
5	56.2	660	1032	11 1		
6	56.0	800	1029	])		
7	56.0	800	1026	li l		
8	56.1	780	1030	]] ]		
9	57.0	820	1022	11 1		
10	58.0	500	1025	8.91	0.412	1.48
11	57.1	580	1029	11 1		
12	57.3	820	1031	$\Pi$		
13	57.8	580	1027	h		
14	57.5	740	1027	h l		
15	57.5	1680	1015	]]		
16	57.9	900	1021	11 1		
17	<b>57.4</b>	780	1013	7.96	0.418	1.50
18	56.9	800	1026	11		
19	<b>57.4</b>	730	1025			
. 20	57. <b>4</b>	800	1028	li li	•	
21	57.4	1310	1017	7.35	0.819	
. 22	57.0	1870	1014	ر الم	0.010	

### PHYSIOLOGICAL ECONOMY IN NUTRITION 197 DAVIS.

		Body-			Urine.		
Date.	•	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>5</sub> .
1903		kilos	c.c.		grams	gram	grams
Dec.	23	57.0	980	1020	h	_	1
1 :	24	57.0	865	1016	11		
	25	57.4	1300	1017	7.35	0.319	
1 :	26	58.0	1390	1017	daily	daily	
	27	57.8	1340	1018	average	average	
	28	57.6	875	1019	1		
! :	29	57.0	680	1020	li l		
	80	56.8	685	1029	11		
	81	56.7	815	1026		0.410	
1904	1				6.83	0.413	0.99
Jan.	1	57.4	1150	1015	11		daily
ì	2	57.9	1610	1023			average
1	8	58.6	1120	1023	ן ען		
	4	58.3	1350	1011	li l		
	5	57.8	1255	1017	11 1		
	6	58.0	925	1022	11 1		
	7	58.6	1120	1016	7.04	0.372	0.95
	8	58.4	1350	1020	[]		
	9	<b>5</b> 7. <b>7</b>	890	1026	]]		
	10	58.6	1510	1012			
	11	57.9				• • •	
:	12	57.2	835	1017	8.82	0.408	• • •
Daily average from Oct. 28		1045	1021	8.61	0.414	1.42	

Any elaborate discussion of the individual results, tabulated in the foregoing tables, seems hardly necessary. To any one interested in the details of the work, a study of the tables themselves will give the necessary information. It may be well, however, to emphasize at once a few of the fundamental points most striking in character which bear upon the main problem. Take, for instance, the case of Oakman, where the data are recorded for every day from October 4, 1903, to April 4, 1904. The degree of proteid metabolism, as indicated by the excretion of nitrogen through the kidneys, is here shown for each day of the six months. Up to November 3, practically for a month, there was not observable any very noticeable change in the rate of proteid metabolism. but commencing with November 4, the nitrogen excretion dropped very rapidly, and as one glances through the daily records for month after month, it is seen that the daily nitrogen output through the kidneys fell to 6 to 8 grams per day, with a daily average excretion of 7.42 grams of nitrogen, as figured from October 21 to April 4.

Further, it will be noted that while the body-weight gradually declined during the first three months, falling from 66.7 kilos down to 62.3 kilos on January 18, from that date on to the close of the experiment the body-weight was practically constant. Here, then, we see, under the use of a prescribed diet quite sufficient in amount to satisfy the cravings of the appetite, a lowering of proteid metabolism equal to that obtained by the individuals of the preceding group. Again, if the body-weight of Oakman is placed at 64 kilos, as representing the average between the initial body-weight and the weight during the last three months, it will be found that the output of metabolized nitrogen per kilo of body-weight amounted to 0.116 gram; not widely different from similar data obtained with men of the preceding group.

Moreover, these results obtained with Oakman are practically duplicated by every other member of this group of soldiers detailed from the United States Army. No exception whatever is to be seen, but every man shows the same lowered



FRITZ.

Photograph taken at the close of the experiment.

,

proteid metabolism, with practical maintenance of bodyweight, with complete satisfying of the appetite, and without loss of health, strength, or vigor, as will be enlarged upon later.

While in the case of Oakman, the nitrogen excreted was determined by an analysis made each day, in the case of the other men of this group the data are given mainly for weekly periods; the results being expressed, however, in the average daily amount for each seven days' period. By comparing the figures for the daily excretion of nitrogen with the daily dietary, it is easy to trace out the influence of the changes in diet on the extent of nitrogen metabolism.

In considering the changes in the rate of proteid metabolism shown by all the members of this group, it will suffice for present purposes to deal mainly with the average results.

Compare now the average daily output of nitrogen through the urine from November 1 until April 4—a period of five months—of each of these men.

## AVERAGE DAILY OUTPUT OF NITROGEN.

```
Steltz
                 7.13
Broyles . . .
                 7.26
Loewenthal. . .
                 7.38
Oakman . . .
                 7.42
Fritz . . . . 7.84 = 7.80 grams Nitrogen as
Cohn. . . . 8.05
                          the grand average.
Coffman . . . 8.17
Zooman . . . 8.25
Sliney. . . . .
                 8.39
Henderson .
                 8.91
                 8.08 — Oct. 27-Feb. 11.
Bates . . .
         . . 8.61 — Oct. 26-Jan. 12.
Davis. .
```

The figures given show an astonishingly low proteid metabolism for the five months' period; 7.5 grams of nitrogen in the day's urine correspond to 46.8 grams of proteid metabolized, while 8.5 grams of nitrogen mean the metabolism of 53.1

grams of proteid. In other words, all of these men during a period of five months practically averaged a daily output of nitrogen through the kidneys corresponding to the metabolism of less than one-half the 105 grams of absorbable proteid called for by the so-called standard diets. Further, close scrutiny of the results in the individual tables shows that during many weeks much lower results were obtained than is indicated by the general averages.

Just here emphasis should be laid upon one fact, doubtless of primary importance in any modification of the rate or extent of metabolism in any individual, viz., that in this experiment the daily diet was prescribed, thereby taking from the individual freedom of choice in the selection of food. The writer has no question in his own mind that an intelligent choice of food, coupled with the satisfying of a natural or acquired appetite in moderation, will lead to better results than any system of prescription as to what shall be eaten each day and in what quantity. Still, this is the method necessarily made use of in the present experiment, the writer having prescribed the character and amount of each meal throughout the entire six months' period, with due regard, of course, to the expressed likes and dislikes of the men.

One feature in the diet compelled by the circumstances of the case also needs to be referred to, namely, the necessity of keeping the men thoroughly satisfied, so that there should be no feeling of hunger, no craving for food. With a person interested in the experiment and desirous of ascertaining the effect of a low nitrogen intake, there would naturally be a willingness to endure, if necessary, for a time some personal discomfort; but with this detail of the United States Army it could not be expected that the men would remain satisfied if they were compelled to undergo the pangs of hunger even for a day or two. Consequently, it was necessary in prescribing the daily diet to see that the quantity of the food was such as to completely satisfy the appetite. This necessitated the use of considerable bulky food of low fuel, and low nitrogen, value. In this way only was it feasible to reduce the

nitrogen intake, as well as the fuel value of the food, to the desired level. Practically during the entire six months' period, with the possible exception of a few days, the men were given sufficient food to completely satisfy their appetites. Throughout the entire period of the experiment, the men all manifested a keen appetite and utilized their food to good advantage, with establishment of the nitrogen metabolism indicated by the foregoing results.

Still confining our attention to the average results regarding nitrogen excretion, we may ask the question, what was the excretion of metabolized nitrogen per kilo of body-weight in the different individuals? This is easily calculated and the data are shown in the following table, in which the figures standing for body-weight are either the data for the month of April, 1904, or else the average between the October and April weights, as in those cases where the body-weight fell off during the experiment.

			F	lody-weight.	November-April.	Body-weight.
Fritz				kilos 74.0	7.84	974m 0.106
Oakman .				64.0	7.42	0.116
Bates				68.0	8.08	0.118
Morris		•		59.0	7.03	0.119
Broyles .				60.0	7.26	0.120
Henderson				71.0	8.91	0.125
Loewenthal			•	59.0	7.38	0.125
Cohn			•	63.5	8.05	0.126
Steltz		•		53.0	7.13	0.184
Sliney	•			60.0	8.39	0.138
Coffman .		•		58.0	8.17	0.140
Davis		•	•	<b>58.0</b>	8.61	0.148
Zooman .	•	•	•	55.0	8.25	0.150

Scrutiny of these results shows that the daily excretion of metabolized nitrogen ranged in this period of five months from 0.106 gram per kilo of body-weight up to 0.150 gram per kilo of body-weight. Since these men were on essentially the

same diet, it is obvious that there were some peculiarities, or personal idiosyncrasies, that tended to modify the exact extent of proteid metabolism, and in some cases at least constituted a more potent factor than body-weight in determining the rate of metabolism. This fact is clearly emphasized in the case of Morris, who, with a body-weight of 59 kilos, showed a proteid metabolism equivalent to only 7.03 grams of nitrogen per day, and Coffman, who, with a body-weight of 58 kilos, showed under the same conditions an average excretion of 8.17 grams of nitrogen per day.

What was the effect of this lowered proteid metabolism upon the bodily condition, especially body-weight? To answer this question the weights of the men are given in the following table, the one column of figures showing the body-weight at the commencement of the experiment, the other column showing the weight at the close of the experiment.

TABLE OF BODY-WEIGHTS.

				0	ctober, 1903.	April, 1904.
Steltz.					52.3	53.0
Zooman					<b>54.0</b>	<b>55.</b> 0
Coffman					59.1	58.0
Morris					<b>59.2</b>	59.0
Broyles					59.4	61.0
Loewent	hal				60.1	59.0
Sliney					61.3	60.6
Cohn .					65.0	62.6
Oakman					66.7	62.1
Henderso	n				71.3	71.0
Fritz .					76.0	72.6
Bates .			•		72.7	64.3 February
Davis.			•		59.3	57.2 January

Comparison of these figures shows that six of the men were practically of the same body-weight at the close of the experiment as at the beginning. Of the others, Oakman, Fritz, Cohn, and Bates lost somewhat. Bates, however, was quite stout, and

the work in the gymnasium as well as the change of diet tended to reduce his weight. In fact, with the possible exception of Oakman, the slight loss of weight these latter men underwent was a distinct gain to their bodily condition. The most significant feature, however, is to be found on scrutiny of the tables of results for each man, where is seen the body-weight for each day of the six months. Here it will be noticed that, as the experiment progressed, there was, as a rule, a tendency for the body-weight to increase; this was followed by a decline, after which the weight remained practically stationary. This is well illustrated in Oakman's case. Starting with a body-weight of 66.7 kilos on October 4, he reached 67.6 kilos on November 29, then declining in weight to 62.3 on January 18, after which he remained practically stationary, or varied only within narrow limits.

Finally, it should be said that the low proteid metabolism on which these men were apparently maintaining body-weight was not associated with any increase of non-nitrogenous food. At no time did the fuel value of the daily food reach 3000 calories; in fact, about January 12 the average fuel value of the food was only a little over 2000 calories per day.

#### FIRST NITROGEN BALANCE.

Early in January, the body-weight of the men having become practically stationary and the proteid metabolism, as measured by the output of metabolized nitrogen, having been reduced to a low level, an attempt was made to see if the men were truly in a condition of nitrogenous equilibrium. A six-day period was used, during which all the food for each meal was carefully analyzed, the nitrogen in every sample of food being determined in duplicate by the Kjeldahl-Gunning process. The fæces of the period were separated by lamp-black, given at appropriate times in gelatin capsules, and all necessary precautions observed to insure an accurate nitrogen balance. The accompanying tables of results give all the necessary data.

## 204 PHYSIOLOGICAL ECONOMY IN NUTRITION

The average nitrogen intake per day was 8.23 grams, and the average fuel value of the food per day was 2078 calories. These are the figures for Oakman. With some of the other men there were slight variations from these figures, as shown in the tables - owing to slight variations in diet. Without exception, all the men showed a minus balance, indicating that on this diet the body was losing nitrogen. In other words, the diet as a whole was insufficient for the needs of the body in every case. Whether this was due to lack of sufficient proteid or to an insufficient amount of fat and carbohydrate, i. e., a too low fuel value, cannot be said definitely, but there is every reason to believe that the amount of non-nitrogenous food was insufficient to furnish the requisite fuel value, and consequently body proteid was drawn upon to supply the The loss of nitrogen to the body per day varied as a rule from 0.5 to 2.0 grams. In one case only, viz., Broyles, was there a close approach to a balance. In his case there was a minus balance of only 0.12 gram of nitrogen per day.

The average daily output of nitrogen through the urine for this six-days period (Oakman's case) was 7.52 grams.



FRITZ

Photograph taken at the close of the experiment.



## Tuesday, January 12, 1904.

Breakfast. — Fried Indian-meal 100 grams, syrup 50 grams, one cup coffee 350 grams, bread 50 grams, butter 15 grams.

Dinner. — Boiled macaroni 250 grams, stewed tomato 250 grams, bread 75 grams, one cup coffee 350 grams.

Supper. — Potato chips 100 grams, fried bacon 25 grams, bread 75 grams, jam 75 grams, one cup tea 350 grams.

Food.		Grams.	Per cent Nitrog	en. Total Nitrogen.
Fried Indian-mea	al	. 100	× 0.38	= 0.380 gram.
Syrup		. 60	× 0.051	= 0.025
Coffee (breakfast		. 850	× 0.075	= 0.262
Butter		. 15	× 0.10	= 0.015
Macaroni		. 250	× 0.53	= 1.825
Stewed tomato		. 250	× 0.15	= 0.375
Bread 50	) + 75 + 75 =	= 200	× 1.48	= 2.960
Potato chips .		. 100	× 1.09	= 1.090
Bacon		. 25	× 3.13	= 0.780
Jam		. 75	× 0.089	= 0.029
Tea		. 350	× 0.067	= 0.234
Coffee (dinner)		. 850	× 0.091	= 0.318
	Total nitroger			7.798 grams. 6.000

Fuel value of the food . . . . 2404 calories.

•

#### OAKMAN.

## Wednesday, January 13, 1904.

Breakfast.—Boiled rice 150 grams, milk 125 grams, sugar 30 grams, butter 10 grams, bread 30 grams, one cup coffee 350 grams.

Dinner.— Hamburg steak with plenty of bread, fat, and onions chopped together 150 grams, boiled potato 200 grams, apple sauce 200 grams, bread 75 grams, one cup coffee 850 grams.

Supper. — Fried rice 100 grams, syrup 50 grams, one cup tea 350 grams, bread 50 grams, butter 15 grams.

Food.						(	Grams.	Per	cei	at Nitrog	en.	Total Nitrogen.
Boiled rice							150	×		0.36	==	0.540 grams.
Milk							125	×		0.55	=	0.687
Sugar							30	×		0.00	=	0.000
Butter (breakfast	j						10	×		0.10	=	0.010
Bread " .	•						30	×		1.66	=	0.498
Coffee "							850	×		0.066	=	0.281
Hamburg steak,	tc.						150	×		2.80	=	4.200
Potato							200	×		0.29	=	0.580
Apple sauce .							200	×		0.067	=	0.134
Bread (dinner)							75	×		1.66	=	1.245
Coffee (dinner)							350	×		0.076	=	0.266
Fried rice							100	×		0.50	=	0.500
Syrup							50	×		0.051	=	0.025
Tea			• .				350	×		0.066	=	0.281
Bread (supper)							50	×		1.66	=	0.830
Butter							15	×		0.10	=	0.015
	To	ta	l n	itr	oge	n i	n food					9.992 grams.
	To	ota	l n	itr	oge	n	in urine					. 7.830

Fuel value of the food . . . . 2133 calories.

## Thursday, January 14, 1904.

Breakfast. — Boiled hominy 150 grams, milk 125 grams, sugar 30 grams, butter 10 grams, bread 30 grams, one cup coffee 350 grams.

Dinner. — Split pea soup (thick) 200 grams, bread 75 grams, mashed potato 100 grams, pickles 30 grams, one cup coffee 350 grams, pie 120 grams.

Supper. — Suet pudding 150 grams, apple sauce 125 grams, crackers 25 grams, one cup tea 350 grams.

Food.						•	lrams.	Per c	ent Nitroge	n. To	tal Nitrogen.
Boiled hominy .							150	X	0.21	=	0.815 gram.
Milk							125	×	0.52	=	0.650
Sugar							30	×	0.00	=	0.000
Butter							10	×	0.10	=	0.010
Bread (breakfast)							30	×	1.59	=	0.477
Coffee (breakfast)	}						350	×	0.096	=	0.886
Split pea soup .							200	×	0.69	=	1.880
Bread (dinner)							75	×	1.59	=	1.192
Mashed potato							100	×	0.81	=	0.310
Pickles							30	×	0.054	=	0.016
Coffee (dinner)							850	×	0.075	=	0.262
Pie							120	×	0.50	=	0.600
Suet pudding .							150	×	0.78	=	1.170
Apple sauce .							125	×	0.058	=	0.072
Crackers							25	×	1.61	=	0.402
Tea							350	X	0.063	=	0.220
	T	ota	d n	itr	og	en	in food				7.412 grams.
					_		in urin				8.290

Fuel value of the food . . . . 2000 calories.

## Friday, January 15, 1904.

Breakfast. — Wheat griddle cakes 200 grams, syrup 50 grams, one cup coffee 350 grams.

Dinner. — Codfish-balls (4 parts potato, 1 part fish, fried in pork fat) 150 grams, stewed tomato 200 grams, bread 75 grams, one cup coffee 350 grams, apple pie 95 grams.

Supper. — Apple fritters 200 grams, stewed prunes 125 grams, bread 50 grams, butter 15 grams, one cup tea 350 grams.

Food.							Grams.	Pe	r ce	nt Nitrog	en. I	Total Nitrogen.
Wheat griddle ca	ke	:8					200	×		0.78	=	1.560 grams.
Syrup							50	×		0.051	=	0.025
Coffee (breakfast	)						350	×		0.075	=	0.262
Codfish-balls, etc.							150	×		1.77	=	2.655
Stewed tomato							200	×		0.14	=	0.280
Bread (dinner)					•		75	×		1.59	=	1.192
Coffee (dinner)							350	×		0.075	=	0.262
Apple pie							95	×		0.40	=	0.380
Apple fritters .							200	×		0.40	=	0.800
Stewed prunes							125	×		0.15	=	0.187
Bread (supper)							50	X		1.59	=	0.795
Butter							15	×		0.10	=	0.015
Tea							350	×		0.042	=	0.147
	T	ota	d n	itr	oge	n	in food					8.560 grams.
	T	ota	l n	itr	oge	n i	in urine					7.140

Fuel value of the food . . . . 2030 calories.

## Saturday, January 16, 1904.

Breakfast. — Soft oat-meal 150 grams, milk 100 grams, sugar 30 grams, bread 30 grams, butter 10 grams, one cup coffee 350 grams.

Dinner. — Baked macaroni with a little cheese 200 grams, stewed tomato 200 grams, bread 50 grams, tapioca-peach pudding 150 grams, one cup coffee 350 grams.

Supper. — French fried potato 100 grams, fried bacon 20 grams, bread 75 grams, jam 75 grams, one cup tea 350 grams.

Food.						Grams.	Per	cent Nitro	gen.	Total Nitrogen.
Soft oat-meal .						150	×	0.27	=	0.405 grams.
Milk						100	×	0.44	=	0.440
Sugar						30	X	0.00	=	0.000
Bread (breakfast)						30	×	1.61	=	0.483
Butter						10	×	0.10	=	0.010
Coffee (breakfast	) .					850	×	0.075	=	0.262
Baked macaroni,	etc.					200	×	0.72	=	1.440
Stewed tomatoes						200	×	0.15	=	0.300
Bread (dinner)						50	×	1.61	=	0.805
Tapioca-peach pu	ddin	g				150	×	0.087	=	0.130
Coffee (dinner)						350	×	0.093	=	0.325
French fried pota	to.					100	×	0.47	=	0.470
Fried bacon .						20	X	8.15	=	0.630
Bread (supper)						75	X	1.61	=	1.207
Jam						75	X	0.039	=	0.029
Tea						850	×	0.099	=	0.346
	Tota	al n	itr	oge	en	in food				. 7.282 grams.
•	Tota	d n	itr	oge	n	in urine			. •	. 8.230

Fuel value of the food . . . . 1824 calories.

## Sunday, January 17, 1904.

Breakfast. — Boiled Indian-meal 125 grams, milk 125 grams, sugar 30 grams, butter 10 grams, bread 30 grams, one cup coffee 350 grams.

Dinner. — Bean soup (thick) 200 grams, bread 75 grams, mashed potato 100 grams, pickles 25 grams, one cup coffee 350 grams, custard pie 105 grams. Supper. — Crackers 50 grams, butter 15 grams, stewed prunes 125 grams, sponge cake 100 grams, one cup tea 350 grams.

Food.						Grams.	Per	cent Nitro	gen.	Total Nitrogen.
Boiled Indian-meal						125	×	0.24	=	0.300 grams.
Milk						125	X	0.50	=	0.625
Sugar						80	X	0.00	=	0.000
Butter (breakfast)						10	X	0.10	=	0.010
Bread (breakfast)						30	X	1.61	=	0.488
Coffee (breakfast)						850	X	0.087	=	0.804
Bean soup						200	X	0.86	=	1.720
Bread (dinner) .						75	X	1.61	=	1.207
Mashed potato .						100	X	0.28	=	0.280
Pickles						25	X	0.054	= -	0.018
Coffee (dinner) .						350	X	0.081	=	0.288
Custard pie						105	X	0.88	=	0.924
Crackers						50	X	1.61	=	0.805
Butter (supper) .						15	×	0.10	=	0.015
Stewed prunes .						125	×	0.17	=	0.212
Sponge cake						100	X	1.00	=	1.000
Tea						850	X	0.048	=	0.168
T	ote	ıl n	itr	oge	en i	in food				. 8.349 grams.
T	ots	d n	itr	oge	n	in urine	•			. 8.140

Fuel value of the food . . . . 2081 calories.

## NITROGEN BALANCE. - Oakman.

	Nitrogen	Ot	ıtput.
	Taken in.	Nitrogen in Urine.	Weight of Faces (dry).
Jan. 12	7.793 grams.	6.00 grams	
18	9.992	7.83	•••
14	7.412	8.29	5.0 grams.
15	8.560	7.14	76.0
16	7.282	8.23	13.5
17	8.849	8.14	9.5
			104.0 grams contain
			7.39% N.
	49.888	45.18 +	7.685 grams nitrogen.
	49.388 grams	nitrogen. 52.	315 grams nitrogen.

Nitrogen balance for six days = -3.427 grams. Nitrogen balance per day = -0.571 gram.

## Average Intake.

Calories per day . . . . . . . . . . . 2078. Nitrogen per day . . . . . . . 8.23 grams.

#### NITROGEN BALANCE. - Loewenthal.

	Nitrogen	Out	rat.
		rogen in Urine.	Weight of Frees (dry).
Jan. 12	7.793 grams.	7.33 grams	
18	9.992	7.64	21 grams.
14	7.888	6.21	28
15	8.560	8.18	52
16	7.282	7.92	. 37
17	8.349	7.26	<u>11</u>
	•		147 grams contain
			6.97 % N.
	49.864	44.54 +	- 10.24 grams nitrogen.
	49.364 grams n	itrogen. 54.	78 grams nitrogen.

Nitrogen balance for six days = -5.416 grams. Nitrogen balance per day = -0.902 gram.

Daily diet same as Oakman's, except that a smaller amount of coffee was taken at breakfast on the 14th. Nitrogen correction made accordingly.



COFFMAN STELTZ
Photographs taken at the close of the experiment.



## NITROGEN BALANCE. - Steltz.

	Nitrogen	Outpu	at.
		ogen in Urine. \	Weight of Freces (dry).
Jan. 12	7.798 grams.	4.61 grams.	15.0 grams
13	9.463	7.90	34.0
14	7.412	4.99	21.0
15	8.560	9.05	81.5
16	7.282	7.56	33.0
17	8.849	8.55	18.0
			152.5 grams contain
			6,52% N.
	48.859	42.66 +	9.94 grams nitrogen.
	48.859 grams n	itrogen. 52.6	0 grams nitrogen.

Nitrogen balance for six days = -3.741 grams. Nitrogen balance per day = -0.628 gram.

Daily diet same as Oakman's, except on January 18, when 8 grams of boiled rice were uneaten at breakfast and 100 grams of fried rice at supper. Correction in nitrogen-content made accordingly.

## 214 PHYSIOLOGICAL ECONOMY IN NUTRITION

## NITROGEN BALANCE. - Bates.

	Nitrogen Taken in. Nitr	Out ogen in Urine.	put. Weight of Froces (dry).
Jan. 12	7.706 grams.	7.46 gram	5
18	9.916	7.08	• • •
14	7.875	7.18	36 grams.
15	8.439	8.04	30
16	7.226	7.66	84
17	8.349	7.88	34
			184 grams contain
	•		7.17 % N.
	49.011	44.70 -	- 9.61 grams nitrogen.
	49.011 grams ni	trogen. 54	.81 grams nitrogen.

Nitrogen balance for six days = -5.299 grams. Nitrogen balance per day = -0.888 gram.

## Daily diet same as Oakman's, with the following exceptions:

Jan. 12	Supper:	8 8	gram	s Potato chips,	uneaten.
13	Dinner:	118	"	Apple sauce,	"
14	"	50	"	Coffee,	"
15	Supper:	81	"	Prunes,	**
16	Breakfast:	75	"	Coffee,	44

## NITROGEN BALANCE. - Coffman.

Nitrogen	Output.		
Taken in. N	itrogen in Urine.	Weight of Fraces (dry)	
7.793 grams.	8.82 grams.	• • •	
9.992	8.28	•••	
7.412	8.30	57.00 grams.	
8.560	7.91	41.25	
7.282	7.82	47.00	
8.349	7.44	21.50	
		166.75 grams contain	
		6.66% N	
49.888	48.07 +	11.10 grams nitrogen.	
49.888 grams 1	nitrogen. 59.1	7 grams nitrogen.	
	7.793 grams. 9.992 7.412 8.560 7.282 8.349	Taken in.       Nitrogen in Urine.         7.793 grams.       8.82 grams.         9.992       8.28         7.412       8.30         8.560       7.91         7.282       7.32         8.349       7.44	

Nitrogen balance for six days = -9.782 grams. Nitrogen balance per day = -1.630 grams.

Daily diet same as Oakman's.

# NITROGEN BALANCE. - Fritz.

	Nitrogen	Output.		
	Taken in.	Nitrogen in Urine.	Weight of Fraces (dry).	
Jan. 12	7.793 grams	. 8.99 grams.	• • •	
13	9.992	6.49	• • •	
14	7.412	10.26	64.0 grams.	
15	8.560	7.97	29.5	
16	7.282	5.20	62.0	
17	8.186	9.40	31.0	
			186.5 grams contain	
			6.49% N.	
	49.225	48.31 +	12.10 grams nitrogen	
	49.225 gram	nitrogen. 60	.41 grams nitrogen.	

Nitrogen balance for six days = -11.185 grams. Nitrogen balance per day = -1.864 grams.

Daily diet same as Oakman's, except that on the 17th inst. a portion of the prunes was uneaten. Correction made accordingly.

## NITROGEN BALANCE. -Henderson.

	Nitrogen	Output.			
	Taken in.	Nitrogen in Urine.	Weight o	f Fraces (dry).	
Jan. 12	7.639 grams	. 7.68 gran	ns		
13	9.794	8.22	•••		
14	7.487	8.24	• • •		
15	8.560	7.76	45 g	rams.	
16	7.068	7.56	26		
17	8.045	7.87	22		
		***************************************	98	grams contain	
				6.45% N.	
	48.598	47.88	+ 6.0	0 grams nitrogen.	
	48.593 grams nitrogen.		58.38 grams nitrogen.		
	Nitrogen balance f	or six days :	= <b>-4</b> .787	grams.	
	Nitrogen balance p	er day :	= -0.789	gram.	

Daily diet same as Oakman's, except that on several days smaller amounts of coffee and tea were taken. Corrections made accordingly.

## NITROGEN BALANCE. - Morris.

	Nitrogen	Output.	Output.			
	Taken in. Nitro	gen in Urine. We	eight of Faces (dry).			
Jan. 12	7.255 grams.	4.19 grams.	• • •			
18	9.573	7.92	•••			
14	7.825	7.91	4.75 grams.			
15	8.538	7.44	88.00			
16	7.282	7.88	71.00			
17	8.349	5.28	53.00			
			166.75 grams contain			
	•		6.45% N.			
	48.822	40.12 +	10.75 grams nitrogen.			
	48.322 grams ni	rogen. 50.87	grams nitrogen.			

Nitrogen balance for six days = -2.548 grams. Nitrogen balance per day = -0.424 gram.

Daily diet same as Oakman's except that on the first three days smaller amounts of tea and coffee were taken. Corrections made accordingly.

## NITROGEN BALANCE. - Zooman.

	Nitrogen	Output.	
	Taken in. Nitro	gen in Urine. Weig	tht of Feeces (dry).
Jan. 12	7.466 grams.	11.31 grams.	• • •
18	9.992	11.68	• • •
14	7.412	9.38	• • •
15	8.560	8.44	27.5 grams.
16	7.282	8.89	28.5
17	8.849	8.99	13.0
			69.0 grams contain
			6.46% N.
	49.061	58.64 +	4.45 grams nitrogen.
	49.061 grams ni	trogen. 63.09	grams nitrogen.

Nitrogen balance for six days = -14.029 grams. Nitrogen balance per day = -2.338 gram.

Daily diet same as Oakman's, except that on January 12th 30 grams potato chips were not eaten. Nitrogen intake corrected accordingly.

## NITROGEN BALANCE. - Sliney.

	Nitrogen Taken in. Nitro	Output gen in Urine. V	t. Veight of Fæces (dry).
Jan. 12	6.501 grams.	5.94 grams	22.5 grams.
13	9.975	8.44	26.0
14	7.388	7.42	22.5
15	8.560	7.89	24.0
16	7.282	7.28	13.0
17	8.849	8.15	15.0
			128.0 grams contain
			6.72 % N.
	48.055	45.07	- 8.26 grams nitrogen.
	48.055 grams n	itrogen. 53	.32 grams nitrogen.

Nitrogen balance for six days -5.275 grams. -0.879 gram. Nitrogen balance per day

Daily diet same as Oakman's, except that on the 12th 190 grams stewed tomatoes and 190 grams macaroni were uneaten; on the 13th at breakfast 25 grams of coffee were left, and on the 14th at breakfast 25 grams coffee were not taken. Corrections in intake of nitrogen made accordingly.



COFFMAN STELTZ

Photographs taken at the close of the experiment.



## NITROGEN BALANCE. - Cohn.

	Nitrogen Taken in.	Out; Nitrogen in Urine.	out. Weight of Fraces (dry).
Jan. 12	7.798 grams.	6.49 grams.	• • •
18	9.957	9.10	29.0 grams.
14	7.412	7.33	8.5
15	8.820	8.04	44.0
16	7.282	8.58	19.0
17	8.849	7.44	5.5
			106.0 grams contain
			6.48% N.
	49.118	46.98 +	6.87 grams nitrogen
	49.113 grams n	itrogen. 58.85	grams nitrogen.

Nitrogen balance for six days = -4.737 grams. Nitrogen balance per day = -0.789 gram.

Daily diet same as Oakman's, except that on the 15th inst. at supper 60 grams fritters were uneaten, and on the 13th at breakfast 30 grams coffee, and at dinner 22 grams apple sauce were uneaten. Corrections made accordingly.

## NITROGEN BALANCE. - Broyles.

	Nitrogen Taken in.	Out <sub>j</sub> Nitrogen in Urine.	out. Weight of Fraces (dry).
Jan. 12	7.475 grams.	5.88 grams.	•••
13	9.495	6.99	• • •
14	6.814	7.47	• • •
15	8.086	9.67	85 grams.
16	6.695	7.65	14
17	7.762	5.28	20 69 grams contain
			6.61 % N.
	46.277	42.44 +	4.56 grams nitrogen.
	46.277 grams nitrogen.		grams nitrogen.

Nitrogen balance for six days = -0.723 gram. Nitrogen balance per day = -0.120 gram.

Daily diet same as Oakman's, except that each day coffee was omitted from breakfast and dinner. Corrections made accordingly.

## SECOND NITROGEN BALANCE.

Commencing February 29, a second nitrogen balance was attempted with very different results. This balance period extended through seven days and was characterized by an average daily intake of 9.50 grams of nitrogen, an increase of 1.8 grams of nitrogen per day over the preceding period, together with an average fuel value of the food equal to 2500 calories per day.

The daily diet with its content of nitrogen, etc., is detailed under Oakman. Any variations (only slight ones) from this diet are indicated on the individual balance-sheets, corrections being made in the nitrogen intake.

In all of the men, with one exception, a distinct plus balance is seen, showing that under the conditions of this experiment the men were all storing up nitrogen. The plus balance per day varied from 0.132 gram to 1.231 grams of nitrogen.

The average daily output of nitrogen through the urine for this seven days' period (Oakman's case) was 7.24 grams, being 0.28 gram less per day than in the preceding period. Sliney, the one man who failed to show a positive balance, was apparently losing nitrogen at the rate of 0.48 gram per day during this period.

All of the individual data are shown in the accompanying tables of results:

## Monday February 29, 1904.

Breakfast. — Boiled rice 175 grams, milk 125 grams, sugar 25 grams, baked potato 150 grams, one cup coffee 350 grams, butter 10 grams.

Dinner. — Baked spaghetti 250 grams, mashed potato 250 grams, bread 75 grams, boiled tomato 150 grams, apple pie 112 grams, one cup coffee 350 grams.

Supper. — Biscuit 175 grams, fried bacon 20 grams, fried sweet potato 150 grams, butter 20 grams, one cup tea 350 grams.

Food.	G <sub>1</sub>	rams. Per ce	nt Nitrogen.	Total Nitrogen.
Boiled rice	1	75 ×	0.84 =	0.595 grams.
Milk	1	125 ×	0.55 =	0.687
Sugar		25 ×	0.00 =	0.000
Baked potato	1	150 ×	0.84 =	0.510
Coffee (breakfast)	8	350 ×	0.082 =	0.287
Butter	10 + 20 =	80 ×	0.16 =	0.480
Spaghetti	2	250 ×	0.73 =	1.825
Mashed potato		250 ×	0.30 =	0.750
Bread		75 ×	1.61 =	1.207
Tomato	:	150 🗙	0.16 =	0.240
Apple pie	1	112 ×	0.46 =	0.515
Biscuit	1	175 ×	1.21 =	2.117
Fried bacon		20 ×	3.80 =	0.760
Fried sweet potato	1	150 ×	0.22 =	0.330
Tea		850 ×	0.06 =	0.210
Coffee (dinner) .		850 ×	0.11 =	0.385
To	otal nitrogen in	food		10.466 grams.
To	otal nitrogen in	urine		6.880

Fuel value of the food . . . . 2670 calories.

## Tuesday, March 1, 1904.

- Breakfast.—Fried rice 150 grams, syrup 50 grams, baked potato 150 grams, butter 10 grams, one cup coffee 350 grams.
- Dinner. Thick pea soup 250 grams, boiled onions 150 grams, boiled sweet potato 150 grams, bread 75 grams, butter 20 grams, one cup coffee 350 grams.
- Supper. Celery-lettuce-apple salad 120 grams, crackers 82 grams, American cheese 20 grams, Saratoga chips 79 grams, one cup tea 850 grams, rice custard 100 grams.

Food.	(	Frams.	Per cent Nitroge	n. Tota	d Nitrogen.
Fried rice ,		150	× 0.84	= 0.0	510 grams.
Syrup		50	× 0.00	= 0.0	000
Baked potato		150	× 0.33	= 0.4	195
Butter	10 + 20 =	30	× 0.16	= 0.0	<b>)48</b>
Coffee (breakfast)		350	× 0.042	= 0.3	147
Pea soup		250	× 0.54	= 1.3	350
Boiled onions		150	× 0.27	= 0.4	105
Boiled sweet potato		150	× 0.18	= 0.	195
Bread		75	× 1.65	= 1.5	237
Coffee (dinner) .		350	× 0.084	= 0.3	294
Salad		120	× 0.14	= 0.3	168
Crackers		82	× 1.50	= 0.	<b>480</b>
Cheese		20	× 8.92	= 0.	78 <del>4</del>
Saratoga chips .		79	× 1.22	= 0.9	963
Tea		350	× 0.054	= 0.	189
Rice custard		100	× 0.56	= 0.6	560
T	otal nitrogen i	n food .	. <b></b> .	7.5	325 grams.
$\mathbf{T}$	otal nitrogen i	n urine		7.	120

Fuel value of the food . . . . 2279 calories.

## Wednesday, March 2, 1904.

- Breakfast. Wheat griddle cakes 200 grams, syrup 50 grams, butter 10 grams, one cup coffee 350 grams, banana 75 grams.
- Dinner.—Boiled salt mackerel 25 grams, boiled potato 250 grams, boiled turnip 150 grams, bread 75 grams, one cup coffee 850 grams, apple sauce 150 grams.
- Supper. Chopped fresh cabbage with salt, pepper, and vinegar 100 grams, bread 75 grams, butter 20 grams, chocolate cake 150 grams, cranberry sauce 100 grams, one cup tea 850 grams.

Food.		Grams.	Per o	ent Nitroge	m. ?	Total Nitrogen.
Griddle cakes .		200	X	0.80	=	1.600 grams.
Syrup		. 50	×	0.00	=	0.000
Butter	10 + 20 =	30	×	0.16	=	0.048
Coffee (breakfast		350	×	0.066	=	0.281
Banana		75	×	0.20	=	0.150
Mackerel		25	×	3.92	=	0.980
Boiled potato .		250	×	0.27	=	0.675
Boiled turnip .		150	×	0.071	=	0.106
Bread	.75 + 75 =	150	×	1.68	=	2.445
Coffee (dinner)		850	×	0.066	=	0.281
Apple sauce .		150	×	0.048	=	0.072
Cabbage		. 100	×	0.28	=	0.280
Chocolate cake		. 150	×	0.95	=	1.425
Cranberry sauce		100	×	0.045	=	0.045
Tea		. 850	×	0.057	=	0.199
	Total nitrogen	in food				8.487 grams.
	Total nitroger					7.580

Fuel value of the food . . . . 2391 calories.

# Thursday, March 3, 1904.

Breakfast. — Boiled hominy 175 grams, milk 125 grams, sugar 25 grams, baked potato 150 grams, butter 10 grams, one cup coffee 350 grams.

Dinner.—Hamburg steak with much bread, fat, and onions 150 grams, boiled potato 250 grams, bread 75 grams, butter 10 grams, one cup coffee 350 grams.

Supper. — Tapioca-peach pudding 250 grams, bread 75 grams, jam 75 grams, butter 20 grams, one cup tea 350 grams.

Food.	Grams.	Per e	ent Nitro	gen.	Total Nitrogen.
Boiled hominy	. 175	×	0.19	=	0.830 grams.
Milk	. 125	×	0.40	=	0.500
Sugar	. 25	×	0.00	=	0.000
Baked potato	. 150	×	0.28	=	0.420
Butter $10 + 10 + 20 =$	= 40	×	0.16	=	0.064
Coffee (breakfast)	. 350	×	0.075	=	0.262
Hamburg steak	. 150	×	2.12	=	8.180
Boiled potato	. 250	×	0.27	=	0.675
Bread 75 + 75 =	= 150	×	1.48	=	2.220
Coffee (dinner)	. 350	×	0.093	=	0.325
Tapioca-peach pudding	. 250	×	0.22	=	0.550
Jam	. 75	×	0.034	=	0.025
Tea	. 850	×	0.057	=	0.199
Total nitroge	n in food	ı			8.750 grams.
Total nitroge	n in urin	e			6.850

Fuel value of the food . . . . 2375 calories.

# Friday, March 4, 1904.

Breakfast. — Fried hominy 150 grams, syrup 50 grams, baked potato 150 grams, one cup coffee 350 grams, butter 10 grams.

Dinner.—Codfish-balls (1 part fish, 4 parts potatoes, fried in pork fat) 150 grams, stewed tomato 200 grams, stewed potato 150 grams, bread 75 grams, one cup coffee 350 grams, apple pie 180 grams.

Supper. — French fried potato 200 grams, fried sausage 50 grams, butter 10 grams, bread 50 grams, stewed prunes 125 grams, sponge cake 35 grams, one cup tea 350 grams.

Food.		Grams.	Per cent Nitr	ogen.	Total Nitrogen.
Fried hominy .		150	× 0.82	=	0.480 grams.
Syrup		50	× 0.00	=	0.000
Baked potato .		150	× 0.37	=	0.555
Coffee (breakfast)	;)	850	× 0.099	=	0.846
Butter	10+10 =	20	× 0.16	=	0.032
Fish-balls		150	× 1.78	=	2.595
Stewed tomato		200	× 0.19	=	0.380
Bread	.75 + 50 =	125	× 1.54	=	1.925
Coffee (dinner)		350	× 0.069	=	0.241
Apple pie		130	× 0.88	=	0.494
French fried pota	ito	200	× 0.49	=	0.980
Sausage		50	$\times$ 2.75	=	1.375
Prunes		125	× 0.17	=	0.212
Sponge cake .		85	× 0.88	=	0.290
Tea		850	× 0.072	=	0.252
Stewed potato.		150	× 0.18	=	0.270
	Total nitrogen	in food			10.427 grams.
,	Total nitrogen	in urine			7.950

Fuel value of the food . . . . 2374 calories.

# Saturday, March 5, 1904.

Breakfast. — Boiled Indian-meal 200 grams, milk 125 grams, sugar 25 grams, one cup coffee 350 grams, fried sweet potato 150 grams, butter 10 grams.
 Dinner. — Tomato soup, thick with potatoes and onions boiled together 325 grams, bread 100 grams, scrambled eggs 50 grams, mashed potato 160 grams, one cup coffee 350.

Supper. — Bread pudding with raisins 250 grams, stewed peaches 150 grams, bacon 20 grams, French fried potato 150 grams, bread 50 grams, butter 10 grams, one cup tea 350 grams.

Food.	Grams.	Per o	ent Nitrog	en.	Total Nitrogen.
Boiled Indian-meal	200	×	0.20	=	0.400 gram.
Milk	125	X	0.52	=	0.650
Sugar	25	X	0.00	=	0.000
Coffee (breakfast)	850	X	0.10	=	0.850
Fried sweet potato	150	X	0.22	= '	0.830
Butter $10 + 10 =$	20	X	0.16	=	0.032
Tomato soup	825	X	0.10	=	0.325
<b>Bread</b> $100 + 50$	150	X	1.57	=	2.355
Scrambled eggs	50	×	2.22	=	1.110
Mashed potato	150	×	0.25	=	0.375
Coffee (dinner)	350	X	0.075	= .	0.262
Bread pudding	250	×	0.80	=	2.000
Stewed peaches	150	X	0.24	=	0.360
Bacon	20	X	4.00	=	0.800
French fried potato	150	X	0.56	=	0.840
Tea	850	X	0.084	=	0.294
Total nitrogen	in food		• .• •		10.488 grams.
Total nitrogen	in urine				6.100

Fuel value of the food . . . . 2302 calories.

# Sunday, March 6, 1904.

Breakfast.—Fried Indian-meal 150 grams, syrup 50 grams, sliced banana 100 grams, baked potato 150 grams, one cup-coffee 350 grams, butter 10 grams.\_

Dinner. — Corned beef 50 grams, boiled cabbage 200 grams, mashed potato 250 grams, bread 75 grams, one cup coffee 350 grams, fried rice 100 grams, jam 75 grams.

Supper. — Sponge cake 150 grams, apple sauce 150 grams, crackers 32 grams, butter 10 grams, one cup tea 350 grams, sardine 14 grams.

Food,						(	Grams.	Per c	ent Nitroge	m.	Total Nitrogen.
Fried Indian-mea	1						150	×	0.88	=	0.570 gram.
Syrup							50	×	0.00	=	0.000
Banana							100	X	0.19	=	0.190
Baked potato .							150	X	0.87	=	0.555
Coffee (breakfast	)						<b>350</b>	X	0.072	=	0.252
Butter		10	+ (	10	)	=	20	×	0.16	=	0.082
Corned beef .							50	X	5.24	=	2.620
Cabbage							200	X	0.34	=	0.680
Mashed potato							250	×	0.82	=	0.800
Bread							75	×	1.67	=	1.252
Coffee (dinner)							350	×	0.098	=	0.325
Fried rice							100	×	0.23	=	0.280
Jam							75	×	0.034	=	0.025
Sponge cake .							150	X	1.02	=	1.580
Apple sauce .	•						150	×	0.044	=	0.066
Crackers							82	×	1.50	=	0.480
Tea						•	850	×	0.054	=	0.189
Sardine							14	×	8.85	=	0.469
	T	ota	d n	itr	oge	n i	in food				10.265 grams.
	T	ota	d r	itr	og	en	in urine				7.960

Fuel value of the food . . . . 3173 calories.

### NITROGEN BALANCE. - Oakman.

	Nitrogen	Output.				
	Taken in.	Nitrogen in	Urine.	Weight of Faces * (dry).		
Feb. 29	10.466 grams.	6.88	grams.	• • •		
Mar. 1	7.825	7.42		• • •		
2	8.487	7.58		17 grams		
8	8.750	6.85		72		
4	10.427	7.95		89		
5	10.488	6.10		19		
6	10.265	7.96		<u>85</u>		
	•			182 grams contain		
				6.81% N.		
	66.703	50.74	+	12.894 grams nitrogen.		
	66.703 grams 1	nitrogen.	68.18	4 grams nitrogen		
Nitro	ogen balance for se	ven days	=	+3.569 grams.		
Nitro	ogen balance per de	ay .	=	+0.509 gram.		
	<b>A</b> v	erage Intake.				
Cal	ories per day			2509.		
Nita	rogen per day			9.50 grams.		

The figures given for weight of faces during this balance period are tabulated for convenience as above, but naturally the last yield was not obtained until the 8th of March. The total of 182 grams, however, is the exact amount of air-dry faces collected between the two charcoal layers, marking off accurately the balance period.

### NITROGEN BALANCE. - Henderson.

	Nitrogen Taken in.	O Nitrogen in Urine.	utput. Weight of Freces (dry).
	Taken in.	Marogen in Orme.	weight of Pasces (dry).
Feb. 29	10.261 grams.	8.36 grams.	• • •
Mar. 1	7.88 <del>4</del>	6.80	•••
2	8.487	8.28	85 grams.
8	8.555	7.87	•••
4	10.427	8.22	• • •
5	10.488	8.09	•••
6	10.265	8.20	28
			118 grams contain
			6.21 % N.
	65.862	55.32 +	7.017 grams nitrogen.
	65.862 grams n	itrogen. 62.	887 grams nitrogen.

Nitrogen balance for seven days = +3.525 grams. Nitrogen balance per day = +0.508 grams.

### NITROGEN BALANCE. - Morris.

		Nitrogen Taken in.	Nitrogen in Urine.	Output. ' Weigl	nt of Faces (dry).
Feb. 2	9	10.466 grams.	6.40 gram	•	, ••
Mar.	1	7.720	6.64	4	l5 grams.
	2	8.819	<b>5.40</b>	4	13
	8	8.750	6.55	4	<b>!</b> 1
	4	10.427	4.99	. 4	15
	5	10.483	5.38		38
	6	10.265	7.01	1	13
				2	— 25 grams contain
					7.18% N.
		66.480	42.37	+ 1	6.155 grams nitrogen.
		66.430 grams	nitrogen.	58.525 gran	ns nitrogen.

Nitrogen balance for seven days = +7.905 grams. Nitrogen balance per day = +1.129 grams.

# NITROGEN BALANCE. - Coffman.

		Nitrogen		output			
		Taken in.	Nitrogen in Urine.	Weight of Free	se (dry).		
Feb. 2	29	10.466 grams.	8.63 gram	s			
Mar.	1	7.825	7.48	•••			
	2	8.487	8.62	•••			
	8	8.750	7.18	•••			
	4	10.427	7.93	65 grams	, ,		
	5	10.483	7.67	88			
	6	10.265	7.95	20			
				168 grams	contain		
				e	3.14% N.		
		66.708	55.46	+ 10.815 gram	s nitrogen.		
		66.703 grams	nitrogen. 65	.775 grams nitroge	n.		

Nitrogen balance for seven days = +0.928 gram. Nitrogen balance per day = +0.182 gram.



ZOOMAN COHN
Photographs taken at the close of the experiment.

• .

## NITROGEN BALANCE. - Steltz.

		Nitrogen		Output.	
		Taken in.	Nitrogen in Urine.	Weight of F	zeces (dry).
Feb.	29	9.989 grams.	7.90 gram	s. 23 gra	ms.
Mar.	1	8.090	5.98	22	
	2	8.487	6.58	21	
	8	9.800	8.10	45	
	4	11.107	4.66	18	
	5	10.024	8.69	86	
	6	10.392	8.20	31	
				10	
				206 gras	ms contain
					6.66 % N.
		67.389	50.11	+ 13.719	grams nitrogen.
		67.389 grams	nitrogen. 6	3.829 grams nit	rogen.

Nitrogen balance for seven days = +3.560 grams. Nitrogen balance per day = +0.508 gram,

### NITROGEN BALANCE. - Loewenthal.

		Nitrogen		Output.			
		Taken in.	Nitrogen in	Urine.	Weight of Faces (dry).		
Feb.	29	10.466 grams.	7.38 gr	ams.	16.5 grams.		
Mar.	1	7.825	7.05		88.0		
	2	8.487	7.07		21.0		
•	8	8.750	7.83				
	4	10.427	7.85		15.0		
	5	10.654	4.29		<b>22.0</b> ·		
	6	10.886	8.07		56.0		
					168.5 grams contain		
					7.11% N.		
		67.495	49.04	_+_	11.980 grams nitrogen.		
		67.495 grams n	itrogen.	61.020	grams nitrogen.		

Nitrogen balance for seven days = +8.475 grams. Nitrogen balance per day = +0.925 gram.

### NITROGEN BALANCE. -- Cohn.

		Nitrogen	Outp	at.		
		Taken in.	Nitrogen in Urine.	Weight of Fæces (dry).		
Feb.	29	10.442 grams.	9.74 grams.	• • •		
Mar.	1	7.825	6.86	• • •		
	2	8.487	7.29	43 grams.		
	4	10.215	5.59	24		
	5	10.483	9.55	57		
	6	8.164	6.77	25		
				149 grams contain		
				6.63 % N. = 9.878 gr.		
			•	-1.401 =		
		55.616	45.80 +	8.477 grams nitrogen.		
		55.616 grams n	itrogen. 54.277	grams nitrogen.		

Nitrogen balance for six days = +1.339 grams. Nitrogen balance per day = +0.223 gram.

<sup>•</sup> This balance is complicated by the loss of the urine on March 8. Consequently one-seventh of the total nitrogen of the fæces of the period, namely 1.401 grams, was deducted from the total fæcal nitrogen.

### NITROGEN BALANCE. - Zooman.

		Nitrogen	WI I- V	Output.			
Feb.	29	Taken in. 10.466 gram	Ni rogen in U 5 7.02 grs	•	of Faces (dry). grams.		
Mar.	1	7.604	6.66	25			
	2	8.487	7.26	52			
	3	8.750	7.56	40	1		
	4	10.427	7.51	14			
	5	10.483	7.08	30			
	6	10.265	6.81	10 189	grams contain		
					6.54 % N.		
		66.482	49.90	+ 12.36	grams nitrogen.		
		66.482 gram	s nitrogen.	62.26 grams nitrogen.			

Nitrogen balance for seven days = +4.222 grams. Nitrogen balance per day = +0.608 gram.

## NITROGEN BALANCE. - Sliney.

		Nitrogen	(	Output.		
		Taken in.	Mitrogen in Urine	. Weight of Fraces (dry).		
Feb.	29	10.466 grams.	9.50 grams.	15 grams.		
Mar.	1	7.825	8.09	<b>35</b>		
	2	8.487	7.97	25		
	8	8.73 <b>4</b>	7.78	45		
	4	10.427	7.49	28		
	5	10.475	7.54	21		
	6	10.265	8.23	32		
				201 grams contain		
				6.70% N.		
		66.679	56.60 <b>+</b>	13.467 grams nitrogen.		
		66.679 grams ni	trogen. 70.067	grams nitrogen.		

Nitrogen balance for seven days = -3.388 grams. Nitrogen balance per day = -0.484 gram.

# 240 PHYSIOLOGICAL ECONOMY IN NUTRITION

# NITROGEN BALANCE. - Broyles.

	Nitrogen Taken in, Nitroge	Outp on in Urine.	ut. Weight in Fæ	es (dry).
Feb. 29	10.179 grams.	5.74 gra	ms	
Mar. 1	7.468	9.26		
2	8.487	6.97	13 g	rams.
3	8.750	6.18	4	
4	10.427	7.68	46	
<b>6</b>	10.483	5.56	29	
6	10.265	7.69	42	
			134 g	rams contain
				6.24% N.
	66.059	49.08	+ 8.861	grams nitrogen.
	66.059 grams ni	trogen. 5	7.441 grams	nitrogen.

Nitrogen balance for seven days = +8.618 grams. Nitrogen balance per day = +1.231 grams.



ZOOMAN COHN

Photographs taken at the close of the experiment.



### NITROGEN BALANCE. - Fritz.

	Nitrogen Taken in.	Mitrogen in U	Output. rine. Weight of	Faces (dry).	
Feb. 29	10.466 grams.	6.22 grams.			
Mar. 1	7.825	7.44			
2	8.487	4.00	71		
3	8.750	7.05	28		
5	10.483	8.71	49		
6	10.265	4.78	44		
			192 grams	contain	
			6.61%	N. =12.691 gr.	
				= 1.818 *	
	56.276	88.20	+ '	10.878 grm. N.	
	56.276 grams n	itrogen.	49.078 grams nitrogen.		
	Nitrogen balance i	or six days	- +7.198	grams.	
	Nitrogen balance	per day	- +1.199	grams.	

<sup>\*</sup> This balance is somewhat complicated by the fact that on March 4 the urine was lost, so that this day had to be thrown out. Correction on the fæces, however, was made by deducting one-seventh of the total fæcal nitrogen, on the assumption that the nitrogen-content was essentially the same for each day of the seven-day period.

### THIRD NITROGEN BALANCE.

The last of March, a third nitrogen balance was tried on a slightly lowered nitrogen intake and with a slight increase in the fuel value of the daily food. In this period of five days, March 28 to April 1, the nitrogen taken in per day averaged 8.62 grams, or nearly one gram per day less than in the preceding period. The fuel value of the food averaged 2840 calories per day, or about 300 calories more than in the preceding period.

The daily diet, with its content of nitrogen, etc., is detailed for each day under Oakman. Any deviation from this diet in the cases of the other men is indicated on the accompanying balance-sheets.

The results obtained in this balance period indicate that the lowest level had been practically reached, at least under the conditions of body-weight, food, and work prevailing. Cohn, Fritz, and Broyles showed a distinct positive balance. Steltz and Loewenthal were practically in equilibrium, the deviation being within the limits of error. The remaining six men showed a minus balance, although in no case was it very marked.

It is interesting to note in this connection that the average daily output of nitrogen through the urine for this five days' period (Oakman's case) amounted to 7.04 grams, being 0.2 gram less per day than in the preceding period. This figure for nitrogen in the urine means the metabolism daily of 44 grams of proteid.

Undoubtedly, the rate of proteid metabolism for these men could have been lowered considerably beyond the present level by increasing largely the intake of carbohydrates and fats, but it has been the intent throughout all of these experiments to learn the minimal proteid requirement under conditions precluding the use of any excess of non-nitrogenous foods; also, to study the effect of a general physiological economy in nutrition, with a view to ascertaining the real necessities of the body for both proteid and non-proteid foods

with maintenance of bodily strength and vigor. Hence, we may again emphasize the fact that the low proteid metabolism maintained by all these men throughout the period of the experiment, with establishment of nitrogenous equilibrium on a consumption of proteid or albuminous food averaging one-half the amount ordinarily specified as the daily requirement of the healthy man, has been accomplished with even less total food—fats and carbohydrates—than the ordinary standards call for, *i. e.*, considerably less than 3000 calories per day in fuel value.

#### OAKMAN.

# Monday, March 28, 1904.

Breakfast.—Fried rice 150 grams, syrup 75 grams, baked potato 250 grams, butter 20 grams, one cup coffee 350 grams.

Dinner. — Thick pea soup 200 grams, boiled onions 100 grams, boiled sweet potato 250 grams, bread 50 grams, mashed potato 200 grams, butter 20 grams, one cup coffee 350 grams.

Supper. — Sliced banana 150 grams, biscuit 125 grams, fried bacon 20 grams, French fried potato 200 grams, butter 25 grams, one cup tea 350 grams.

Food.			Grams.	Per	cent Nitrogen.	Total Nitrogen.
Fried rice			150	×	0.40 =	0.600 grams.
Syrup			75	×	0.051 =	0.038
Baked potato .			250	×	0.33 =	0.825
Butter 25				×	0.15 =	0.097
Coffee (breakfast	) .		350	×	0.10 =	0.350
Pea soup			200	×	0.50 =	1.000
Boiled onion .			100	×	0.30 =	0.300
Sweet potato .			250	×	0.12 =	0.300
Bread			50	×	1.57 =	0.785
Mashed potato			200	×	0.80 =	0.600
Coffee (dinner)			350	×	0.075 =	0.262
Banana			150	×	0.27 =	0.405
Biscuit			125	×	1.21 =	1.513
Bacon			20	×	8.85 =	0.770
Fried potato .			200	×	0.46 =	0.920
Tea			850	×	0.075 =	0.262
	Total	l nitrogen	in food			9.027 grams.
	Total	nitrogen	in urine			. 6.640

Fuel value of the food . . . . 2935 calories.

# Tuesday, March 29, 1904.

Breakfast. — Boiled hominy 175 grams, milk 75 grams, sugar 25 grams, baked potato 250 grams, butter 20 grams, one cup coffee 350 grams.

Dinner. — Hamburg steak with much bread, fat, and onions 125 grams, boiled potato 300 grams, butter 10 grams, one cup coffee 350 grams, bread 85 grams, boiled carrots 125 grams.

Supper. — Tapioca-peach pudding 300 grams, bread 35 grams, Saratoga chips 75 grams, butter 20 grams, jam 75 grams, one cup tea 350 grams.

Food.	Grams.	Per	cent Nitro	gen.	Total Nitrogen.
Boiled hominy	. 175	X	0.23	=	0.403 grams.
Milk	. 75	×	0.55	=	0.418
Sugar	. 25	×	0.00	=	0.000
Baked potato	. 250	×	0.36	=	0.900
Butter $20 + 10 + 20 =$	= 50	×	0.15	=	0.075
Coffee (breakfast)	. 350	×	0.057	=	0.200
Hamburg steak, etc	. 125	×	2.50	=	3.180
Boiled potato	. 800	×	0.26	=	0.780
Coffee (dinner)	. 350	×	0.051	=	0.179
Carrots	. 125	×	0.13	=	0.163
Tapioca-peach pudding	. 300	×	0.28	=	0.840
Bread $85 + 35 =$	= 70	×	1.51	=	1.057
Saratoga chips	. 75	×	0.79	=	0.598
Jam	. 75	×	0.039	=	0.029
Tea	. 350	×	0.06	=	0.210
Total nitroge	n in food				8.972 grams.
Total nitroger	ı in urine				8.840

Fuel value of the food . . . . 2840 calories.

# Wednesday, March 30, 1904.

- Breakfast.—Fried hominy 150 grams, syrup 75 grams, butter 10 grams, sliced banana 250 grams, one cup coffee 350 grams.
- Dinner. Codfish-balls (1 part fish, 5 parts potatoes, fried in pork fat) 125 grams, mashed potato 250 grams, stewed tomato 200 grams, bread 35 grams, apple sauce 200 grams, one cup coffee 350 grams.
- Supper. Chopped fresh cabbage with salt, pepper, and vinegar 75 grams, bread 50 grams, butter 20 grams, fried sweet potato 250 grams, cranberry sauce 200 grams, sponge cake 50 grams, one cup tea 350 grams.

Food.	Grams.	Per	oent Nitrogen.	Total Nitrogen.
Fried hominy	. 150	X	0.35 =	0.525 grams.
Syrup	. 75	×	0.051 =	0.038
Butter $10 + 20 =$	: 30	×	0.15 =	0.045
Banana	. 250	×	0.25 =	0.625
Coffee (breakfast)	. 350	×	0.066 =	0.281
Codfish-balls, etc	. 125	×	3.25 =	4.063
Mashed potato	. 250	×	0.26 =	0.650
Tomato	200	×	0.18 =	0.860
Bread $35 + 50 =$	<b>85</b>	×	1.50 =	1.280
Apple sauce	<b>20</b> 0	×	0.058 =	0.106
Coffee (dinner)	. 350	×	0.051 =	0.179
Cabbage	. 75	×	0.22 =	0.165
Fried sweet potato	250	×	0.15 =	0.375
Cranberry sauce	. 200	×	0.066 =	0.182
Sponge cake	. 50	×	0.87 =	0.435
Tea	. 850	×	0.042 =	0.147
Total nitrogen Total nitrogen				9.366 grams.

Fuel value of the food . . . . 2657 calories.

# Thursday, March 31, 1904.

Breakfast. — Fried Indian meal 100 grams, syrup 75 grams, baked potato 250 grams, one cup coffee 350 grams, butter 20 grams.

Dinner. — Tomato soup, thick, with potatoes and onions boiled together 300 grams, mashed potato 200 grams, scrambled egg 50 grams, bread 50 grams, butter 10 grams, one cup coffee 350 grams.

Supper. — Bread pudding 150 grams, sliced banana 200 grams, fried bacon 20 grams, boiled potato 200 grams, butter 10 grams, one cup tea 850 grams.

Food.							Grams.	Pe	r cent Nitro	gen.	Total Nitrogen.
Fried Indian mea	ı						100	×	0.44	=	0.440 grams.
Syrup							75	×	0.051	=	0.038
Baked potato .							250	×	0.82	=	0.800
Coffee (breakfas	t)						350	X	0.066	=	0.281
Butter 10	+ (	20	+	10	)	=	40	×	0.15	=	0.060
Tomato soup .							300	X	0.48	=	1.440
Mashed potato							200	×	0.26	=	0.520
Scrambled egg							50	X	2.15	=	1.075
Bread							50	X	1.48	=	0.740
Coffee (dinner)							350	X	0.057	=	0.200
Bread pudding							150	X	0.92	=	1.880
Banana							200	X	0.24	=	0.480
Bacon							20	X	1.95	=	0.890
Boiled potato .							200	×	0.25	=	0.500
Tea	•	•			•		<b>350</b>	×	0.086	=	0.126
	T	otal	ni	tro	oge	en :	in food				8.420 grams.
	T	otal	n	itr	og	en	in urine				. 7.100

Fuel value of the food . . . . 2466 calories.

# Friday, April 1, 1904.

- Breakfast. Fried hominy 150 grams, syrup 75 grams, baked potato 200 grams, butter 20 grams, one cup coffee 850 grams.
- Dinner.—Baked spaghetti 250 grams, mashed potato 250 grams, boiled turnip 150 grams, bread 85 grams, butter 10 grams, apple sauce 200 grams, one cup coffee 850 grams.
- Supper. Apple-tapioca pudding 300 grams, fried sweet potato 200 grams, butter 20 grams, jam 100 grams, fried bacon 25 grams, bread 35 grams, one cup tea 350 grams.

Food.	Grams.	Per	oent Nitrogen.	Total Nitrogen.
Fried hominy	150	×	0.24 =	0.360 grams.
Syrup		×	0.051 =	0.038
Baked potato	200	×	0.33 =	0.660
Butter $20 + 10 + 20 =$	50	×	0.15 =	0.075
Coffee (breakfast)	350	×	0.06 =	0.210
Spaghetti	250	×	0.64 =	1.600
Mashed potato	250	×	0.28	0.700
Boiled turnip	150	×	0.61 =	0.915
Bread $35 + 35 =$	70	×	1.56 =	1.092
Apple sauce	200	×	0.058 =	0.106
Coffee (dinner)	350	×	0.072 =	0.252
Apple-tapioca pudding	800	×	0.043 =	0.129
Fried sweet potato	200	×	0.15 =	0.300
Jam	100	×	0.089 =	0.039
Bacon	25	×	2.96 =	0.740
Tea	350	×	= 880.0	0.126
Total nitrogen	in food			. 7.842 grams.
Total nitrogen				. 6.830

Fuel value of the food . . . . 3248 calories.

### NITROGEN BALANCE. - Oakman.

	Nitrogen	Output.					
		rogen in Urine.	Weight of Fraces (dry).				
Mar. 28	9.027 grams	6.64 gram	•				
29	8.972	8.84	89 grams.				
30	9.856	6.80	88				
81	8.420	7.10	31				
Apr. 1	7.342	6.83	29				
			137 grams contain				
			6.84 % N.				
	43.117	35.21 -	+ 9.87 grams nitrogen.				
	43.117 grams ni	trogen. 44.	44.580 grams nitrogen.				

Nitrogen balance for five days = -1.468 grams. Nitrogen balance per day = -0.292 gram.

#### Average Intake.

## NITROGEN BALANCE. - Broyles.

	Nitrogen	Outp	eut.
		Nitrogen in Urine.	Weight of Fraces (dry).
Mar. 28	9.027 grams.	6.79 grams.	27.0 grams.
29	8.972	7.06	45.5
30	9.356	7.27	41.0
81	8.640	6.21	25.0
Apr. 1	7.342	5.86	20.0
			158.5 grams contain
			5.92 % N.
	43.837	33.19 +	9.383 grams nitrogen.
	43.837 grams	nitrogen. 42.5	73 grams nitrogen.

Nitrogen balance for five days = +0.764 gram. Nitrogen balance per day = +0.163 gram.

## NITROGEN BALANCE. - Fritz.

	Nitrogen Taken in. Nit		tput. Weight of Freces (dry).
Mar. 29	7.229 grams.	5.09 gram	<b>15.</b>
30	9.235	7.10	25 grams.
31	8.640	5.18	43
<b>A</b> pr. 1	7.439	5.74	34
			102 grams contain
			6.42% N.
	32.543	23.11	+ 6.548 grams nitrogen.
	32.543 grams r	itrogen. 2	9.658 grams nitrogen.

Nitrogen balance for four days = +2.885 grams. Nitrogen balance per day = +0.721 gram.

# 250 PHYSIOLOGICAL ECONOMY IN NUTRITION

### NITROGEN BALANCE. - Loewenthal.

	Nitrogen	Output.	
	Taken in. Nit	rogen in Urine.	Weight of Fraces (dry).
Mar. 28	9.027 grams.	6.27 grams	s. 13.5 grams.
29	8.760	7.54	52.0
30	9.341	7.07	41.0
31	8.112	5.24	21.0
Apr. 1	7.842	6.43	17.0
			144.5 grams contain
			7.00% N.
	42.582	32.55 +	10.115 grams nitrogen
	42.582 grams ni	trogen. 42.6	65 grams nitrogen.

Nitrogen balance for five days = -0.083 gram. Nitrogen balance per day = -0.016 gram.

### NITROGEN BALANCE. - Cohn.

	Nitrogen Taken in.		tput. Weight of Faces (dry).	
Mar. 28	8.825 grams	•	` ` ` ` ` `	
29	8.151	6.48	27 grams.	
80	9.211	6.36	25	
31	8.030	7.47	23	
Apr. 1	7.246	4.35	39	
			114 grams contain	
			6.84 % N.	
	41.463	31.27 +	7.797 grams nitrogen.	
	41.463 grams	nitrogen. 89.	067 grams nitrogen.	

Nitrogen balance for five days = +2.896 grams. Nitrogen balance per day = +0.479 gram.

### NITROGEN BALANCE. - Coffman.

	Nitrogen Taken in. Nita		tput. Weight of Fæces (dry).
Mar. 28	9.027 grams.	8.38 gram	ı
29	8.972	8.06	42 grams.
30	9.356	6.88	24
31	8.640	7.78	29
Apr. 1	7.842	7.22	32
			127 grams contain
			6.54% N.
	43.387	38.32 +	- 8.306 grams nitrogen.
	43.387 grams ni	trogen. 46.	626 grams nitrogen.

Nitrogen balance for five days = -3.289 grams. Nitrogen balance per day = -0.657 gram.

### NITROGEN BALANCE. - Sliney.

	Nitrogen	Output.		
	Taken in. Nits	rogen in Urine.	Weight of Freces (dry).	
Mar. 28	8.527 grams.	7.09 gram	s. 28.0 grams.	
29	8.630	7.37	31.5	
80	9.856	8.10	33.0	
31	8.420	7.47	26.0	
Apr. 1	7.342	6.78	50.0	
			168.5 grams contain	
			6.42% N.	
	42.275	36.81 +	10.82 grams nitrogen.	
	42.275 grams n	itrogen. 47.	63 grams nitrogen.	

Nitrogen balance for five days = -5.355 grams. Nitrogen balance per day = -1.071 grams.

## 252 PHYSIOLOGICAL ECONOMY IN NUTRITION

### NITROGEN BALANCE. - Steltz.

Nitrogen	Output.		
	trogen in Urine.	Weight of Freces (dry).	
8.026 grams.	3.60 gram	ns. 32.0 grams.	
8.252	6.79	29.0	
9.255	7.20	44.0	
8.720	7.11	22.0	
7.857	7.82	30.5	
		157.5 grams contain	
		5.97 % N.	
41.610	82.52 -	+ 9.403 grams nitroge	n
41.610 grams	nitrogen. 41.	923 grams nitrogen.	
	8.028 grams. 8.252 9.255 8.720 7.357	Taken in.         Nitrogen in Urine.           8.026 grams.         3.60 gram           8.252         6.79           9.255         7.20           8.720         7.11           7.357         7.82           41.610         82.52	Taken in.         Nitrogen in Urine.         Weight of Faces (dry).           8.026 grams.         3.60 grams.         32.0 grams.           8.252         6.79         29.0           9.255         7.20         44.0           8.720         7.11         22.0           7.357         7.82         30.5           167.5 grams contain         5.97 % N.           41.610         82.52         +         9.403 grams nitroget

Nitrogen balance for five days = -0.813 gram. Nitrogen balance per day = -0.062 gram.

#### NITROGEN BALANCE. - Zooman.

	Nitrogen Taken in. Nitr	Outpu ogen in Urine. W	t. Teight of Fæces (dry).
Mar. 28	9.027 grams.	7.47 grams.	88.0 grams.
29	8.379	7.80	45.0
30	9.856	5.40	30.0
81	8.420	8.04	28.0
Apr. 1	7.269	8.44	11.5
	<del></del>		147.5 grams contain
			6.39% N.
	42.451	37.15 +	9.425 grams nitrogen,
	42.451 grams ni	trogen. 46.57	5 grams nitrogen.

Nitrogen balance for five days = -4.124 grams. Nitrogen balance per day = -0.824 gram.

### NITROGEN BALANCE. - Henderson.

	Nitrogen	Output.	
	Taken in. Ni	trogen in Urine.	Weight of Faces (dry).
Mar. 28	9.027 grams.	8.40 grams	
29	8.972	9.04	• • •
30	9.856	5.95	83 grams.
31	8.640	5.42	36
Apr. 1	7.342	6.60	<u>58</u>
			177 grams contain
			6.48% N.
	43.837	85.41 +	11.469 grams nitrogen.
	43.387 grams n	itrogen. 46.87	79 grams nitrogen.

Nitrogen balance for five days = -8.542 grams. Nitrogen balance per day = -0.708 gram.

### NITROGEN BALANCE. - Morris.

	Nitrogen	Output.		
		rogen in Urine. V	Veight of Fæces (dry).	
Mar. 28	8.877 grams.	6.68 grams.	12 grams.	
29	8.774	5.69	48	
80	8.941	6.06	34	
31	8.420	6.96	84	
Apr. 1	7.286	7.10	83	
			161 grams contain	
			7.08% N.	
	42.298	32.49 +	11.399 grams nitrogen.	
	42.298 grams ni	itrogen. 43.889	grams nitrogen	

Nitrogen balance for five days = -1.591 grams. Nitrogen balance per day = -0.317 gram.

# SUMMARY REGARDING NITROGEN REQUIREMENT.

These results obtained with this body of United States soldiers, living on a prescribed diet and exposed to the stress and strain of military discipline with its attendant duties, together with the gymnastic work and training required each day, confirm in every detail the conclusions arrived at with the preceding group of professional workers. Once accustomed to a more sparing proteid diet, less rich in nitrogen, each one of these subjects had no difficulty in maintaining body-weight on the simpler and lighter food provided. great difficulty was experienced in establishing a condition of nitrogenous equilibrium with this lowered intake of proteid food, neither was it necessary to increase the amounts of nonnitrogenous foods (fats and carbohydrates) to accomplish this end. The bodies of these men were quite able to adjust themselves to a lowered proteid metabolism, and physiologically speaking, one might well conjecture whether we have not in this condition a nearer approach to the normal and ideal state of the body than when the latter is struggling daily with 118 grams of proteid food, reinforced by fats and carbohydrates correspondingly increased in amount. ever this may be, the members of the soldier detail were able to live for five consecutive months with a proteid metabolism corresponding to 7 to 8 grams of nitrogen per day, with maintenance of body-weight and without discomfort or loss of bodily vigor.

It was easy in most instances to prove the establishment of nitrogen equilibrium with a daily intake of 8.5 to 9.5 grams of nitrogen, and with a total fuel value of the daily food equal to 2500 to 2800 calories. In other words, a metabolism of less than 50 grams of proteid per day was quite sufficient for the needs of the body, and a fuel value of 2500 to 2600 calories was ample to meet the requirements of the men under the then existing conditions of bodily and mental activity. Are we not justified, therefore, in again asking the question, why should we hold and teach the doctrine that the healthy adult

needs to metabolize 105 grams of proteid food daily? As Voit has well said, the smallest amount of food that will serve to maintain physiological equilibrium and keep up health and strength is the ideal diet. The eleven subjects of this Hospital detachment, who remained throughout the experiment, were apparently able to maintain physiological equilibrium and preserve their health and strength under the conditions of diet as described, thereby demonstrating the possibilities of a physiological economy corresponding to a saving of full fifty per cent or more in proteid food; a saving of possibly great physiological import, to say nothing of the possible economic and sociological importance of the saving. Further, we may add that the minimal proteid requirement as evidenced by the results of these experiments is more than fifty per cent lower than the figures quoted by most physiologists as necessary for the maintenance of life and strength; and we are certainly justified in the additional statement that if the figures obtained in these experiments truly represent the minimal proteid requirement of the men under observation, then this minimal requirement is quite sufficient to meet the physiological needs of the body for an indefinite period.

### PHYSICAL CONDITION OF THE MEN.

Recalling the fact that this condition of lowered proteid metabolism was maintained for a period of five months, we may next consider the effect of this changed nutritive condition upon the health and strength of the men. The question of body-weight we have already considered. More pertinent is the question, to how great an extent was the strength and bodily vigor of the men modified by the diminished amount of proteid food? The answer to this question is found in the subjoined report from Dr. William G. Anderson, Director of the Yale University Gymnasium.

YALE UNIVERSITY GYMNASIUM, NEW HAVEN, CONN., April 6, 1904.

To Professor Russell H. Chittenden,
Director of the Sheffield Scientific
School, Yale University,
New Haven, Conn.

Dear Sir, — I hand you herewith a report of the physical training of the squad of soldiers sent by you to the Yale Gymnasium.

These men have taken one hour's exercise daily for six months, Sundays excepted, — October 1, 1903, to April 1, 1904.

Before beginning the bodily development of the men we measured each one and took what are known as the American Collegiate Strength Tests. These measurements and tests have been taken twice each month. For details as to measurements and methods of testing the strength of the body, see the "Notes" attached to this report.

The members of the squad were called to the floor each morning as soon after nine-thirty as they could don the required suit. For twenty minutes they were put through a series of setting up exercises and body-building movements; then followed exercises on the apparatus, such as bars, rings, ladders, etc. This was followed by a jump or game.

The exercises were progressive as to duration, force, extent, and number of movements. At the end of the six months the men were being put through gymnastics that were exacting and fatiguing. As the progression was carefully made, the men did not suffer from soreness to any marked degree. By way of comparison it may be said, the gymnastic training given the soldiers was much more severe than is given to the Freshmen of Yale in their required physical training. Perhaps the Varsity Crew are "put through" as rigid gymnastic training as any of the athletes in college, so we copied the exercises taken by oarsmen in order to "try out" the soldiers. This work was easily taken by all the "Dieters,"—no complaint, so far as I know, having been entered.

The improvement in accuracy and grace of movement has been noticeable, while there has been a gain in skill as well.

I do not consider these men, as a body, well put up. They did not rank favorably with applicants for policeman or fireman and

were noticeably timid in exercises that called for courage. In the vault over a fence there was a very marked fear. This was the poorest and least satisfactory test of all we made. At the end of the six months the hesitancy to vault had entirely disappeared, the event being looked upon by the men as a "cinch."

The gain in self-reliance and courage has been as evident as the increase in the figures of the strength test. I consider this a valuable acquisition as it stands for a good body condition. Among athletes, especially gymnasts, a weakened or tired body is made known by fear or uncertainty. Our students, as well as professional athletes, admit this.

I attribute the timidity of the soldiers in these seemingly simple tests to their general physical condition at the outset, and the increase in self-reliance to the better condition of their bodies at the end of the six months' training. The story told by the measurements is both interesting and significant, but of less importance than the ability to improve under training, which ability cannot be measured with a tape or weighed with scales.

In the majority of cases there was a loss in body-weight, but this we look for, as the larger number of soldiers were fairly well supplied with adipose tissue. Both Sliney and Oakman are exceptions, these men being "fine" at the outset. I mean thin and muscular with little fat. The latter, Oakman, is an old man judged from the physical activity standpoint, and lost noticeably during the tests and exercise. I refer to weight and girth measurements only. He made a very large gain in his strength tests and was among the leading men in all that called for courage and self-control. Sliney, nervous, irritable, and aggressive, balanced his losses with his gains in the tape and scale events, but made great improvement in the use of the dynamometers. Fritz and Cohn were quite fat and showed the expected loss in weight, but not in other respects.

For comparison I have quoted from Dr. J. W. Seaver's anthropometric charts and have selected the "Average Student Measurements" (2390 men) as well as the mean measurements of 500 athletes and gymnasts. A glance at either set of figures will give an idea of how the soldiers compare with men living on a regular diet and surrounded with the very best environment.

The strength tests stand for mental states more than the tape line and calipers, and are suggestive of improvement in body con-

ditions for the reasons mentioned above. These records are far above those made by Academic Freshmen in Yale, but it must be added that the soldiers have taken more exercise than the collegians.

The hearts and lungs of the men are in excellent condition, while the soldiers as a body are in better shape physically, are stronger and healthier than in October. The skin of the men is clear and ruddy.

The figures for comparison in the Strength Test Table are from the records sent me by Dr. Geo. Meylan, the Director of the Columbia University Gymnasium. I understand that others than college men are represented in his data. In the figures given in my own tables I have omitted the small fractions in the main body of the tables, but have recorded them in the totals.

The greater portion of the training of the soldiers has been under the personal supervision of Wm. H. Callahan, M. D., the Medical Assistant at the Gymnasium, to whom I am indebted for help and suggestions. Mr. Wm. Chase, Mr. Anton Muller, Mr. John Stapleton, and Mr. H. R. Gladwin, Assistant Instructors in the Gymnasium, have led the drills and have looked after the actual muscular training of the men. I gladly acknowledge the co-operation of these gentlemen in the physical development of the squad.

A brief summary of my conclusions: —

The men were not above the average standard, physically, when they began their work, this standard being set by applicants for firemen and policemen, not by college students. At the end of their training they were much above the same standard, while their strength tests were far greater than the averages made by college men. They showed less improvement in increase in size than University men do under like gymnastic treatment, but the gain in self-confidence and in body-fibre was very evident. The gain in accuracy and skill was marked. The men showed interest in their work throughout the six months. At the end of the period of training the soldiers were in excellent condition in spite of the fact that in some cases there was a slight loss of bodyweight. This loss is not to be attributed necessarily to the diet, because most men who exercise lose slightly if there is an excess of adipose tissue.

Respectfully yours, (Signed) W. G. Anderson.



LOEWENTHAL MORRIS

Photographs taken at the close of the experiment.



Dr. Anderson furnishes the following "Notes" descriptive of the methods by which the strength tests were made:

"Strength of Back. The subject, standing upon the iron foot-rest with the dynamometer so arranged that when grasping the handles with both hands his body will be inclined forward at an angle of 60 degrees, should take a full breath and, without bending the knees, give one hard lift, mostly with the back.

"Strength of Legs. The subject while standing on the footrest with body and head erect, and chest thrown forward, should sink down, by bending the knees, until the handle grasped rests against the thighs, then taking a full breath, he should lift hard principally with the legs, using the hands to hold the handle in place.

"Strength of Chest. The subject with his elbows extended at the sides until the forearms are on the same horizontal plane and holding the dynamometer so that the dial will face outward and the indicator point upward, should take a full breath and push vigorously against the handles, allowing the back of the instrument to press on the chest.

"Strength of Upper Arms, Triceps. The subject, while holding the position of rest upon the parallel bars, supporting his weight with arms straight, should let the body down until the chin is level with the bars, and then push it up again until the arms are fully extended. Note the number of times that he can lift himself in this manner.

"Strength of Upper Arms, Biceps. The subject should grasp a horizontal bar or pair of rings and hang with the feet clear from the floor while the arms are extended. Note the number of times that he can haul his body up until his chin touches the bar or ring.

"Strength of Forearms. The subject, while holding the dynamometer so that the dial is turned inward, should squeeze the spring as hard as possible, first with the right hand then with the left. The strength of the muscles between the shoulders may be tested with the same instrument. The subject, while holding the dynamometer on a level with the chest, should

grasp it with handles and pull both arms from the centre outward.

"The total strength is ascertained by multiplying the weight by the number of times it has been raised (push up and pull up), to this product we add the strength of hands, legs, back, and chest. The result is the total strength of the man. In some cases the product obtained by multiplying the weight by push up and pull up is divided by ten to reduce the size of the figures. We have not done so here.

"The run, vault, and ladder tests are not figured in. The lung capacity is also omitted from the final figures."

The following tables furnished by Dr. Anderson and Dr. Callahan give (1) the measurements of the eleven men who completed the experiment, taken on October 12, 1903, and For comparison are also given measurements April 2, 1904. of Yale College students, athletes, etc. (2) Strength or dynamometer tests, i. e., the first test taken in October and the final test taken in April, 1904, together with Columbia University strength tests for comparison. (3) A series of eleven tables giving for each man the individual strength tests, two or three each month, taken during the stay of the detachment in New Haven. Study of these individual results is quite interesting, since it shows very strikingly the gradual gain in strength of the men, and at the same time illustrates how temporary conditions, bodily or mental, may influence a record of this character, more noticeable in some individuals than in others. Mental stimulus, as is well known, counts for much in the manifestation of muscular power, but the neuro-muscular mechanism depends for its highest efficiency upon the nutritive condition of the tissues as much as does the muscle tissue alone. In the obtaining of a strength test, it is usually found that the best results are recorded when there is competition among the men; i. e., under the influence of an outside stimulus.

## MEASUREMENTS.

(OCTOBER 12 - APRIL 2.)

		Weight.	Girth Nock.	Waist.	Chest Normal.	Chest Inflated.	Chest Deflated.	Right Bloeps.	Right Thigh.	Left Bloops.	Left Calf	Left Thigh.	Right Calf.
Henderson	{	157 158	362 869	755 731	953 942	1085 1025	890 886	816 311	510 508	291 287	362 357	500 495	351 347
Oakman	{	145 137	344 386	788 727	888 869	947 938	798 795	305 297	508 502	289 282	335 881	512 507	330 331
Morris	{	129 131	340 851	718 719	850 862	982 938	831 834	280 285	519 525	288 290	327 838	502 509	825 832
Zooman	{	120 122	850 860	718 712	868 859	920 914	804 803	290 307	479 486	286 298	812 815	478 484	313 814
Coffman	{	129 127	352 349	788 716	860 857	923 981	815 818	309 314	518 527	811 316	354 359	520 529	850 854
Steltz	{	116 115	330 380	713 717	815 821	850 857	779 771	291 287	487 492	283 281	319 324	483 478	318 822
Loewenthal	{	133 180	338 341	680 684	835 840	881 890	822 819	315 306	510 504	811 302	388 842	512 505	333 887
Sliney	{	135 183	359 361	718 702	824 834	930 926	823 818	314 806	476 475	302 298	331 333	473 475	830 882
Fritz	{	167 161	890 861	860 795	880 892	930 997	860 855	860 858	560 555	343 340	877 372	562 559	375 373
Cohn	{	142 138	363 854	810 771	871 878	912 914	882 818	826 824	586 529	310 308	374 371	528 520	875 871
Broyles	{	130 133	856 870	710 720	851 864	941 966	805 798	281 283	500 508	282 284	841 844	498 501	334 338
Yale College †		139 147	850 857	780 747	861 886	910 940		295 808	515 534		850 358	509 527	850 861

<sup>•</sup> Given here in pounds.

<sup>†</sup> Yale College students, 50% class of the mass of students (2,890 men).

<sup>†</sup> Yale College students, 50% or mean of 500 athletes and gymnasts, picked men.

## 262 PHYSIOLOGICAL ECONOMY IN NUTRITION

## STRENGTH OR DYNAMOMETER TESTS.

FIRST TRIALS OCTOBER, 1908; FINAL TEST APRIL, 1904.

	Weight.	Lang Capacity.	Right Hand.	Left Hand.	Ohpat.	Back.	Legs.	Pull up, Blospe.	Push up, Triceps.	One-fourth mile Run.	Vault.	Ladder.	Total.	
Henderson {	157 153	<b>46</b> 5 555	103 135	105 105	130 185	340 600	560 865	8 9	3 9	1.25 1.08	D† E	E†	2970 4598	Oct. Apr.
Oakman {	145 137	365 410	110 130	115 100	95 120	305 560	500 720	12 15	4 10	1.24 1.21	D E	D E	8445 5055	
Morris {	129 131	320 360	90 100	75 85	75 105	270 450	350 710	9 14	4 12	1.30 1.09	D E	D E	2543 4869	
Zooman {	120 122	400 420	110 115	90 95	130 100	350 440	850 910	8 13	9 18	1.40 1.13	F E	E E	3070 5 <b>4</b> 57	
Coffman {	129 127	290 400	105 105	103 85	100 110	820 440	580 880	7 20	6 17	1.20 1.13	F E	F E	2835 6269	
Steltz {	116 115	800 880	80 100	85 90	105 135	800 410	400 490	10 19	6 10	1.30 1.06	E	E E	2838 4581	
Loewenthal {	133 130	865 425	100 130	95 115	85 115	260 570	460 700	6 16	5 12	1.20 1.08	E	E	2463 5277	
Sliney	135 133	380 420	125 145	130 135	100 115	400 508	600 800	8 15	6 12	1.15 1.08	E	E E	8245 5307	
Fritz {	167 161	480 495	121 140	85 110	120 115	810 720	615 1080	4 11	3 8	2.40 1.17	D E	D E	2504 5178	
Cohn {	142 188	820 400	50 90	75 90	80 97	245 870	340 580	3 9	6 11	2.30 1.14	D E	D E	2210 4002	
Broyles {	130 134	485 515	105 105	95 110	90 135	340 560	560 875	6 15	4 13	1.45 1.15	D E	F E	2560 5530	
Columbia { University * }	138	410	103	100		882	417	9	8					

<sup>\*</sup> Columbia University strength test records for comparison. The 50% or mean test. \*From Dr. Meylan.

 $<sup>\</sup>dagger$  D = difficult; E = easy; F = failure.



Soldiers exercising in the gymnasium.



## INDIVIDUAL STRENGTH TESTS.

## MORRIS.

		Weight.	Pull up.	Push up.	Right Hand.	Left Hand.	Chest.	Back.	Legs.	Vault and Ladder.	Long Capacity.	One-fourth Mile Ran.	Product.	Total.
Oct.	1, '03	129	9	4	90	75	75	270	850	D	820	1.80	1683	2543
Oct.	12, '03	130	7	2	95	75	75	250	430	D	810	1.40	1170	2095
Oct.	26, '03	132	8	5	85	70	80	250	400	D	320		1120	2667
Nov.	9, '03	183	12	7	108	84	85	340	457	F	880	1. <b>4</b> 5	2536	3620
No▼.	23, '08	132	12	8	95	70	90	315	450	F	825	1.18	2046	S066
Dec.	1, '03	184	10	6	95	85	75	350	455	F	350	1.14	2152	3212
Dec.	15, '03	130	7	4	100	100	85	400	490	F	840	1.12	1430	2605
Dec.	29, '03	129	10	6	100	85	90	260	<b>4</b> 75	E	330	1.11	2075	8085
Jan.	12, '04	131	10	6	95	69	90	260	445	E	345	1.15	2108	8067
Jan.	24, '04	132	10	5	80	80	85	340	510	E	850	1.18	1980	3075
Feb.	9, '04	180	7	6	80	70	85	280		E	345	1.14	1699	2214
Feb.	23, '04	134	5	6	100	100	90	810	450	E	360	1.09	1479	2529
Mar.	8, '04	131	10	10	90	85	95	375	550	E	820	1.14	2620	3815
Mar.	22, '04	132	10	9	98	82	100	370	500	E	360	1.15	2508	3658
April	2, '04	181	14	12	100	85	105	450	710	E	360	1.09	3419	4869

# 264 PHYSIOLOGICAL ECONOMY IN NUTRITION

## COFFMAN.

											Ė			
		Weight.	Pall ap.	Push up.	Right Hand.	Left Hand.	Chest.	Back.	Legs.	Vault and Ladder.	Lang Capacity.	One-fourth Mile Run.	Product.	Total.
Oct.	1, '03	129	7	6	105	108	100	320	580	D	290	1.20	1677	2835
Oct.	12, '08	129	9	8	110	85	120	350	500	F	310	1.09	2193	3858
Oct.	26, '08	129	9	12	100	100	110	350	610	F	345	1.11	2719	4119
Nov.	9, '03	181	18	12	105	100	125	370	725	E	350	1.12	8287	4727
Nov.	28, '03	180	12	11	95	85	120	320	620	E	860	1.12	2990	4280
Dec.	1, '03	133	13	18	107	83	111	880	580	E	380	1.12	3471	4632
Dec.	15, ′03	129	15	12	100	98	120	390	649	E	375	1.12	8488	<b>484</b> 0
Dec.	29, '03	126	12	12	100	85	100	250	580	E	865	1.07	3042	4157
Jan.	12, '04	127	12	12	100	86	107	350	575	E	360	1.12	3060	4278
Jan.	26, '04	127	15	11	105	97	100	465	680	E	400	1.13	3812	4759
Feb.	9, '04	126	15	14	95	85	100	410	725	E	390	1.11	3675	5090
Feb.	28, '04	126	17	14	100	95	90	<b>4</b> 60	555	E	380	1.09	3921	5221
Mar.	8, '04	126	18	16	90	95	100	425	675	E	390	1.14	4284	5669
Mar.	22, '04	129	19	16	115	93	100	480	600	E	400	1.10	4515	5918
Apr.	2, '04	127	20	17	105	85	110	440	830	E	400	1.13	4699	<b>626</b> 9

OAKMAN.

	Weight.	Pull up.	Push up.	Right Hand.	Left Hand.	Chest.	Back.	Legs.	Vault and Ladder.	Lung Capacity.	One-fourth Mile Run.	Product.	Total.
Oct. 1, '08	145	12	4	110	115	95	305	500	D	365	1.24	2320	3445
Oct. 12, '03	146	9	5	112	111	120	860	507	D	375	1.30	2044	3254
Oct. 26, '08	148	10	6	120	115	120	480	580	F	400	1.40	2368	8788
Nov. 9, '08	150	18	9	123	121	125	890	620	F	406	1.40	3150	4529
Nov. 28, '08	147	12	5	125		125	410	620	F	360	1.25	2499	8779
Dec. 1, '08	147	12	9	185	105	112	840	590	F	405	1.24	3092	4874
Dec. 15, '03	141	10	6	125	120	115	390	570	E	400	1.25	2264	8584
Dec. 29, '03	139	18	9	105	100	110	850	540	E	880	1.29	3058	4263
Jan. 12, '04	143	9	6	106	104	112	345	5 <b>4</b> 5	E	870	1.27	2152	3865
Jan. 26, '04	142	12	6	120	102	105	350	610	E	410	1.29	2556	8843
Feb. 9, '04	188	18	8	110	105	100	425	700	E	435	1.26	2913	4853
Feb. 23, '04	143	12	7	120	110	100	880	800	E	430	1.28	2716	4226
Mar. 8, '04	137	12	8	105	110	120	400	700	E	440	1.26	2755	4190
Mar. 22, '04	148	12	5	115	90	105	<b>400</b>	575	E	440	1.25	2489	3724
Apr. 2, '04	137	15	10	130	100	120	560	720	E	410	1.21	8425	5055

ZOOMAN.

	Weight.	Pull up.	Push up.	Right Hand.	Left Hand.	Chest.	Back.	Legs.	Vault and Ladder.	Lung Capacity.	One-fourth Mile Run.	Product.	Total.
Oct. 1, '08	120	8	9	110	90	180	850	350	D	400	1.40	2040	8070
Oct. 12. '08	121	9	12	100	95	180	860	870	F	410	1.28	2551	3616
Oct. 26, '03	124	11	17	115	75	130	400	400	F	440	1.11	3472	4592
Nov. 9, '03	126	9	18	115	107	125	365	540	E	420	1.14	3415	4681
Nov. 23, '03	120	13	20	110	100	117	850	470	E	400	1.12	3968	5115
Dec. 1, '08	123	12	18	120	105	103	445	520	E	380	1.10	3690	4988
Dec. 15, '03	120	12	17	100	90	110	360	490	E	410	1.14	3540	<b>4</b> 690
Dec. 29, '08	119	12	17	105	90	115	850	495	E	420	1.14	3472	4627
Jan. 12, '04	122	11	15	112	102	117	820	520	E	420	1.21	3172	4303
Jan. 26, '04	121	11	14	105	95	115	422	535	E	405	1.13	3043	<b>4</b> 315
Feb. 9, '04	122	10	17	95	85	115	<b>4</b> 60	<b>54</b> 5	E	420	1.10	3313	4613
Feb. 23, '04	121	10	17	100	105	112	850	570	E	408	1.12	3260	4497
Mar. 8, '04	119	12	18	105	95	100	425	700	E	410	1.12	8570	4995
Mar. 22, '04	125	12	19	105	94	105	875	650	E	420	1.18	3875	5204
Apr. 2, '04	122	18	18	115	95	100	<b>44</b> 0	910	E	<b>42</b> 0	1.13	8797	5 <b>4</b> 57

STELTZ.

	Weight.	Pull up.	Push up.	Right Hand.	Left Hand.	Chest.	Back.	Legs.	Vault and Ladder.	Lung Capacity.	One-fourth Mile Run.	Product.	Total.
Oct. 1, '03	116	10	6	80	85	105	300	400	D	800	1.30	1868	2838
Oct. 12, '03	118	11	5	95	78	120	310	520	D	320	1.22	1888	2911
Oct. 26, '03	118	15	10	80	80	120	820	450	F	360	1.12	2962	4012
Nov. 9, '08	120	15	10	91	96	127	375	505	F	345	1.20	3000	4194
Nov. 23, '03	118	17	7	95	110	115	310	430	<b>E</b>	875	1.20	2844	3904
Dec. 1, '08	119	10	8	93	85	78	830	865	E	400	1.24	2148	8099
Dec. 15, '03	116	12	4	85	62	115	880	335	E	400	1.15	1984	2961
Dec. 29, '08	116	15	10	75	65	95	250	800	E	380	1.09	2912	8697
Jan. 12, '04	119	11	11	85	90	115	270	415	E	880	1.10	<b>26</b> 18	3593
Jan. 26, '04	118	13	7	72	82	185	365	440	E	880	1.07	2370	8464
Feb. 9, '04	118	17	11	100	80	130	360	500	E	405	1.08	3304	4474
Feb. 23, '04	117	17	8	80	95	120	870	360	E	420	1.08	2937	3962
Mar. 8, '04	116	17	9	90	100	95	875	500	E	400	1.06	3016	4176
Mar. 22, '04	117	19	7	90	90	100	800	820	E	400	1.06	3055	3955
Apr. 2, '04	116	19	10	100	90	185	410	490	E	880	1.06	8856	4581

## HENDERSON.

	Weight.	Pull up.	Push up.	Right Hand.	Left Hand.	Chest.	Back.	Lega.	Vault and Ladder.	Lung Capacity.	One-fourth Mile Run.	Product.	Total.
Oct. 1, '08	157	8	8	108	105	130	340	560	F	465	1.25	1727	2965
Oct. 12, '03	159	в	5	115	115	135	300	500	F	485	1.19	1749	2914
Nov. 9, '03	162	9	9	105	105	140	420	700	E	525	1.07	2925	4428
Nov. 28, '03	157	11	9	185	115	145	500	750	E	515	1.07	31 <b>4</b> 0	8145
Dec. 15, '03												• • •	
Dec. 29, '03	153	5	4	105		125	250	365	E	520	1.16	1379	2109
Jan. 12, '04	153	5	5	85	60	135	320	595	E	530	1.18	1535	2780
Jan. 26, '04	155	5	3	100	65	105	350	460	E	5 <b>4</b> 0	1.14	1244	2382
Feb. 9, '04	151	5	5	85	80	115	350	400	E	465	1.12	1512	2542
Feb. 23, '04	158	9	7	110	115	125	450	610	E	535	1.13	2460	8870
Mar. 8, '04	151	8	7	130	110	120	550	875	E	540	1.18	2265	4050
Mar. 22, '04	155	9	8	122	100	125	870	570	E	550	1.12	2635	3922
Apr. 2, '04	158	9	9	135	105	135	600	865	E	555	1.08	2758	4598

## LOEWENTHAL.

	Weight.	Pull ap.	Push up.	Right Hand.	Left Hand.	Chest.	Back.	Legs.	Vault and Ladder.	Lung Capacity.	One-fourth Mile Bun.	Product.	Total.
Oct. 1, '08	183	в	5	100	95	85	260	460	D	<b>36</b> 5	1.20	1463	2468
Oct. 12, '03	138	12	6	120	115	90	340	470	F	870	1.11	2403	3588
Oct. 26, '08	184	7	٠.	125	110	105	370	490	F	860	1.25	988	2138
Nov. 9, '08	185	6	8	126	123	120	365	555	F	365	1.20	1897	8179
Nov. 23, '03	184	7	8	110	100	120	370	420	E	865	1.14	2021	8187
Dec. 1, '08	185	18	11	125	105	85	350	890	E	380		8240	4195
Dec. 15, '03	130	8	4	115	105	85	845	480	E	360	1.15	1560	2690
Dec. 29, '03	130	10	6	90	80	105	230	300	E	380		2080	2885
Jan. 12, '04	180	12	9	106	104	117	360	480	E	880	1.12	2740	3908
Jan. 26, '04	130			60	108	80	410	465	E	885	1.15		
Feb. 9, '04	180	13	8	115	120	115	455	550	E	420	1.09	2730	4085
Feb. 28, '04	133	14	6	115	105	100	430	500	E	385	1.08	2670	8920
Mar. 8, '04	130	14	9	125	120	115	450	700	E	420	1.14	3007	4517
Mar. 22, '04	132	6	11	103	113	120	870	500	E	420	1.10	2244	3450
Apr. 2, '04	180	16	12	180	115	115	570	700	E	425	1.08	8647	5277

## 270 PHYSIOLOGICAL ECONOMY IN NUTRITION

SLINEY.

							_	_	_	_			
	Weight.	Pull up.	Push up.	Right Hand.	Left Hand.	Chest.	Back.	Legs.	Vault and Ladder.	Lung Capacity.	One fourth Mile Run.	Product.	Total.
Oct. 12,'08	135	8	6	125	130	100	400	600	D	880	1.15	1890	3245
Oct. 26, '08	136	8	9	140	110	100	350	870	F	870	1.13	2260	3450
Nov. 9, '03	139	12	10	150	135	105	460	560	F	420	1.14	3127	4537
Nov. 23, '08	136	7	11	130	110	95	330	570	E	365	1.11	2448	3688
Dec. 1, '03	136	11	11	135	125	110	445	620	E	400	1.18	8071	4506
Dec. 15, '03	131	6	11	130	155	95	400	590	E	400	1.09	2281	3591
Dec. 29, '03	131	14	12	130	120	130	370	555	E	<b>42</b> 0	1.09	3419	4724
Jan. 12, '04	138	11	12	140	185	115	355	690	E	410	1.15	3174	4609
Jan. 26, '04	137	10	10	140	122	100	400	<b>74</b> 5	E	140	1.12	2745	4252
Feb. 9, '04	138	11	11	138	132	105	<b>45</b> 0	575	E	405	1.08	3105	<b>44</b> 05
Feb. 28, '04	139	12	10	140	145	99	430	650	E	405	1.08	3063	4527
Mar. 8, '04	135	14	13	150	180	110	525	825	E	440	1.08	8651	5391
Mar. 22, '04	139	14	. 9	145	138	115	840	565	E	440	1.07	8197	<b>4500</b>
Apr. 2,'04	183	15	12	145	185	115	508	<b>d00</b>	E	420	1.08	8604	5307

## FRITZ.

	Weight.	Pull up.	Push up.	Right Hand.	Left Hand.	Chest.	Back.	Lega.	Vault and Ladder.	Lung Capacity.	One-fourth Mile Run.	Product.	Total.
Oct. 31, '03	167	4	8	121	85	120	310	615	D	480	2.40	1252	2504
Nov. 28, '08	168	6	4	140	90	120	870	1050	F	480	1.80	1685	8455
Dec. 1, '03	172	10	6	130	95	124	345	565	F	480	1.20	2752	4011
Dec. 15, '03	165	7	4	135	85	120	850	720	F	485	1.19	1820	3230
Dec. 29, '08	164	8	9	110	85	90	250	415	E	500	1.20	2788	3588
Jan. 12, '04	165	6	11	118	72	115	425	560	E	525	1.18	2809	4099
Feb. 9, '04	165	11	5	142	117	125	600	880	E	510	1.25	2640	4504
Feb. 23, '04	165	12	5	180	95	120	580	960	E	520	1.18	2818	4678
Mar. 8, '04	162	11	7	125	80	110	550	875	E	500	1.21	2916	4656
Mar. 22, '04	165	10	2*	125	85	110	600	830	E	500	1.16	1986	8786
Apr. 2, '04	161	11	8	140	110	115	720	1080	E	495	1.17	8063	5178

<sup>\*</sup> Sore arm.

## CEX

	W-14/16.	4:11:4	Push up	Right Hand	I all Hand.	- Chres	Back	:	Variet and	A grand no	Our frugeh	Product.	Total.
39, 21 .we	345	\$	f	¥	7.5	Ħ	545	246	=	<b>13</b>	23	<b>1699</b>	2286
New . 25. YE	346	-	ž	316	¥	2	গ্ৰ	434	=		æ	1982	305
Dec. 1, W	147	•	ş	<b>S</b>	٦	16	201	414	7	12	:5	<b>753</b>	3313
Dec. 15, 48	30			舞	8	Ħ	=1	=	7	<b>**</b>	:5		
Dec. 28, 48	HE	ŧ	:	N	N	5	230	<b>D</b> evi	Ξ	<b>3</b>	LE	1866	2523
Jan 12 W	341	:	ş	Ţ	7	N	2W		Ŧ	254	.3	1430	200
Jan 36 We	347	f	ž	:2	7.5	386	XV.		Ŧ		:.#	He	365
Pol. 2. 74	14	f	H	*	٠,	N	25	67	Ŧ	<b>3</b>	1.35	235	3300
Fred 35 I'M	345	•	::	£3	¥	34	25	43/	Ŧ	सम	:J#	255	3586
Mar 22 Ye	:4	¥	૫	æ	я́і	3 <b>M</b> E	<b>3</b>	25	I	<b>7M</b>	: <b>]</b> £		3300
Apr. 2.44	:25	\$	::	N	۶	5	ដា	24	3	4	1.34	5.3	-



SLINEY

Photograph taken at the close of the experiment.



BROYLES.

	Weight.	Pull ap.	Push up.	Right Hand.	Left Hand.	Chest.	Back.	Logs.	Vault and Ladder.	Lung Capacity.	One-fourth Mile Run.	Product.	Total.
Nov. 18, '03	180	6	4	105	95	90	840	560	D	485	1.45	1370	<b>256</b> 0
Nov. 23, '03	127	9	10	95	100	95	370	630	F	465	1.17	2422	8007
Dec. 1, '03	130	13	11	130	120	105	420	680	F	480	1.14	3126	4581
Dec. 15, '08	129	9	13	110	110	125	870	470	F	510	1.12	2848	4023
Dec. 29, '03	123	15	11	105	90	125	300	600	E	480	1.14	8204	4424
Jan. 12, '08	126	15	12	101	95	180	870	580	E	485	1.14	3415	4691
Jan. 26,'04	180	10	8	103	95	120	470	890	E	480	1.15	2345	4022
Feb. 9, '04	130	10	10	105	105	115	510	625	E	500	1.15	2615	4075
Feb. 28, '04	184	12	6	115	100	115	550	850	E	505	1.15	2425	4155
Mar. 8, '04	182	15	8	110	110	120	525	900	E	510	1.17	8047	4812
Mar. 22, '04	188	12	12	110	110	125	470	850	E	510	1.14	8820	4989
Apr. 2, '04	183	15	18	105	110	135	560	875	E	515	1.15	3745	5530

The main things in Dr. Anderson's report especially to be emphasized are (1) the gain in self-reliance and courage of the men under training, indicative as they are of the better physical condition of their bodies, and (2) the marked in-

crease in their strength as indicated by the steady improvement in the strength or dynamometer tests. Obviously, the daily training to which the men were subjected in the Gymnasium is not to be overlooked as one factor in bringing about the gain in accuracy and skill, and indeed this factor must count for something in explaining the general gain in bodily strength, but increased skill alone will not account for the great gain in muscular power.

The results of these systematic tests make it very evident that the men were not being weakened by the lowered intake of proteid food. On the contrary, their ability to do muscular work was greatly increased; a fact which cannot well be connected with anything other than the physiological economy which was being practised. There must be enough food to make good the daily waste of tissue, enough food to furnish the energy of muscular contraction, but any surplus over and above what is necessary to supply these needs is not only a waste, but may prove an incubus, retarding the smooth working of the machinery and detracting from the power of the muscular mechanism to do its best work.

The figures showing the total strength of the men in October, 1903, on their ordinary diet, and on the second day of April, 1904, when the experiment was nearing completion are certainly very impressive.

#### TOTAL STRENGTH

						October	April
Henderso	n					2970	4598
Oakman .						3445	5055
Morris						2543	4869
Zooman						3070	5457
Coffman .						2835	6269
Steltz						2838	4581
Loewenth	al					2463	5277
Sliney						3245	5307
Fritz .						2504	5178
Cohn .						2210	4002
D1	_	_	_		_	2560	5530

Here we see gains in strength of 100 per cent in some cases, while Coffman shows an improvement so marked as to be almost marvellous. While there can be no question that a certain amount of this gain is to be attributed to the practice incidental to these months of work in the Gymnasium, it is equally clear that a large part of the gain is due to the improved physical condition of the men's bodies, for which the change in diet must be considered as responsible. In any event, the change from the ordinary diet to a diet comparatively poor in proteid has not resulted in any physical deterioration. On the contrary, there is every indication of a marked improvement in physical condition. In this connection the following note from Dr. DeWitt, commanding the detachment, is of interest:

882 Temple Street, New Haven, Conn., March 30, 1904.

Professor Russell H. Chittenden, Director Sheffield Scientific School, New Haven, Conn.

Sir, — In compliance with your verbal request I have the honor to inform you that at this date the men of this detachment are all in good physical condition.

Very respectfully,
(Signed) WALLACE DEWITT,
1st Lieut. and Asst. Surgeon U. S. Army,

Commdg. Det. H. C.

Finally, attention may be called to the photographs of the men, taken just prior to the close of the experiment, from which may be gained some idea of their physical condition so far as it can be judged by external appearance. Certainly, there is no indication in these photographs of any lack of bodily vigor. On the contrary, there is good muscular development, without any undue amount of fat, and indeed every indication of a good bodily condition, coupled with that appearance of quickness and alertness that belongs to the well-developed man, in a state of physiological balance. The

photographs scattered through this section of the book, showing the soldiers at work in the gymnasium, likewise give some idea of the lighter forms of exercise they followed each day in the training of bodily movements.

What now is to be said regarding the nervous condition of the men; i. e., their ability to respond to stimulation or, in other words, their mental quickness or reaction time? To study this question, the soldiers were sent with regularity to the Yale Psychological Laboratory, where their reaction time was studied with great care. The results of this investigation are contained in the following report made by Dr. Charles H. Judd, in charge of the Yale Psychological Laboratory.

## REPORT ON REACTION TIME.

In order to test the quickness of the members of the Hospital Corps Detachment, and to determine whether the changes in diet affected in any way their ability to respond promptly to sensory stimulation, each man was carried through a series of reaction experiments at the Yale Psychological Laboratory. method of the experiments was that regularly employed in simple reaction-time experiments. The person whose reaction time is to be measured is seated in a comfortable position with his finger pressing on an electric key. He is told that he is to lift his finger from the key as quickly as possible when he hears a given sound-signal. The sounder which produces this signal and the electric key are placed in a circuit with a standard time-measuring apparatus the Hipp Chronoscope. This chronoscope is arranged so that it begins to record the instant the sound is given and stops the instant the reactor lifts his finger. The dial of the chronoscope shows in thousandths of a second (hereafter designated by the technical term sigmas) the time that elapses between the sound to which the reactor is to respond and the movement of response. The chronoscope was tested at frequent intervals by means of a standard pendulum and errors in the record are well under two sigmas.

The time which is measured by this method is occupied chiefly by nervous processes. The following factors may be specified:

First, the auditory organ is aroused by the sound; second, the afferent nerve transmits the stimulation to the central nervous system; third, the central nervous system carries the energy to the motor nerves; fourth, the efferent motor nerves transmit the stimulus to the muscles; and finally, some time is required by the muscle for its contraction. It will be seen, accordingly, that the chief factors of a reaction are nervous processes, and since the external conditions of successive reactions are in all respects uniform, any variations in the time of a given person's reactions may be regarded as indicating variations in the nervous condition of the reactor. In view of the instability of nervous conditions. it is necessary to eliminate any slight or merely temporary fluctuations by taking each time a given reactor is tested a series of reactions. For this reason, a series of ten reactions was taken with each of the men every two weeks during November, December, and January. After an interval of two months, namely, on March 30, 31, and April 1, two final series were made with each man.

The results are presented in tables 1 to 5. The dates in the first column indicate the day on which each individual set of ten reactions was taken. The second column presents the averages of each ten reactions in sigmas. The third column gives the mean variations of the various single reaction times from the average. The fourth column gives the variations of the day's average, recorded in column 2, from the general average of that individual for all his experiments. These general averages are presented in table 6. At the bottom of each complete column of averages will be found the group average. This quantity is obtained by averaging the results from all the members of the squad for periods of, approximately, two weeks.

The mean variations in the third column make it clear that the subjects did not settle down into what could be regarded as trained subjects. Trained subjects are expected to give mean variations which are consistently within the limits of 10 per cent of the average. While there are, of course, instances in which the mean variation falls within this limit, there is no consistent exhibition of the regularity indicative of thorough practice. This fact is further confirmed by a comparison of the results of March 30, 31, and April 1 with those of the earlier months. At the end of March, all effects of practice in November, December, and January,

except the most general, may be regarded as having disappeared; and yet the averages and variations for the March and April dates resemble closely those of the month immediately preceding and also those of November. The effects of practice may, accordingly, be regarded as insignificant.

This lack of special training accounts for the large variations which appear in some cases. As is usual in reaction experiments, the signal to which the reactors were to respond was in each case preceded by about two seconds by a bell signal to arouse attention. Conditions were thus rendered as nearly uniform as possible, but the variations indicate in three or four cases exceptional lapses of attention. Such exceptional cases can be eliminated without prejudicing the final validity of the results by substituting the median for the averages. In table 7, the medians are grouped together and show even more than the tables of average the absence of any general variation during the period of the tests.

The obvious conclusion from these tests is that the quickness of the members of the squad underwent no general change during the whole period covered by the test. Individual members showed variations from time to time, but these variations are clearly accidental in character, for they show no regular tendencies and are in no way related to the changes in the character and amount of the diet.

(Signed) CHARLES H. JUDD.

April 12, 1904.

The following tables give all the data upon which the foregoing conclusions are based, being furnished by Dr. Judd and Mr. Warren M. Steele, Assistant in Psychology, by whom the observations were made. Dr. Cloyd N. McAllister, Instructor in Psychology, was also associated in the making of these observations.

TABLE 1. — OCTOBER, 1908.

Name.	Date.	Avg.	M. V.	Var. from G. A.	Date.	Avg.	M. V.	Var. from G. A.
I. Coffman  II. Henderson  III. Loewenthal  IV. Morris  V. Oakman  VI. Slincy  VII. Steltz  VIII. Zooman	17 17 17 17 17 17 22 17	207.4 179.9 216.9 227.7 222.7 262.4 167.7 264.9	34.1 31.5 43.5 19.8 65.4 83.1 16.3 32.1	8.7 21.2 7.8 8.7 4.8 49.9 20.4 50.0	  27 29 27	223.5 204.5 195.0	28.8 46.2 21.1	5.6 8.0 6.9
X. Broyles XI. Cohn		218.7						

TABLE 2. - NOVEMBER, 1903.

Name.	Date.	Avg.	M. V.	Var. from G. A.	Date.	Avg.	M. V.	Var. from G. A.	Date.	Avg.	M. V.	Var. from G. A.
I. Coffman .	6	303.7	37.9	87.6	20	215.9	22.8	0.5	ļ			
II. Henderson	5	248.5	72.9	47.4	19	198.2	22.0	2.9	<b> .</b> .	ا ا		
III. Loewenthal	18	190.1	28.7	84.6	30	229.7	37.4	5.0				
IV. Morris					25	248.6	45.0	17.2	ļ			
V. Oakman .	10	185.9	18.9	82.0	24	212.1	21.9	5.8				
VI. Sliney	12	200.1	25.8	11.0	26	209.2	24.6	3.3				
VII. Steltz	10	203.2	45.7	15.1	24	163.8	31.4	24.3				
VIII. Zooman .	5	192.0	29.2	22.9	19	188.8	23.4	81.1				
IX. Fritz	2	240.1	17.3	19.5	17	204.8	18.2	15.8	30	223.8	52.3	8.2
X. Broyles				l	17	184.7	23.0	8.2		'		
XI. Cohn	ļ		<b> </b>		28	279.1	50.7	70.3	ļ			
	_								L			
Group average .						211.8					• • •	

TABLE 8. — DECEMBER, 1903.

Name.	Date.	Avg.	M. v.	Var. from G. A.	Date.	Avg.	M. V.	Var. from G. A.	Date.	Avg.	M. V.	Var. from G. A.
I. Coffman	4	230.1	22.5	14.0	18	184.7	21.4	31.4		.,.		
II. Henderson					81	187.0	10.0	14.1	ļ			
III. Loewenthal	11	224.6	83.0	0.1	26	217.9	10.5	6.8				
IV. Morris	9	237.7	16.4	6.3	23	276.6	80.8	45.2				
V. Oakman	8	238.5	19.5	15.6	22	220.5	17.5	2.6	ļ			
VI. Sliney	10	218.8	25.4	5.8	24	197.1	24.1	15.4				
VII. Steltz	8	178.8	15.9	9.8	22	201.7	20.3	13.6				• •
VIII. Zooman	3	292.7	42.4	77.8	17	248.1	71.1	83.2	81	206.9	35.0	8.0
IX. Fritz	14	187.4	16.1	83.2	28	200.7	22.4	19.9				
X. Broyles	1	180.3	18.8	7.6	15	188.0	25.8	0.1				
XI. Cohn	7	243.8	111.1	34.5	21	177.0	16.2	31.8				٠.
Group averages		222.6	•••		 	209.0			_ 		• • •	

TABLE 4. - JANUARY AND FEBRUARY, 1904.

Name.	Date.	Avg.	M. V.	Var. from G. A.	Date.	Avg.	М. V.	Var. from G. A.	Date.	Атв.	M. V.	Var. from G. A.
I. Coffman	1	188.5	22.4	27.6	15	246.4	60.5	30.3	29 Feb.	198.2	14.2	17.9
II. Henderson	7	206.8	14.2	5.7	21	180.9	8.9	20.2	4	172.1	15.0	29.0
III. Loewenthal		208.0				201.8	1	12.9			10.0	20.0
IV. Morris	1	260.4		1 1	1	222.3			3	244.0	15.6	12.6
V. Oakman .	5				1	215.5			2	197.0		
VI. Sliney	1 -	186.6			1 .	167.9			l			
VII. Steltz	5	211.5	9.1	23.4	19	200.4	29.2	12.3	2	169.9	18.9	18.2
VIII. Zooman	7	237.6	106.2	22.7	21	197.0	13.0	17.9	4	164.5		50.4
IX. Fritz	11	240.5	78.9	19.9	25	183.9	15.9	36.7				
X. Broyles	12	211.6	33.3	23.7	26	208.6	21.7	15.7				
XI. Cohn	4	187.1	84.3	21.7	18	195.8	23.7	13.0	1	195.4	47.2	18.4
	_				L							
Group averages		215.8				202.3						

TABLE 5. - MARCH AND APRIL, 1904.

Name.	Date.	Avg.	K. V.	Var. from G. A.	Date.	Avg.	M. V.	Var. from G. A.
I. Coffman	30	191.9	13.7	24.2	1	194.4	25.9	21.7
II. Henderson	81	195.2	15.2	5.9	1	241.8	27.2	40.2
III. Loewenthal	81	231.9	45.9	7.2	1	291.0	49.6	66.3
IV. Morris	31	174.4	11.4	57.0	1	190.7	18.6	40.7
V. Oakman	31	228.2	47.2	5.3	1	226.3	52.0	8.4
VI. Sliney	30	239.3	47.1	16.8	31	289.6	11.8	17.1
VII. Steltz	30	193.5	34.0	5.4	1	198.3	29.3	5.2
VIII. Zooman	30	196.6	11.6	18.3	1	179.8	15.6	35.1
IX. Fritz	31	244.5	48.9	23.9	1	259.7	89.3	39.1
X. Broyles	30	180.0	19.4	0.1	1	182.6	23.6	5.3
XI. Cohn	81	210.8	21.7	2.0	1	181.7	12.0	27.1
Group averages		207.4				216.4	•••	

TABLE 6.

	Individual General Average.	Av. Var. from Gen. Av.		Individual General Average.	Av. Var. from Gen. Av.					
I. Coffman	216.1	26.4	VII. Steltz	188.1	14.0					
II. Henderson	201.1	20.7	VIII. Zooman	214.9	38.4					
III. Loewenthal	224.7	17.5	IX. Fritz	220.6	23.5					
IV. Morris	231.4	24.5	X. Broyles	187.9	8.8					
V. Oakman	217.9	11.0	XI. Cohn	208.8	26.7					
VI. Sliney	212.5	19.7								
General g	General group average 211.6									

TABLE 7

XI. Cobn.	.naibeM	:	:	251.5	197.0	179.0	180.5	187.0	:	190.5	:	206.5	184.0
	Date.		:	g	7	21	4	18	:		:	31	
X. Broyles.	Median.	:	152.0	:	178.0	187.5	203.5	204.0	:	:	182.0	:	186.5
Ã	.estaG		17		1	16	12	26		:	8	:	1
IX. Fritz.	.naibeM	:	199.0	214.0	179.5	188.0	209.0	184.0	:	:	:	211.0	245.5
	Date.		17	ಜ	14	28	Ξ	8	:	:	:	31	1
VIII. Zooman.	Median,	260.0	185.0	184.0	284.5	229.5	185.0	199.5	:	164.5	197.0	:	174.0
28	Date.	17	9	19	00	17	-	21	:	4	8	:	1
VII. Steltz.	Median,	168.0	173.5	174.5	178.0	199.5	210.0	193.0	:	168.0	189.5	:	185.0
28	Date.	17	2	24	30	22	9	10	:	83	೫	:	-
VI. Sliney.	Median.	272.0	191.0	201.5	210.0	188.0	180.5	171.0	:	:	219.5	237.0	:
· 55	Date.	83	12	88	91	24	14	88	:	:	8	31	:
V. Oakman.	Median.	171.0	180.5	199.0	234.0	227.0	239.5	209.5	:	193.0	:	216.5	204.6
õ	Date.	17	10	24	<b>∞</b>	22	9	19	:	63	<u>:</u>	31	1
IV. Morria.	.naibeM	238.0	:	224.5	230.0	283.6	251.0	222.5	:	243.0	:	173.0	183.0
Ř	Date.	17	:	25	6	23	9	8	:	တ	:	31	-
III. Loewenthal.	.naibeM	196.0	181.0	217.0	218.5	219.5	212.0	209.0	:	:	:	206.5	296.0
100	Date.	17	13	30	11	26	00	ន	:	:	:	31	-
II. Henderson.	Median.	181.5	214.0	198.0	:	185.5	204.5	181.5	:	172.5	:	201.0	237.0
Hem	Date.	17	10	19	:	81	-	21	:	4	:	81	-
I. Coffman.	.naibe <b>M</b>	207.4	305.5	216.0	224.5	176.0	178.5	204.4	201.5	:	192.0	:	196.0
8	Date.	17	8	8	4	18	_	15	83	:	စ္က	31	-
	<b>м</b> опср•	Oct.	Nov.	Nov.	Dec.	Dec.	Jan.	Jan.	Jan.	Feb.	Mar.	Mar.	Apr.

## CHARACTER OF THE BLOOD.

One question that naturally arises in considering the possible effects of a diminished proteid intake upon bodily health is whether a continued diminution of proteid food will have any influence upon the character and composition of the blood. It might be claimed, for example, that a lowering of the quantity of proteid food below the ordinarily accepted standards will eventually result in a deterioration in the character of the blood. Obviously, if such should prove to be the case, it would at once emphasize the necessity for higher standards of proteid feeding. Further, there might result marked changes in the hæmoglobin-content of the blood in connection with a lowered proteid metabolism long continued. thoughts in mind, a careful study of the blood of the soldiers has been made from time to time, with special reference to determining the number of erythrocytes and leucocytes in the fluid, attention also being paid to the percentage of hæmoglobin.

Four distinct observations were made, as a rule, upon each man, namely, in the months of October, December, January, The results are tabulated in the accompanying tables. Examination of these results shows that, as a rule, the number of erythrocytes, or red blood corpuscles, was somewhat increased during this period of lowered proteid feeding. We are not disposed, however, to lay very much stress upon this apparent increase, because it is not sufficiently marked to carry much weight, especially in view of the difficulties attending the obtaining of great accuracy in blood counts in general. Regarding the leucocytes, the figures are less definite, but may be fairly interpreted as indicating practically no appreciable change in the number of white corpuscles. Similarly, the hæmoglobin-content shows no distinct alteration. Hence, the conclusion is that the physiological economy practised by the soldiers during their six months' stay in New Haven, and especially the marked diminution in the amount of proteid food consumed, did not result in any deterioration of the blood, so far as it can be measured by the number of contained erythrocytes and leucocytes, and by the content of hæmoglobin.

# 284 PHYSIOLOGICAL ECONOMY IN NUTRITION

Date.	Erythrocytes per cmm.	Leucocytes per cmm.	Hæmoglobin per cent.
Oakman.			
Oct. 23, 1903	5,480,000	7,300	- 82
Dec. 1, 1908	6,000,000	9,500	82
Jan. 26, 1904	4,670,000	12,500	80-81
Mar. 22, 1904	6,560,000	6,000	84-85
Slinet.			
Oct. 22, 1903	5,450,000	11,500	85
Nov. 20, 1908	6,070,000	8,800	84
Jan. 19, 1904	4,058,000	11,100	76
Mar. 15, 1904	6,208,000	8,400	82
Bates.		•	
Oct. 18, 1903	5,088,000	12,000	77
Nov. 10, 1903	7,344,000	13,600	96
•	0,022,000	20,000	"
Coffman.			
Oct. 13, 1908	6,024,000	9,300	84
Nov. 10, 1908	7,544,000	8,600	94
Feb. 2, 1904	3,160,000	11,800	78
Mar. 18, 1904	5,568,000	10,000	74
Сони.			
Nov. 28, 1908	5,952,000	17,200	94
Feb. 2, 1904	6,000,000	13,600	87-88
Mar. 18, 1904	7,000,000	9,000	85-86
Loewenthal.		•	
Oct. 16, 1908	6,392,000	5,900	84
Nov. 20, 1908	6,780,000	5,000	85
Jan. 25, 1904	6,500,000	11,200	85
Mar. 22, 1904	7,000,000	10,200	86-87
Morris.	.,,		
Oct. 14, 1903	6,728,000	11,300	84-85
Nov. 10, 1903	6,620,000	8,100	90-91
Feb. 8, 1904	6,000,000	9,600	87–88
Mar. 15, 1904	5,000,000	10,200	85
	.,,	,	



Soldiers exercising in the gymnasium.

1 . 

Date.	Erythrocytes per cum.	Leucocytes per cmm.	Hæmoglobin per cent.
Steltz.			
Oct. 16, 1903	6,792,000	12,400	85-86
Nov. 20, 1903	5,500,000	18,800	88
Feb. 8, 1904	5,000,000	14,700	86–87
Mar. 15, 1904	7,000,000	14,800	85
Broyles.	l		
Nov. 24, 1908	5,310,000	9,200	89
Jan. 19, 1904	5,200,000	6,100	80
Mar. 15, 1904	5,600,000	8,800	85
Zooman.			
Oct. 22, 1903	6,024,000	9,300	91
Nov. 24, 1908	5,136,000	6,700	94
Feb. 8, 1904	7,760,000	16,000	87-88
Mar. 22, 1904	4,800,000	18,600	88
Davis.			
Oct. 13, 1903	4,160,000	5,700	86-87
Nov. 10, 1903	5,850,000	9,200	88
FRITZ.			
Nov. 2, 1908	4,776,000	9,800	87-88
Dec. 1, 1903	6,048,000	9,200	94
Jan. 19, 1904	5,848,000	10,000	84
Mar. 15, 1904	5,784,000	6,400	92
Henderson.			
Oct. 16, 1903	7,192,000	16,000	87
Nov. 20, 1908	5,760,000	10,200	84
Jan. 25, 1904	6,800,000	8,000	
Mar. 18, 1904	8,144,000	15,000	79–80

## GENERAL CONCLUSIONS.

Careful consideration of the foregoing data, taken in their entirety, must lead the unbiassed thinker to admit the possibilities of physiological economy in nutrition. That there is no real need for a daily diet containing 118 grams of proteid food seems clearly indicated. The members of the soldier detachment lived without discomfort for a period of five months on amounts of proteid food not more than one-half that called

for by the ordinary standard dietaries, and this without increasing the amount of non-nitrogenous food. Body-weight, nitrogen equilibrium, physical strength and vigor, ability to respond to sensory stimulation, the composition and general condition of the blood, all remained unimpaired under a daily diet involving the metabolism of only 7 to 8 grams of nitrogen per day and with a fuel value of less than 2800 calories per day.

Further, the practice of such economy led to marked improvement in the working of the neuro-muscular machinery, sufficiently noticeable to attract the attention of the men themselves, apart from the records of the dynamometer, etc. Indeed, it has been the universal feeling among all the subjects of experiment that they were less conscious of fatigue than formerly, or that they could do more work without the feeling of fatigue that is usually so conspicuous after heavy work, or long-continued muscular strain. We thus have for consideration an added factor, viz., the possible improvement of the physical condition of the body under a lowered proteid intake. This question, however, we shall discuss more fully It is enough for the present to simply emphasize the fact that with a greatly diminished proteid metabolism the body suffers no harm, the muscular machinery is as well able to perform its work as usual, and consequently there would seem to be no adequate reason why our daily dietary should be cumbered with such quantities of proteid matter as are generally considered necessary for health and strength.

There is one point of great importance in this connection that should not be overlooked, viz., whether the power of resistance toward disease is diminished in any way by a continued low proteid intake. This is surely a proper question, and one that must be carefully considered. Fortunately or unfortunately, we have no facts at our disposal. We have the belief, however, engendered by the results so far obtained, that there is no good ground for assuming the body to be any more susceptible to disease under conditions of low proteid metabolism than when supplied with an excess of proteid food. Indeed, it has been somewhat remarkable how free from all

troubles — even during a very trying winter — the subjects of this experiment have been. We believe that economy in the use of proteid food, curtailment of proteid metabolism to a degree commensurate with the real needs of the body, will prove helpful to health, but we have no convincing facts to present, — only the simple statement that all the men have been well and remarkably free from colds and other minor ailments all through the experiment.

It is a remarkable and suggestive fact that when a person has once practised physiological economy in his diet sufficiently long for it to have become in a measure a habit, he has no desire to return to a fuller dietary rich in proteid matter. This, it seems to the writer, is convincing proof that both body and mind are fully satisfied with the smaller amounts of food, and argues in favor of the latter being quite adequate for the physiological needs of the organism. In this connection, the writer presents a few lines received during the summer from one of the soldier detachment. Nine of these men, after completing their work at New Haven early in April, 1904, were detailed for service at St. Louis, and the letter which is quoted was written simply to ask concerning some photographs that had been promised them. In the letter, however, occur two or three sentences which are interesting and suggestive.

World's Fair Grounds, St. Louis, Mo., July 8, 1904.

#### Professor Russell H. Chittenden:

Dear Sir, — On behalf of the men that were undergoing the "Food Test" conducted by you last winter, I write these few lines asking whether we are entitled to any of the photographs that were taken of us in the Yale Gymnasium the last two days we were there. . . . The men are all in first-class condition as regards their physical condition, and are all very thankful to you. We eat very little meat now as a rule, and would willingly go on another test. Enclosed you will find a list of the men as follows: Private 1st Class Jonah Broyles; Private 1st Class William E. Coffman; Private 1st Class James D. Henderson; Private 1st Class Maurice D. Loewenthal; Private 1st Class

William Morris; Private 1st Class William F. Sliney; Private 1st Class John J. B. Steltz; Private 1st Class Ben Zooman; Private 1st Class William Oakman.

Trusting I may hear from you in the near future, I am, Very respectfully,

(Signed) JOHN J. B. STELTZ.

Medical Department Exhibit U. S. Army, World's Fair Station.

# Daily Dietary of the Soldier Detail from October 2, 1903, to April 4, 1904.

For the first two weeks of their stay in New Haven, the soldiers were given their ordinary army ration, which is rich in meat and consequently had a high content of proteid or The detachment had their own cook and helper, and their food was prepared for them as they had always been Further, they had at this time perfect accustomed to it. freedom as to the quantity of food to be eaten, the figures given in the earlier days representing their own choice of quantity. Later, by the beginning of the third week, the diet was modified somewhat by the introduction of other articles in place of meat, especially at breakfast, so that the total nitrogen intake was diminished in some degree, but the men were still allowed freedom as to quantity. From November to the close of the experiment in April, both the character and quantity of the food for each meal were prescribed, but great care was exercised to see that the men were fully satisfied. Changes were made gradually and no discomfort was felt, or at least no complaint was made, although the men were frequently questioned and encouraged to comment upon the dietary and to make suggestions.

The dietary, however, speaks for itself, and a careful perusal of the daily record, with reference both to the character of the food and the quantities employed, will give clearer and more exact information as to the changes introduced than any verbal description. The only statement that need be made is that the heavier proteid foods were greatly reduced in

amount, and replaced in a measure by the lighter carbohydrate foods. Finally, it may be said that while vegetable foods eventually predominated, there was at no time a complete change to a vegetable diet.

#### Friday, October 2, 1903.

- Breakfast. Beefsteak 222 grams, fried potatoes 234 grams, onions 34 grams, gravy 68 grams, bread 144 grams, coffee 679 grams, sugar 18 grams.
- Dinner. Beef 171 grams, boiled potatoes 350 grams, onions 55 grams, bread 234 grams, coffee 916 grams, sugar 27 grams.
- Supper. Corned beef 195 grams, potatoes 170 grams, onions 21 grams, bread 158 grams, coffee 450 grams, sugar 21 grams, fruit jelly 107 grams.

#### Saturday, October 3, 1903.

- Breakfast. Bacon 162 grams, fried cake 215 grams, bread 72 grams, sugar 21 grams, coffee 550 grams.
- Dinner. Roast beef 250 grams, gravy 133 grams, bread 234 grams, sugar 21 grams, coffee 667.
- Supper. Frankfurters 171 grams, bread 128 grams, milk 71 grams, sugar 21 grams, coffee 450 grams.

#### Sunday, October 4, 1903.

- Breakfast. Beefsteak 299 grams, onions 21 grams, gravy 175 grams, bread 222 grams, milk 83 grams, sugar 21 grams, coffee 491 grams.
- Dinner. Roast beef 221 grams, potatoes 517 grams, gravy 154 grams, bread 148 grams, pie 184 grams, sugar 18 grams, milk 46 grams, coffee 621 grams.
- Supper. Roast beef 96 grams, potatoes 260 grams, onions 32 grams, jam 92 grams, bread 32 grams, coffee 360 grams, milk 65 grams, sugar 18 grams.

## Monday, October 5, 1903.

- Breakfast. Bacon 185 grams, fried potatoes 277 grams, gravy 98 grams, bread 140 grams, coffee 538 grams, sugar 18 grams, milk 65 grams.
- Dinner. Cabbage 304 grams, corned beef 200 grams, potatoes 309 grams, bread 145 grams, milk 55 grams, sugar 18 grams, coffee 457 grams.
- Supper. Cabbage 130 grams, potatoes 248 grams, onions 27 grams, bacon 35 grams, bread 200 grams, butter 30 grams, milk 55 grams, coffee 500 grams, sugar 20 grams, blackberry jam 135 grams.

#### Tuesday, October 6, 1903.

- Breakfast. Bologna sausage 150 grams, bread 230 grams, butter 25 grams, milk 55 grams, sugar 20 grams, coffee 334 grams.
- Dinner.— Beans 180 grams, onions 27 grams, bacon 90 grams, bread 160 grams, milk 55 grams, sugar 30 grams, coffee 500 grams.
- Supper. Beans 70 grams, beef liver 160 grams, onions 100 grams, bread 182 grams, milk 56 grams, sugar 20 grams, coffee 500 grams.

#### Wednesday, October 7, 1903.

- Breakfast. Beefsteak 290 grams, gravy 116 grams, bread 142 grams, milk 55 grams, sugar 20 grams, coffee 500 grams.
- Dinner. Roast beef 240 grams, onions 20 grams, gravy 166 grams, bread 170 grams, milk 56 grams, sugar 20 grams, coffee 550 grams.
- Supper. Potatoes 280 grams, beef 110 grams, onions 32 grams, bread 185 grams, pie 60 grams, milk 55 grams, butter 35 grams, sugar 20 grams, coffee 500 grams, blackberry jam 60 grams.

#### Thursday, October 8, 1903.

- Breakfast. Meat 107 grams, eggs 120 grams, bread 117 grams, milk 55 grams, sugar 20 grams, coffee 500 grams.
- Dinner. Bacon 170 grams, cabbage 297 grams, potatoes 360 grams, bread 120 grams, milk 40 grams, sugar 12 grams, coffee 300 grams.
- Supper. Peaches 100 grams, bread 347 grams, butter 35 grams, milk 55 grams, sugar 52 grams, coffee 416 grams.

### Friday, October 9, 1903.

- Breakfast. Beef 120 grams, potatoes 220 grams, onions 50 grams, butter 35 grams, milk 55 grams, bread 175 grams, sugar 20 grams, coffee 500 grams.
- Dinner. Roast beef 203 grams, potatoes 143 grams, gravy 144 grams, bread 108 grams, sugar 18 grams, milk 55 grams, coffee 451 grams.
- Supper. Beef liver 188 grams, onions 93 grams, bacon 86 grams, bread 154 grams, butter 38 grams, sugar 19 grams, milk 55 grams, coffee 500 grams.

### Saturday, October 10, 1903.

- Breakfast. Eggs 86 grams, bacon 89 grams, potatoes 187 grams, bread 128 grams, milk 55 grams, sugar 18 grams, coffee 500 grams.
- Dinner. Fish 233 grams, bacon 65 grams, onions 49 grams, potatoes 140 grams, bread 226 grams, milk 55 grams, sugar 19 grams, coffee 465 grams.
- Supper. Hamburg steak 224 grams, onions 23 grams, butter 28 grams, bread 147 grams, pie 128 grams, milk 55 grams, sugar 18 grams, coffee 500 grams.

#### Sunday, October 11, 1903.

- Breakfast. Beefsteak 243 grams, bread 105 grams, milk 55 grams, sugar 18 grams, coffee 335 grams.
- Dinner. Roast pork 208 grams, turnips 159 grams, potatoes 201 grams, gravy 138 grams, apple pie 168 grams, bread 89 grams, milk 55 grams, sugar 18 grams, coffee 340 grams.
- Supper. Stewed peaches 235 grams, bread 291 grams, milk 55 grams, butter 28 grams, sugar 18 grams, coffee 475 grams.

#### Monday, October 12, 1903.

- Breakfast. Potatoes 275 grams, beef 131 grams, onions 37 grams, bread 135 grams, milk 50 grams, sugar 18 grams, coffee 350 grams.
- Dinner. Beans 850 grams, bacon 70 grams, onions 39 grams, pickles 39 grams, bread 147 grams, milk 55 grams, sugar 18 grams, coffee 500 grams.
- Supper. Frankfurters 149 grams, butter 28 grams, bread 149 grams, black-berry jam 63 grams, milk 55 grams, sugar 18 grams, coffee 500 grams.

#### Tuesday, October 13, 1903.

- Breakfast. Beef liver 149 grams, bacon 68 grams, bread 100 grams, milk 55 grams, sugar 19 grams, coffee 375 grams.
- Dinner. Roast beef 187 grams, potatoes 131 grams, gravy 167 grams, tomatoes 151 grams, bread 112 grams, milk 55 grams, sugar 18 grams, coffee 410 grams.
- Supper. Roast beef 140 grams, apple sauce 350 grams, bread 144 grams, butter 33 grams, milk 55 grams, sugar 18 grams, coffee 500 grams.

### Wednesday, October 14, 1903.

- Breakfast. Bacon 93 grams, apple sauce 299 grams, syrup 58 grams, bread 271 grams, milk 55 grams, sugar 85 grams, coffee 417 grams.
- Dinner. Hamburg steak 186 grams, potatoes 336 grams, gravy 100 grams, onions 37 grams, bread 187 grams, milk 55 grams, sugar 18 grams, coffee 350 grams.
- Supper. Beef 224 grams, potatoes 242 grams, onions 28 grams, prunes 147 grams, bread 135 grams, butter 28 grams, milk 55 grams, sugar 44 grams, coffee 500 grams.

### Thursday, October 15, 1903.

- Breakfast. Beef liver 159 grams, bacon 72 grams, bread 138 grams, milk 55 grams, sugar 18 grams, coffee 500 grams.
- Dinner. Cabbage 401 grams, bacon 156 grams, potatoes 201 grams, bread 121 grams, milk 55 grams, sugar 19 grams, coffee 480 grams.
- Supper. Bologna sausage 154 grams, rice 140 grams, eggs 18 grams, bread 138 grams, butter 28 grams milk 55 grams, sugar 63 grams, coffee 500 grams.

### Friday, October 16, 1903.

- Breakfast. Beefsteak 285 grams, bread 140 grams, milk 61 grams, sugar 20 grams, coffee 545 grams.
- Dinner. Fish 226 grams, potatoes 287 grams, tomatoes 135 grams, bread 128 grams, milk 55 grams, sugar 18 grams, coffee 500 grams.
- Supper. Pork sausage 244 grams, apple sauce 204 grams, bread 189 grams, butter 31 grams, milk 61 grams, sugar 20 grams, coffee 545 grams.

#### Saturday, October 17, 1903.

- Breakfast. Ham 183 grams, potatoes 298 grams, bread 115 grams, sugar 20 grams, milk 61 grams, coffee 545 grams.
- Dinner. Beef 204 grams, potatoes 290 grams, onions 18 grams, bread 145 grams, milk 61 grams, sugar 20 grams, coffee 375 grams.
- Supper. Roast beef 142 grams, apple sauce 112 grams, butter 35 grams, bread 183 grams, pie 104 grams, milk 61 grams, sugar 66 grams, coffee 545 grams.

#### Sunday, October 18, 1903.

- Breakfast. Hamburg steak 234 grams, onions 31 grams, bread 155 grams, milk 61 grams, sugar 20 grams, coffee 455 grams.
- Dinner. Chicken 326 grams, dressing 142 grams, potatoes 290 grams, tomatoes 453 grams, bread 122 grams, milk 61 grams, sugar 20 grams, coffee 545 grams.
- Supper. Apple sauce 244 grams, syrup 100 grams, bread 518 grams, milk 61 grams, sugar 20 grams, coffee 500 grams.

### Monday, October 19, 1903.

- Breakfast. Eggs 79 grams, bacon 43 grams, bread 127 grams, milk 61 grams, sugar 20 grams, coffee 514 grams.
- Dinner. Roast beef 214 grams, sweet potatoes 374 grams, tomatoes 305 grams, onions 23 grams, bread 140 grams, milk 61 grams, sugar 25 grams, coffee 545 grams.
- Supper. Roast beef 173 grams, apple sauce 214 grams, bread 163 grams, butter 30 grams, milk 61 grams, sugar 25 grams, coffee 509 grams.

### Tuesday, October 20, 1903.

- Breakfast. Oatmeal 316 grams, bread 95 grams, butter 19 grams, bacon 95 grams, coffee 600 grams, milk 245 grams, sugar 75 grams.
- Dinner. Roast beef 187 grams, boiled potatoes 366 grams, tomatoes 156 grams, bread 79 grams, coffee 600 grams, milk 101 grams, sugar 36 grams.
- Supper. Cold roast beef 176 grams, apple sauce 277 grams, bread 159 grams, butter 36 grams, coffee 370 grams, sugar 89 grams, milk 63 grams.

#### Wednesday, October 21, 1903.

- Breakfast. Fried oatmeal 142 grams, syrup 86 grams, bacon 62 grams, biscuits 155 grams, butter 35 grams, coffee 436 grams, milk 186 grams, sugar 46 grams.
- Dinner. Hamburg steak 275 grams, potatoes 399 grams, onions 63 grams, gravy 145 grams, bread 84 grams, coffee 500 grams, milk 140 grams, sugar 46 grams.
- Supper. Baked beans 836 grams, bread 148 grams, butter 43 grams, stewed prunes 193 grams, coffee 518 grams, milk 178 grams, sugar 48 grams.

#### Thursday, October 22, 1903.

- Breakfast. Boiled hominy 178 grams, French fried potatoes 168 grams, toasted bread 109 grams, butter 36 grams, coffee 473 grams, milk 163 grams, sugar 53 grams.
- Dinner. Corned beef 149 grams, boiled cabbage 191 grams, potatoes 189 grams, bread 87 grams, coffee 518 grams, sugar 51 grams, milk 76 grams.
- Supper. Bologna sausage 104 grams, Saratoga chips 69 grams, fried hominy 214 grams, syrup 91 grams, bread 75 grams, butter 86 grams, coffee 500 grams, sugar 40 grams, milk 91 grams.

#### Friday, October 23, 1903.

- Breakfast. Boiled rice with sugar and milk 221 grams, biscuits 158 grams, butter 38 grams, coffee 586 grams, milk 182 grams, sugar 71 grams.
- Dinner. Fish 288 grams, potatoes 265 grams, tomatoes 193 grams, bread I07 grams, coffee 545 grams, sugar 71 grams, milk 173 grams.
- Supper. Oyster stew with crackers 361 grams, apple sauce 102 grams, bread 43 grams, butter 35 grams, coffee 409 grams, sugar 46 grams, milk 309 grams.

### Saturday, October 24, 1903.

- Breakfast. Egg omelette 71 grams, with wheat flour 28 grams, bread 97 grams, butter 27 grams, coffee 545 grams, sugar 63 grams, milk 159 grams.
- Dinner. Hamburg steak made with bread 163 grams, fat 10 grams, and onions for flavor 90 grams, tomatoes 283 grams, bread 244 grams, butter 48 grams, coffee 454 grams, milk 182 grams, sugar 48 grams.
- Supper. Bacon 79 grams, potato chips 170 grams, stewed prunes 61 grams, biscuits 178 grams, butter 42 grams, coffee 545 grams, milk 182 grams, sugar 69 grams.

### Sunday, October 25, 1903.

- Breakfast. Apple 125 grams, fried rice 242 grams, syrup 64 grams, biscuits 127 grams, butter 33 grams, coffee 363 grams, milk 154 grams, sugar 31 grams.
- Dinner. Roast pork 252 grams, apple sauce 145 grams, potatoes 234 grams,

#### 294 PHYSIOLOGICAL ECONOMY IN NUTRITION

bread 66 grams, tapioca pudding 265 grams, coffee 363 grams, sugar 88 grams, milk 164 grams.

Supper. — Toasted bread 75 grams, blackberry jam 81 grams, bread 75 grams butter 46 grams, coffee 368 grams, milk 160 grams, sugar 46 grams.

#### Monday, October 26, 1903.

- Breakfast. Griddle cakes 305 grams, syrup 67 grams, bread 35 grams, coffee 454 grams, milk 145 grams, butter 23 grams, sugar 41 grams.
- Dinner. Beef stew with potatoes, onions, and thickened with corn starch 560 grams, bread 94 grams, milk 154 grams, coffee 454 grams, sugar 41 grams.
- Supper. Macaroni with cheese 226 grams, stewed tomatoes 282 grams, bread 114 grams, butter 41 grams, stewed prunes 127 grams, coffee 445 grams, milk 90 grams, sugar 20 grams.

#### Tuesday, October 27, 1903.

- Breakfast. Boiled rice, milk, and sugar 311 grams, toasted bread 114 grams, butter 31 grams, coffee 545 grams, milk 190 grams, sugar 79 grams.
- Dinner. Codfish-balls 369 grams, mashed potatoes 269 grams, pickles 43 grams, bread 72 grams, apple pie 117 grams, coffee 545 grams, milk 91 grams, sugar 25 grams.
- Supper. Apple-rice pudding 397 grams, biscuit 252 grams, butter 48 grams, coffee 500 grams, milk 91 grams, sugar 25 grams.

### Wednesday, October 28, 1903.

- Breakfast. Apple 252 grams, fried hominy 168 grams, syrup 86 grams, bread 79 grams, coffee 445 grams, milk 100 grams, sugar 25 grams.
- Dinner. Bean porridge with bread 415 grams, boiled onions 99 grams, coffee 545 grams, milk 91 grams, sugar 25 grams, bread 63 grams, bread pudding 282 grams.
- Supper. Apple fritters 371 grams, syrup 67 grams, biscuit 87 grams, butter 36 grams, pickles 23 grams, coffee 454 grams, milk 91 grams, sugar 25 grams.

### Thursday, October 29, 1903.

- Breakfast. Fried rice cakes 201 grams, syrup 54 grams, coffee 545 grams.
- Dinner. Hamburg steak\* with bread, fat, and onions 230 grams, boiled potatoes 211 grams, stewed tomatoes 257 grams, bread 61 grams, coffee 363 grams.
- Supper. Apple sauce 277 grams, biscuits 293 grams, butter 56 grams, apple pie 117 grams, tea 527 grams.

<sup>\*</sup> Hamburg steak contained 52 grams meat, 4 grams fat, 63 grams onions, 111 grams bread, each man eating 280 grams.

#### Friday, October 30, 1903.

- Breakfast. Boiled hominy 364 grams, sugar 47 grams, milk 91 grams, coffee 455 grams.\*
- Dinner. Fish 219 grams, French fried potatoes 158 grams, boiled onions 58 grams, bread pudding 338 grams, coffee 436 grams.
- Supper. Bacon 61 grams, Saratoga chips 119 grams, stewed prunes 206 grams, bread 155 grams, butter 58 grams, coffee 454 grams.

#### Saturday, October 31, 1903.

- Breakfast. Steamed oatmeal (soft) 349 grams, sugar 76 grams, milk 182 grams, biscuits 109 grams, butter 53 grams, coffee 409 grams.
- Dinner. Macaroni flavored with cheese 345 grams, stewed tomatoes 149 grams, bread 58 grams, apple pie 112 grams, coffee 416 grams.
- Supper. Boiled cabbage 278 grams, bologna sausage 159 grams, bread 79 grams, rice pudding 224 grams, coffee 500 grams.

#### Sunday, November 1, 1903.

- Breakfast. Apple 240 grams, rice croquettes 271 grams, syrup 67 grams, bread 41 grams, coffee 417 grams.
- Dinner. Roast pork 294 grams, apple sauce 217 grams, potatoes 352 grams, tapioca pudding 116 grams, coffee 417 grams.
- Supper. Biscuits 415 grams, butter 58 grams, blackberry jam 133 grams, pickles 33 grams, tea 416 grams.

### Monday, November 2, 1903.

- Breakfast. Steamed oatmeal 448 grams, milk 208 grams, sugar 65 grams, bread 70 grams, coffee 375 grams.
- Dinner. Beef stew 187 grams, potatoes 261 grams, onions 51 grams, thickened with corn starch 14 grams, bread 140 grams, coffee 500 grams.
- Supper. Macaroni 149 grams, stewed tomatoes 271 grams, pickles 72 grams, apple pie 109 grams, bread 189 grams, butter 53 grams, coffee 516 grams.

### Tuesday, November 3, 1903.

- Breakfast. Boiled rice 303 grams, sugar 72 grams, milk 225 grams, bread 67 grams, coffee 450 grams.
- Dinner. Baked bean porridge 326 grams, boiled onions 128 grams, potatoes 287 grams, bread 105 grams, coffee 508 grams.
- Supper. Boiled cabbage 217 grams, Saratoga chips 53 grams, stewed prunes 67 grams, fried rice 149 grams, syrup 58 grams, coffee 516 grams, bacon 37 grams, potatoes 179 grams.

<sup>\*</sup> One cup, total weight 454 grams, but containing small portions of milk and sugar.

#### Wednesday, November 4, 1903.

Breakfast. — Apple 250 grams, fried hominy 100 grams, syrup 50 grams, coffee 385 grams.

Dinner. — Hamburg steak with bread, fat, and onions 200 grams, boiled potatoes 250 grams, stewed tomatoes 250 grams, bread 75 grams, coffee 1 cup.

Supper. — Apple fritters 226 grams, syrup 50 grams, biscuit 95 grams, butter 30 grams, coffee 1 cup.

#### Thursday, November 5, 1903.

Breakfast. — Banana 114 grams, boiled rice \* 250 grams, with milk 181 grams, and sugar 76 grams, coffee 1 cup.

Dinner. — Macaroni and cheese 300 grams, bread 50 grams, apple sauce 200 grams, custard pie 112 grams, coffee 1 cup.

Supper. — Sausage 50 grams, potato chips 100 grams, stewed prunes 165 grams, bread 50 grams, butter 25 grams, coffee 1 cup.

#### Friday, November 6, 1903.

Breakfast. - Rice croquettes 200 grams, syrup 50 grams, coffee 1 cup.

Dinner. — Clam chowder with onions, tomatoes, and potatoes 350 grams, bread 75 grams, coffee 1 cup, tapioca-peach pudding 125 grams.

Supper. — Bread 127 grams, butter 40 grams, jam 125 grams, tea 1 cup.

#### Saturday, November 7, 1903.

Breakfast. — Soft oatmeal 300 grams, milk 150 grams, sugar 50 grams, bread 30 grams, coffee 1 cup.

Dinner. — Bean porridge with onions 294 grams, stewed prunes 66 grams, bread 75 grams, coffee 1 cup.

Supper. — Bread pudding 292 grams, stewed peaches 97 grams, crackers 50 grams, butter 25 grams, coffee 1 cup.

### Sunday, November 8, 1903.

Breakfast. — Apple 197 grams, stewed hominy 248 grams, milk 150 grams, sugar 50 grams, coffee 1 cup.

Dinner. — Beef stew thickened with corn starch, onions, and potatoes 405 grams, bread 75 grams, apple sauce 125 grams, coffee 1 cup.

Supper. — Pie 107 grams, chocolate 1 cup, biscuit 200 grams, butter 80 grams, stewed prunes 160 grams.

<sup>\*</sup> Rice or hominy on being moistened and cooked gains in weight 120 per cent; or rather, after the excess of moisture has evaporated and the rice is fried, it shows a gain of that amount. But for boiled rice or hominy, without drying or frying, there is an increase in weight of 230 per cent, as usually prepared.



Soldiers exercising in the gymnasium.

		٠	
		•	

#### Monday, November 9, 1903.

Breakfast. — Wheat griddle cakes 200 grams, syrup 40 grams, coffee 1 cup. Dinner. — Corned beef 125 grams, cabbage 200 grams, potatoes 175 grams, bread 75 grams, coffee 1 cup.

Supper. — Rice pudding 150 grams, stewed peaches 100 grams, crackers 50 grams, butter 25 grams, coffee 1 cup.

#### Tuesday, November 10, 1903.

- Breakfast. Toasted bread 200 grams, butter 40 grams, boiled egg 50 grams, coffee 1 cup.
- Dinner. Macaroni baked with cheese 300 grams, bread 50 grams, apple sauce 200 grams, custard pie 112 grams, coffee 1 cup.
- Supper. Bread pudding 300 grams, stewed peaches 100 grams, crackers 50 grams, butter 25 grams, coffee 1 cup.

#### Wednesday, November 11, 1903.

- Breakfast. Apple 196 grams, boiled rice 247 grams, milk 125 grams, sugar 50 grams, coffee 1 cup.
- Dinner. Hamburg steak with bread, fat, and onions 200 grams, boiled potatoes 250 grams, stewed tomatoes 250 grams, bread 75 grams, coffee 1 cup.
- Supper. Fried rice 100 grams, syrup 50 grams, biscuit 173 grams, butter 80 grams, tea 1 cup.

### Thursday, November 12, 1903.

- Breakfast. Banana 114 grams, toasted bread 179 grams, butter 50 grams, coffee 1 cup.
- Dinner.—Sausage 96 grams, French fried potatoes 200 grams, pickles 50 grams, bread 50 grams, apple and rice pudding 175 grams, coffee 1 cup.
- Supper. Boiled hominy 200 grams, milk 125 grams, sugar 47 grams, stewed prunes 109 grams, bread 50 grams, coffee 1 cup.

### Friday, November 13, 1903.

Breakfast. - Fried hominy 100 grams, syrup 50 grams, coffee 1 cup.

Dinner. — Clam chowder with onions, potatoes, and tomatoes 350 grams, bread 75 grams, coffee 1 cup.

Supper. — Biscuit 277 grams, butter 50 grams, jam 125 grams, sardines 85 grams, coffee 1 cup.

#### Saturday, November 14, 1903.

- Breakfast. Boiled rice 250 grams, milk 125 grams, sugar 50 grams, coffee 1 cup.
- Dinner. Beef stew with onions, potatoes, thickened with corn starch 850 grams, bread 75 grams, apple sauce 125 grams, coffee 1 cup.
- Supper. Rice croquettes 125 grams, syrup 40 grams, biscuit 175 grams, butter 25 grams, tea 1 cup,

#### Sunday, November 15, 1903.

- Breakfast. Apple 224 grams, soft boiled oatmeal 200 grams, milk 100 grams, sugar 40 grams, coffee 1 cup.
- Dinner. Macaroni and cheese 300 grams, stewed tomatoes 150 grams, bread 50 grams, pie 92 grams, coffee 1 cup.
- Supper. Fried bacon 30 grams, fried egg 40 grams, potato chips 100 grams, bread 50 grams, coffee 1 cup.

#### Monday, November 16, 1903.

- Breakfast. Wheat griddle cakes 150 grams, syrup 40 grams, coffee 1 cup. Dinner. - Corned beef 75 grams, cabbage 200 grams, mashed potatoes 200
- grams, bread 50 grams, coffee 1 cup.
- Supper. Rice pudding 150 grams, stewed peaches 100 grams, crackers 50 grams, butter 20 grams, coffee 1 cup.

### Tuesday, November 17, 1903.

- Breakfast. Indian-meal pudding 200 grams, milk 125 grams, coffee 1 cup.
- Dinner. Tomato soup with potatoes and onions boiled together 837 grams, bread 100 grams, sausage 44 grams, baked potato 200 grams, coffee 1 cup.
- Supper. Fried Indian-meal pudding 100 grams, syrup 50 grams, butter 35 grams, coffee 1 cup, apple sauce 100 grams, biscuit 150 grams.

### Wednesday, November 18, 1903.

- Breakfast. Boiled hominy 150 grams, milk 125 grams, sugar 80 grams, coffee
- Dinner. Bean soup (thick) 200 grams, pickles 85 grams, bread pudding 250 grams, bread 75 grams, coffee 1 cup, stewed peaches 75 grams.
- Supper. Fried hominy 150 grams, butter 25 grams, syrup 50 grams, bread 75 grams, stewed prunes 100 grams, coffee 1 cup.

### Thursday, November 19, 1903.

Breakfast. — Boiled rice 300 grams, milk 150 grams, sugar 50 grams, coffee 1 cup.

Dinner. — Hamburg steak made with plenty of chopped bread, fat, and onions 200 grams, baked potato 250 grams, bread 75 grams, stewed tomatoes 250 grams, coffee 1 cup.

Supper. — Biscuit 275 grams, butter 50 grams, apple sauce 175 grams, tea 1 cup.

#### Friday, November 20, 1903.

Breakfast. — Apple 200 grams, fried rice 150 grams, syrup 50 grams, coffee 1 cup.

Dinner. — Codfish-balls made with plenty of potatoes 200 grams, boiled onions 200 grams, bread 75 grams, apple pie 105 grams, coffee 1 cup.

Supper. — Banana fritters 200 grams, bread 75 grams, butter 50 grams, pickles 30 grams, coffee 1 cup.

#### Saturday, November 21, 1903.

Breakfast. — Wheat griddle cakes 200 grams, syrup 50 grams, bread 50 grams, coffee 1 cup.

Dinner. — Split pea soup 200 grams, bread 75 grams, pickles 30 grams, tapiocapeach pudding 150 grams, coffee 1 cup.

Supper. — Biscuit 275 grams, stewed prunes 100 grams, butter 50 grams, tea 1 cup.

### Sunday, November 22 1903.

Breakfast.—Apple 217 grams, Johnny cake made of corn meal 200 grams, butter 50 grams, coffee 1 cup.

Dinner.—Beef stew with onions, potatoes, and corn starch 350 grams, bread 75 grams, coffee 1 cup, pie 118 grams.

Supper. — Chocolate 1 cup, bread 150 grams, milk 800 grams.

### Monday, November 23, 1903.

Breakfast.—Boiled rice 300 grams, milk 125 grams, sugar 50 grams, coffee 1 cup.

Dinner. — Macaroni boiled 300 grams, stewed tomatoes 250 grams, bread 75 grams, pie 114 grams, coffee 1 cup.

Supper. — Fried rice 150 grams, syrup 50 grams, jam 75 grams, bread 75 grams, tea 1 cup.

#### Tuesday, November 24, 1903.

- Breakfast. Boiled hominy 150 grams, milk 125 grams, sugar 30 grams, coffee 1 cup, orange 200 grams.
- Dinner. Tomato soup with potatoes and onions boiled together 325 grams, bread 100 grams, fried sausage 50 grams, baked potato 200 grams, coffee 1 cup.
- Supper. Fried hominy 100 grams, syrup 50 grams, biscuit 150 grams, butter 35 grams, apple sauce 100 grams, coffee 1 cup.

#### Wednesday, November 25, 1903.

- Breakfast. Boiled Indian-meal 200 grams, milk 125 grams, coffee 1 cup, orange 225 grams.
- Dinner.— Split pea soup (thick) 200 grams, bread 75 grams, pickles 30 grams, apple pie 120 grams, coffee 1 cup.
- Supper. Bread pudding 250 grams, stewed peaches 100 grams, crackers 50 grams, butter 25 grams, tea 1 cup.

#### Thursday, November 26, 1903.

- Breakfast. Biscuit 250 grams, butter 50 grams, apple sauce 150 grams, coffee 1 cup.
- Dinner. Roast turkey (sliced) 100 grams, cranberry sauce 150 grams, mashed potatoes 150 grams, bread crumb stuffing 100 grams, boiled onions 200 grams, bread 75 grams, corn-starch pudding 125 grams, orange 200 grams, coffee 1 cup.
- Supper. Crackers 50 grams, tea 1 cup, stewed prunes 150 grams, butter 50 grams, wheat bread 100 grams.

### Friday, November 27, 1903.

- Breakfast.—Boiled rice 250 grams, milk 125 grams, sugar 50 grams, coffee 1 cup.
- Dinner.—Clam chowder with onions, potatoes, and tomatoes 350 grams, bread 75 grams, coffee 1 cup.
- Supper. Biscuit 275 grams, butter 50 grams, jam 125 grams, sardine 60 grams, coffee 1 cup.

#### Saturday, November 28, 1903.

- Breakfast.—Fried rice 100 grams, syrup 50 grams, coffee 1 cup, apple 200 grams.
- Dinner. Boiled macaroni 200 grams, stewed tomatoes 250 grams, bread 50 -grams, apple pie 150 grams, coffee 1 cup.
- Supper. Potato chips 100 grams, fried bacon 30 grams, bread 75 grams, jam 75 grams, tea 1 cup.

#### Sunday, November 29, 1903.

Breakfast. — Wheat griddle cakes 200 grams, syrup 50 grams, coffee 1 cup. Dinner. — Bean soup (thick) 200 grams, bread 75 grams, boiled potato 150 grams, bread pudding 250 grams, coffee 1 cup.

Supper. — Stewed peaches 100 grams, butter 85 grams, bread 75 grams, fried sausage 33 grams, coffee 1 cup.

#### Monday, November 30, 1903.

- Breakfast. Boiled oatmeal 200 grams, milk 125 grams, sugar 30 grams, coffee 1 cup.
- Dinner. Corned beef 75 grams, cabbage 200 grams, mashed potatoes 200 grams, bread 50 grams, coffee 1 cup.
- Supper. Rice pudding 150 grams, stewed peaches 100 grams, crackers 50 grams, butter 25 grams, coffee 1 cup.

#### Tuesday, December 1, 1903.

- Breakfast. Boiled hominy 150 grams, milk 125 grams, sugar 30 grams, coffee 1 cup.
- Dinner. Tomato soup with potatoes and onions boiled together 325 grams, bread 100 grams, shaved dried beef 30 grams, baked potato 147 grams, coffee 1 cup.
- Supper. Fried hominy 100 grams, syrup 50 grams, crackers 50 grams, butter 80 grams, coffee 1 cup, apple sauce 100 grams.

### Wednesday, December 2, 1903.

- Breakfast. Boiled Indian-meal 200 grams, milk 125 grams, coffee 1 cup, orange 200 grams.
- Dinner. Split pea soup (thick) 200 grams, bread 75 grams, pickles 30 grams, apple pie 125 grams, coffee 1 cup.
- Supper. Stewed peaches 100 grams, bread pudding 250 grams, crackers 50 grams, butter 25 grams, coffee 1 cup.

### Thursday, December 3, 1903.

Breakfast. — Wheat griddle cakes 200 grams, syrup 50 grams, coffee 1 cup.

Dinner. — Hamburg steak with bread, fat, and onions 150 grams, boiled potatoes 250 grams, stewed tomatoes 250 grams, bread 75 grams, coffee 1 cup.

Supper. — Boiled rice 150 grams, milk 125 grams, sugar 30 grams, coffee 1 cup.

#### Friday, December 4, 1903.

Breakfast. - Fried rice 100 grams, syrup 50 grams, coffee 1 cup.

Dinner. — Clam chowder with onions, tomatoes, and potatoes 350 grams, bread 75 grams, coffee 1 cup.

Supper. — Biscuit 275 grams, butter 50 grams, coffee 1 cup, jam 125 grams, sardines 75 grams.

#### Saturday, December 5, 1903.

Breakfast. — Boiled oatmeal 175 grams, milk 125 grams, sugar 30 grams, coffee

Dinner. —Boiled macaroni 200 grams, stawed tomatoes 250 grams, bread 50 grams, pie 117 grams, coffee 1 cup.

Supper. — Potato chips 100 grams, fried bacon 30 grams, corn-starch custard 125 grams, bread 40 grams, tea 1 cup.

#### Sunday, December 6, 1903.

Breakfast. — Banana 125 grams, toasted bread 150 grams, butter 50 grams, coffee 1 cup.

Dinner. — Sausage 50 grams, French fried potatoes 200 grams, pickles 30 grams, apple-rice pudding 200 grams, coffee 1 cup.

Supper. — Stewed prunes 150 grams, crackers 75 grams, butter 40 grams, coffee 1 cup.

### Monday, December 7, 1903.

Breakfast. — Corn-meal Johnny-cake 200 grams, butter 50 grams, coffee 1 cup.
 Dinner. — Bean soup (thick) 200 grams, boiled potatoes 200 grams, bread 75 grams, pie 146 grams, coffee 1 cup.

Supper. — Crackers 80 grams, milk 200 grams, stewed peaches 150 grams, coffee 1 cup.

### Tuesday, December 8, 1903.

Breakfast. — Boiled rice 150 grams, milk 125 grams, sugar 80 grams, coffee 1 cup.

Dinner. — Meat pie (made with a little meat, flour, etc.) 150 grams, mashed potato 250 grams, stewed prunes 100 grams, bread 75 grams, coffee 1 cup.

Supper. — Fried rice 100 grams, syrup 50 grams, bread 100 grams, cocoa 1 cup.

### Wednesday, December 9, 1903.

- Breakfast. Boiled oatmeal 175 grams, milk 100 grams, sugar 25 grams, coffee 1 cup.
- Dinner. Boiled macaroni without cheese 200 grams, stewed tomatoes 250 grams, bread 75 grams, pie 110 grams, coffee 1 cup.
- Supper. Bread 200 grams, milk 200 grams, stewed peaches 150 grams, cocoa 1 cup.

#### Thursday, December 10, 1903.

- Breakfast. Boiled hominy 125 grams, milk 100 grams, sugar 30 grams, coffee 1 cup.
- Dinner. Roast beef 50 grams, boiled potato 200 grams, bread 75 grams, boiled onions 100 grams, coffee 1 cup.
- Supper. Suet pudding 150 grams, stewed prunes 150 grams, crackers 50 grams, cocoa 1 cup.

#### Friday, December 11, 1903.

- Breakfast. Fried hominy 100 grams, syrup 50 grams, coffee 1 cup.
- Dinner. Clam chowder with onions, potatoes and tomatoes 850 grams, bread 100 grams, coffee 1 cup.
- Supper. Biscuit 275 grams, butter 50 grams, sardine 50 grams, jam 125 grams, cocoa 1 cup.

### Saturday, December 12, 1903.

- Breakfast. Griddle cakes (with egg) 200 grams, syrup 50 grams, coffee 1 cup. Dinner. Cold roast beef 50 grams, French fried potatoes 200 grams, applerice pudding 200 grams, pickles 30 grams, coffee 1 cup.
- Supper. Bread 100 grams, butter 50 grams, stewed prunes 150 grams, cocoa 1 cup.

### Sunday, December 13, 1903.

- Breakfast. Boiled Indian-meal 200 grams, milk 100 grams, coffee 1 cup.
- Dinner. Tomato soup with potatoes and onions boiled together 325 grams, bread 100 grams, shaved dried beef 12 grams, baked potato 110 grams, coffee 1 cup.
- Supper. Bread 150 grams, butter 50 grams, apple sauce 175 grams, cocoa 1 cup.

### Monday, December 14, 1903.

- Breakfast. Fried Indian-meal 100 grams, syrup 50 grams, coffee 1 cup.
- Dinner. Split-pea soup (thick) 175 grams, bread 75 grams, boiled onions 100 grams, mashed potato 150 grams, apple pie 121 grams, coffee 1 cup.
- Supper. Bread 100 grams, milk 200 grams, stewed peaches 150 grams, cocoa 1 cup.

#### Tuesday December 15, 1903.

- Breakfast. Boiled rice 150 grams, milk 125 grams, sugar 30 grams, coffee 1 cup.
- Dinner. Baked macaroni with a little cheese 200 grams, stewed tomatoes 200 grams, bread 50 grams, pie 115 grams, coffee 1 cup.
- Supper. Bread pudding 250 grams, stewed peaches 100 grams, crackers 50 grams, butter 15 grams, coffee 1 cup.

#### Wednesday, December 16, 1903.

- Breakfast. Fried rice 100 grams, syrup 50 grams, coffee 1 cup.
- Dinner. Hamburg steak with bread, fat, and onions 150 grams, boiled potatoes 200 grams, apple sauce 200 grams, bread 75 grams, coffee 1 cup.
- Supper. Biscuits 150 grams, butter 20 grams, stewed prunes 150 grams, tea 1 cup.

#### Thursday, December 17, 1903.

- Breakfast. Boiled hominy 150 grams, milk 125 grams, sugar 30 grams, coffee 1 cup.
- Dinner. Vegetable soup (thickened with flour) containing potatoes, onions, and tomatoes 300 grams, bread 75 grams, apple-rice pudding 150 grams, coffee 1 cup.
- Supper. Fried bacon 80 grams, baked potato 150 grams, bread 50 grams, butter 15 grams, coffee 1 cup.

#### Friday, December 18, 1903.

- Breakfast. Fried hominy 100 grams, syrup 50 grams, coffce 1 cup.
- Dinner. Clam chowder with onions, potatoes, and tomatoes 350 grams, bread 75 grams, coffee 1 cup.
- Supper. Shaved dried beef 30 grams, biscuit 150 grams, butter 20 grams, apple sauce 150 grams, tea 1 cup.

### Saturday, December 19, 1903.

- Breakfast.—Boiled Indian-meal 150 grams, milk 125 grams, sugar 80 grams, bread 35 grams, coffee 1 cup.
- Dinner. Corned beef 75 grams, cabbage 200 grams, boiled potatoes 175 grams, bread 40 grams, coffee 1 cup.
- Supper. Rice pudding with raisins 150 grams, stewed peaches 100 grams, crackers 50 grams, butter 15 grams, tea 1 cup.

#### Sunday, December 20, 1903.

Breakfast. \_\_Apple 150 grams, fried Indian-meal 100 grams, bread 35 grams, syrup 50 grams, coffee 1 cup.

Dinner. — Split-pea soup (thick) 150 grams, bread 40 grams, boiled carrots 100 grams, mashed potato 150 grams, apple pie 125 grams, coffee 1 cup.

Supper. — Sausage 50 grams, French fried potatoes 100 grams, bread 50 grams, butter 15 grams, tea 1 cup.

#### Monday, December 21, 1903.

Breakfast. — Wheat griddle cakes 150 grams, syrup 40 grams, coffee 1 cup.

Dinner. — Beef stew (with onions and potatoes, thickened with corn starch) 350 grams, bread 75 grams, stewed prunes 125 grams, coffee 1 cup.

Supper. — Suet pudding 150 grams, apple sauce 125 grams, tea 1 cup.

#### Tuesday, December 22, 1903.

Breakfast. — Apple 150 grams, boiled rice 150 grams, milk 125 grams, sugar 30 grams, bread 80 grams, coffee 1 cup.

Dinner. — Baked macaroni with cheese 200 grams, stewed tomatoes 200 grams, bread 50 grams, pie 110 grams, coffee 1 cup.

Supper. — Bread pudding 250 grams, stewed peaches 100 grams, crackers 50 grams, butter 15 grams, tea 1 cup.

### Wednesday, December 23, 1903.

Breakfast. — Fried rice 100 grams, syrup 50 grams, crackers 30 grams, butter 10 grams, coffee 1 cup.

Dinner — Bean soup (thick) 200 grams, bread 75 grams, pickles 80 grams, tapioca-peach pudding 150 grams, coffee 1 cup.

Supper. — Apple fritters 200 grams, stewed prunes 125 grams, bread 50 grams, butter 15 grams, tea 1 cup.

### Thursday, December 24, 1903.

Breakfast. — Apple 200 grams, boiled hominy 150 grams, milk 125 grams, sugar 30 grams, coffee 1 cup.

Dinner — Tomato soup with potatoes and onions boiled together 325 grams, fried sausage 40 grams, bread 75 grams, baked potato 150 grams, coffee 1 cup.

Supper — Biscuit 175 grams, butter 30 grams, jam 125 grams, tea 1 cup.

#### Friday, December 25, 1903.

- Breakfast. Apple 200 grams, fried hominy 130 grams, syrup 50 grams, coffee 1 cup.
- Dinner.—Roast turkey 100 grams, bread-crumb stuffing 100 grams, cranberry sauce 150 grams, boiled onions 200 grams, mashed potatoes 150 grams, bread 75 grams, orange 200 grams, corn-starch custard 125 grams, coffee 1 cup.
- Supper. Bread 100 grams, crackers 25 grams, stewed prunes 150 grams, butter 40 grams, tea 1 cup.

#### Saturday, December 26, 1903.

- Breakfast. Boiled rice 200 grams, milk 125 grams, sugar 30 grams, coffee 1 cup.
- Dinner. Clam chowder with onions, potatoes, and tomatoes 350 grams, bread 75 grams, pickles 85 grams, coffee 1 cup.
- Supper. Potato chips 100 grams, fried bacon 25 grams, bread 75 grams, jam 75 grams, tea 1 cup.

### Sunday, December 27, 1903.

- Breakfast. Wheat griddle cakes 200 grams, syrup 50 grams, coffee 1 cup, apple 200 grams.
- Dinner.—Split-pea soup (thick) 200 grams, bread 75 grams, stewed tomatoes 150 grams, pie 115 grams, coffee 1 cup.
- Supper. Suet pudding (plum duff) 150 grams, apple sauce 125 grams, tea 1 cup.

### Monday, December 28, 1903.

- Breakfast.—Boiled Indian-meal 150 grams, milk 125 grams, sugar 30 grams, coffee 1 cup, apple 200 grams.
- Dinner. Corned beef 75 grams, cabbage 200 grams, mashed potatoes 200 grams, bread 50 grams, coffee 1 cup.
- Supper. Rice pudding 150 grams, stewed peaches 100 grams, crackers 40 grams, butter 15 grams, tea 1 cup.

### Tuesday, December 29, 1903.

- Breakfast. Fried Indian-meal 100 grams, syrup 50 grams, coffee 1 cup, apple 210 grams.
- Dinner. Boiled macaroni 200 grams, stewed tomatoes 250 grams, bread 50 grams, apple pie 140 grams, coffee 1 cup.
- Supper. Potato chips 100 grams, fried bacon 35 grams, bread 75 grams, jam 75 grams, tea 1 cup.

#### Wednesday, December 30, 1903.

Breakfast. — Boiled rice 150 grams, milk 125 grams, sugar 30 grams, coffee 1 cup, apple 150 grams, baked potato 100 grams.

Dinner. — Hamburg steak with bread, fat, and onions 150 grams, boiled potatoes 200 grams, apple sauce 200 grams, bread 75 grams, coffee 1 cup.

Supper. — Biscuit 160 grams, butter 20 grams, stewed prunes 150 grams, tea 1 cup.

#### Thursday, December 31, 1903.

Breakfast. — Apple 175 grams, fried rice 100 grams, syrup 56 grams, coffee 1 cup, baked potato 90 grams.

Dinner. — Split-pea soup (thick) 200 grams, bread 75 grams, pickles 80 grams, boiled potato 100 grams, coffee 1 cup, pie 115 grams.

Supper. — Bread pudding 250 grams, stewed peaches 100 grams, crackers 20 grams, butter 10 grams, tea 1 cup.

#### Friday, January 1, 1904.

Breakfast. — Wheat griddle cakes 200 grams, syrup 50 grams, apple 175 grams, coffee 1 cup, baked potato 100 grams.

Dinner. — Clam chowder with onions, potatoes, and tomatoes 350 grams, bread 75 grams, ice cream 200 grams, coffee 1 cup.

Supper. — Suet pudding 150 grams, apple sauce 125 grams, crackers 25 grams, tea 1 cup.

### Saturday, January 2, 1904.

Breakfast. — Boiled rice 150 grams, milk 125 grams, sugar 30 grams, coffee 1 cup, apple 190 grams, baked potato 100 grams.

Dinner.—Baked macaroni with small amount of cheese 200 grams, stewed tomatoes 200 grams, bread 50 grams, pie 115 grams, coffee 1 cup.

Supper. — Biscuit 125 grams, butter 20 grams, fried bacon 80 grams, baked potato 150 grams, stewed prunes 150 grams, tea 1 cup.

### Sunday, January 3, 1904.

Breakfast.—Apple 190 grams, fried rice 100 grams, syrup 50 grams, baked potato 90 grams, butter 10 grams, coffee 1 cup.

Dinner. — Baked potato 150 grams, dried beef 50 grams, stewed with milk 50 grams, bread 75 grams, butter 20 grams, coffee 1 cup, pickles 20 grams, tapioca-peach pudding 125 grams.

Supper. — Apple fritters 200 grams, stewed prunes 125 grams, bread 50 grams, butter 15 grams, tea 1 cup.

### Monday, January 4, 1904.

Breakfast. — Boiled Indian-meal 150 grams, milk 125 grams, sugar 30 grams, bread 35 grams, coffee 1 cup.

Dinner.—Tomato soup with potatoes and onions boiled together 325 grams, bread 75 grams, coffee 1 cup, bread pudding 150 grams.

Supper. — Biscuit 175 grams, butter 80 grams, jam 125 grams, tea 1 cup.

#### Tuesday, January 5, 1904.

Breakfast.—Boiled hominy 150 grams, milk 125 grams, sugar 30 grams, baked potato 150 grams, butter 10 grams, coffee 1 cup.

Dinner. — Split-pea soup (thick) 800 grams, bread 75 grams, pickles 80 grams, coffee 1 cup, pie 100 grams.

Supper. — Fried bacon 30 grams, potato chips 100 grams, bread 75 grams, jam 60 grams, tea 1 cup.

#### Wednesday, January 6, 1904.

Breakfast. — Fried hominy 100 grams, syrup 50 grams, coffee 1 cup, apple 200 grams.

Dinner. — Hamburg steak with plenty of bread, fat, and onions 150 grams, boiled potatoes 200 grams, apple sauce 200 grams, bread 75 grams, coffee 1 cup.
 Supper. — Biscuit 150 grams, butter 20 grams, stewed prunes 150 grams, tea 1 cup.

### Thursday, January 7, 1904.

Breakfast. — Apple 190 grams, boiled rice 150 grams, milk 125 grams, sugar 30 grams, coffee 1 cup, baked potato 100 grams.

Dinner. — Baked macaroni with small amount of cheese 200 grams, stewed tomatoes 200 grams, bread 50 grams, coffee 1 cup, pie 130 grams.

Supper. — Suet pudding 150 grams, apple sauce 125 grams, crackers 25 grams, tea 1 cup.

### Friday, January 8, 1904.

Breakfast. — Apple 150 grams, wheat griddle cakes 200 grams, syrup 50 grams, coffee 1 cup.

Dinner. — Fish-balls with creamed potatoes 150 grams, stewed tomatoes 200 grams, bread 75 grams, coffee 1 cup, tapioca-peach pudding 125 grams.

Supper. — Biscuit 150 grams, butter 20 grams, stewed peaches 150 grams, tea 1 cup, baked potato 100 grams.

### Saturday, January 9, 1904.

Breakfast. — Apple 200 grams, boiled rice 150 grams, milk 125 grams, sugar 80 grams, butter 10 grams, baked potato 100 grams, coffee 1 cup.

Dinner. — Baked potato 150 grams, dried beef 50 grams, stewed with milk 20 grams, bread 75 grams, butter 20 grams, pickles 20 grams, coffee 1 cup.

Supper. — Apple fritters 200 grams, stewed prunes 125 grams, bread 50 grams, butter 15 grams, tea 1 cup.

### Sunday, January 10, 1904.

Breakfast. — Fried rice 100 grams, syrup 50 grams, coffee 1 cup, apple 250 grams.

Dinner.—Tomato soup with potatoes and onions boiled together 325 grams, bread 75 grams, bread pudding 150 grams, coffee 1 cup.

Supper. — Toasted bread 100 grams, butter 20 grams, sardine 25 grams, stewed prunes 150 grams, tea 1 cup.

### Monday, January 11, 1904.

Breakfast. — Boiled Indian-meal 150 grams, milk 125 grams, sugar 30 grams, bread 35 grams, butter 10 grams, coffee 1 cup.

Dinner. — Boiled fresh beef 75 grams, boiled cabbage 200 grams, mashed potatoes 200 grams, bread 50 grams, coffee 1 cup.

Supper. — Rice pudding 150 grams, stewed peaches 100 grams, crackers 30 grams, butter 10 grams, tea 1 cup.

#### Tuesday, January 12, 1904.

Breakfast.—Fried Indian-meal 100 grams, syrup 50 grams, coffee 1 cup = 350 grams or 867 cc., bread 50 grams, butter 15 grams.

Dinner. — Boiled macaroni 250 grams, stewed tomatoes 250 grams, bread 75 grams, coffee 1 cup, 367 grams.

Supper. — Potato chips 100 grams, fried bacon 25 grams, bread 75 grams, jam 75 grams, tea 1 cup = 850 cc.

Total nitrogen, 7.793 grams.

Fuel value, 2404 calories.

### Wednesday, January 13, 1904.

Breakfast. — Boiled rice 150 grams, milk 130 cc. 125 grams, sugar 30 grams, butter 10 grams, bread 30 grams, coffee 1 cup = 350 cc.

Dinner. — Hamburg steak with plenty of bread, fat, and onions chopped together 150 grams, boiled potatoes 200 grams, apple sauce 200 grams, bread 75 grams, coffee 1 cup = 350 cc.

Supper. — Fried rice 100 grams, syrup 50 grams, tea 350 cc., bread 50 grams, butter 15 grams.

Total nitrogen, 9.992 grams.

Fuel value, 2133 calories.

### Thursday, January 14, 1904.

Breakfast. — Boiled hominy 150 grams, milk 125 grams, sugar 30 grams, butter 10 grams, bread 30 grams, coffee 1 cup = 850 cc.

Dinner. — Split-pea soup (thick) 200 grams, bread 75 grams, mashed potatoes 100 grams, pickles 30 grams, coffee 1 cup = 350 cc., pie 120 grams.

Supper. — Suet pudding 150 grams, apple sauce 125 grams, crackers 25 grams, tea 1 cup = 350 cc.

Total nitrogen, 7.412 grams.

Fuel value, 2000 calories.

#### Friday, January 15, 1904.

- Breakfast. Wheat griddle cakes 200 grams, syrup 50 grams, coffee 1 cup = 350 cc.
- Dinner. Codfish-balls (4 parts potato, 1 part fish, fried in pork fat) 158 grams, stewed tomatoes 200 grams, bread 75 grams, coffee 1 cup = 350 cc., apple pie 95 grams.
- Supper. Apple fritters 200 grams, stewed prunes (stones not included) 125 grams, bread 50 grams, butter 15 grams, tea 1 cup = 350 cc.

Total nitrogen, 8.560 grams.

Fuel value, 2030 calories.

### Saturday, January 16, 1904.

- Breakfast.—Soft oatmeal 150 grams, milk 100 grams, sugar 30 grams, bread 30 grams, butter 10 grams, coffee 1 cup = 350 cc.
- Dinner. Baked macaroni with a little cheese 200 grams, stewed tomatoes 200 grams, bread 50 grams, tapioca-peach pudding 150 grams, coffee 1 cup = 350 cc.
- Supper.—French fried potatoes 100 grams, fried bacon 20 grams, bread 75 grams, jam 75 grams, tea 1 cup = 350 cc.

Total nitrogen, 7.282 grams.

Fuel value, 1824 calories.

#### Sunday, January 17, 1904.

- Breakfast. Boiled Indian-meal 125 grams, milk 125 grams, sugar 30 grams, butter 10 grams, bread 30 grams, coffee 1 cup = 350 cc.
- Dinner.—Bean soup (thick) 200 grams, bread 75 grams, mashed potato 100 grams, pickles 25 grams, coffee 1 cup = 350 cc., custard pie 105 grams.
- Supper. Crackers 50 grams, butter 15 grams, stewed prunes (without stones) 125 grams, sponge cake 100 grams, tea 350 cc.

Total nitrogen, 8.849 grams.

Fuel value, 2081 calories.

### Monday, January 18, 1904.

- Breakfast. Apple 150 grams, fried Indian-meal 100 grams, syrup 50 grams, baked potato 90 grams, butter 10 grams, coffee 1 cup.
- Dinner.—Beef stew with potatoes and onions thickened with corn starch 300 grams, bread 75 grams, coffee 1 cup.
- Supper. Bread pudding 250 grams, stewed peaches 100 grams, crackers 25 grams, butter 10 grams, tea 1 cup.

### Tuesday, January 19, 1904.

- Breakfast. Apple 180 grams, boiled rice 150 grams, milk 125 grams, sugar 30 grams, baked potato 100 grams, butter 10 grams, coffee 1 cup.
- Dinner. Tomato soup with potatoes and onions boiled together 350 grams, bread 75 grams, mashed potatoes 150 grams, pickles 30 grams, coffee 1 cup, pie 129 grams.
- Supper. Biscuit 125 grams, butter 20 grams, apple sauce 175 grams, crackers 16 grams, tea 1 cup.

### Wednesday, January 20, 1904.

- Breakfast. Apple 150 grams, fried rice 100 grams, syrup 50 grams, biscuit 50 grams, butter 10 grams, coffee 1 cup.
- Dinner. Baked beans with a little salt pork 150 grams, bread 75 grams, boiled onions 100 grams, coffee 1 cup.
- Supper. French fried potatoes 100 grams, fried bacon 20 grams, bread 50 grams, butter 10 grams, tea 1 cup.

### Thursday, January 21, 1904.

- Breakfast. Wheat griddle cakes 150 grams, syrup 50 grams, butter 10 grams, bread 50 grams, coffee 1 cup.
- Dinner. Mashed potatoes 200 grams, fried egg 33 grams, bread 75 grams, butter 15 grams, apple pie 117 grams, coffee 1 cup.
- Supper. Crackers 50 grams, butter 10 grams, stewed prunes 125 grams, tea 1 cup, sponge cake 50 grams.

### Friday, January 22, 1904.

- Breakfast. Boiled Indian-meal 150 grams, milk 125 grams, sugar 30 grams, coffee 1 cup, baked potato 100 grams, butter 10 grams.
- Dinner. Clam chowder with onions, potatoes, and tomatoes 200 grams, bread 75 grams, mashed potato 100 grams, coffee 1 cup.
- Supper. Apple fritters 200 grams, jam 75 grams, tea 1 cup, gingerbread 30 grams.

### Saturday, January 23, 1904.

- Breakfast. Boiled rice 150 grams, milk 125 grams, sugar 30 grams, coffee 1 cup, butter 10 grams, baked potato 150 grams.
- Dinner. Boiled macaroni 250 grams, stewed tomatoes 250 grams, bread 75 grams, coffee 1 cup.
- Supper. French fried potatoes 125 grams, fried bacon 30 grams, bread 100 grams, jam 75 grams, tea 1 cup.

### Sunday, January 24, 1904.

- Breakfast.—Fried rice 150 grams, syrup 50 grams, baked potato 125 grams, coffee 1 cup, apple 150 grams.
- Dinner. Split-pea soup 200 grams, bread 100 grams, pickles 30 grams, mashed potatoes 200 grams, coffee 1 cup, apple pie 100 grams.
- Supper. Baked apple with sugar 150 grams, crackers 50 grams, butter 20 grams, tea 1 cup, potato chips 50 grams.

#### Monday, January 25, 1904.

Breakfast. - Boiled hominy 150 grams, milk 125 grams, sugar 30 grams, coffee 1 cup, baked potato 100 grams, butter 10 grams.

Dinner. — Meat pie 200 grams (with 30 grams meat), boiled cabbage 200 grams, mashed potatoes 200 grams, coffee 1 cup.

Supper. - Sponge cake 100 grams, stewed peaches 100 grams, crackers 25 grams, butter 10 grams, tea 1 cup, baked potato 115 grams.

#### Tuesday, January 26, 1904.

Breakfast. - Apple 140 grams, wheat griddle cakes 200 grams, syrup 50 grams, coffee 1 cup, bread 50 grams, butter 10 grams.

Dinner. - Baked beans with a little salt pork 150 grams, boiled onions 100 grams, bread 75 grams, bread pudding 150 grams, coffee 1 cup.

Supper. — Biscuit 175 grams, butter 20 grams, apple sauce 125 grams, tea 1 cup.

### Wednesday, January 27, 1904.

Breakfast. - French fried potatoes 150 grams, fried bacon 20 grams, bread 75 grams, butter 10 grams, coffee 1 cup, apple 170 grams.

Dinner. - Split-pea soup (thick) 200 grams, bread 75 grams, mashed potatoes 100 grams, boiled onions 150 grams, coffee 1 cup.

Supper. — Rice pudding with raisins 200 grams, apple sauce 150 grams, crackers 25 grams, tea 1 cup.

### Thursday, January 28, 1904.

Breakfast. — Fried rice 100 grams, syrup 50 grams, bread 50 grams, coffee 1 cup, banana 75 grams.

Dinner. — Baked potato 170 grams, dried beef 40 grams, stewed with milk 30 grams, bread 75 grams, butter 20 grams, pickles 20 grams, coffee 1 cup.

Supper. — Apple fritters 200 grams, stewed prunes 125 grams, bread 50 grams, butter 15 grams, peach pie 120 grams, tea 1 cup.

### Friday, January 29, 1904.

Breakfast. — Wheat griddle cakes 200 grams, syrup 50 grams, coffee 1 cup, French fried potatoes 50 grams.

Dinner. — Oyster soup \* 800 grams, crackers 50 grams, bread 100 grams, coffee 1 cup, salad made of lettuce leaves, chopped apple, and celery with oil, salt, and pepper 75 grams.

Supper. — Biscuit 175 grams, butter 20 grams, jam 125 grams, tea 1 cup, apple 175 grams.

In each bowl of soup were 90 grams of oysters, 20 grams of butter, and 190 grams of milk.

#### Saturday, January 30, 1904.

- Breakfast. Boiled oatmeal 125 grams, milk 100 grams, sugar 25 grams, coffee 1 cup, baked potato 50 grams, butter 10 grams.
- Dinner. French fried potatoes 200 grams, cold roast beef 40 grams, bread 75 grams, pickles 30 grams, carrots 125 grams, coffee 1 cup.
- Supper. Apple-rice pudding 200 grams, stewed prunes 150 grams, bread 100 grams, butter 30 grams, tea 1 cup.

### Sunday, January 31, 1904.

- Breakfast. Boiled Indian-meal 200 grams, milk 100 grams, sugar 25 grams, coffee 1 cup, bread 50 grams.
- Dinner. Tomato soup with potatoes and onions boiled together 325 grams, bread 100 grams, fried egg 30 grams, baked potato 140 grams, coffee 1 cup.
- Supper. Lettuce-apple-celery salad 100 grams, bread 100 grams, butter 20 grams, stewed peaches 150 grams, tea 1 cup.

#### Monday, February 1, 1904.

- Breakfast. Fried Indian-meal 100 grams, syrup 50 grams, bread 50 grams, butter 10 grams, coffee 1 cup.
- Dinner. Baked beans 120 grams, with salt pork 30 grams, boiled onions 100 grams, mashed potatoes 200 grams, bread 75 grams, apple pie 120 grams, coffee 1 cup.
- Supper. Apple fritters 200 grams, stewed prunes 125 grams, crackers 82 grams, butter 15 grams, tea 1 cup.

### Tuesday, February 2, 1904.

- Breakfast. Fried hominy 100 grams, syrup 50 grams, apple 160 grams, bread 50 grams, butter 10 grams, coffee 1 cup.
- Dinner. Boiled macaroni 250 grams, stewed tomatoes 250 grams, bread 75 grams, pie 120 grams, coffee 1 cup.
- Supper. Biscuit 175 grams, butter 20 grams, crackers 25 grams, stewed prunes 125 grams, tea 1 cup.

### Wednesday, February 3, 1904.

- Breakfast. French fried potatoes 150 grams, fried bacon 20 grams, bread 50 grams, butter 10 grams, coffee 1 cup.
- Dinner.—Corned beef 40 grams, cabbage 200 grams, mashed potatoes 200 grams, bread 75 grams, coffee 1 cup, boiled rice 200 grams, syrup 50 grams.
- Supper. Lettuce-apple-celery salad 150 grams, bread 100 grams, butter 20 grams, jam 75 grams, tea 1 cup.

#### Thursday, February 4, 1904.

- Breakfast.—Wheat griddle cakes 200 grams, syrup 50 grams, coffee 1 cup, baked potato 150 grams, butter 10 grams.
- Dinner. Barley broth with potatoes and onions 250 grams, wheat flour dumplings 150 grams, boiled turnips 200 grams, bread 75 grams, tapiocapeach pudding 200 grams, coffee 1 cup.
- Supper. Suet pudding 150 grams, baked apple with sugar 150 grams, crackers 25 grams, stewed prunes 150 grams, tea 1 cup.

### Friday, February 5, 1904.

- Breakfast. Banana 100 grams, French fried potatoes 200 grams, biscuit 175 grams, butter 20 grams, coffee 1 cup.
- Dinner. Boiled codfish 60 grams, mashed potatoes 250 grams, boiled onions 200 grams, bread 75 grams, coffee 1 cup.
- Supper. Bread pudding with raisins 250 grams, stewed peaches 150 grams, crackers 25 grams, butter 10 grams, tea 1 cup.

#### Saturday, February 6, 1904.

- Breakfast. Boiled rice 175 grams, milk 125 grams, sugar 25 grams, coffee 1 cup, banana 90 grams.
- Dinner. Baked beans 70 grams, with salt pork 30 grams, bread 75 grams, boiled sweet potato 150 grams, butter 10 grams, coffee 1 cup, apple pie 100 grams.
- Supper. French fried potatoes 125 grams, celery-lettuce-apple salad 150 grams, bread 100 grams, butter 20 grams, jam 75 grams, tea 1 cup.

### Sunday, February 7, 1904.

- Breakfast. Fried rice 150 grams, syrup 50 grams, baked potato 140 grams, butter 10 grams, coffee 1 cup.
- Dinner.—Hamburg steak with much bread, fat, and onions 150 grams, boiled potato 150 grams, butter 10 grams, bread 75 grams, coffee 1 cup.
- Supper. Tapioca-peach pudding 250 grams, sponge cake 75 grams, tea 1 cup.

#### Monday, February 8, 1904.

- Breakfast. Boiled hominy 150 grams, milk 125 grams, sugar 25 grams, bread 50 grams, stewed prunes 150 grams, coffee 1 cup.
- Dinner. Baked spaghetti with a little grated cheese 200 grams, mashed potato 200 grams, bread 75 grams, boiled tomato 150 grams, pickles 20 grams, fruit pie, 130 grams, coffee 1 cup.
- Supper. Biscuit 175 grams, fried bacon 20 grams, French fried potatoes 150 grams, butter 20 grams, tea 1 cup.

#### Tuesday, February 9, 1904.

- Breakfast. Fried hominy 125 grams, syrup 50 grams, baked potato 150 grams, butter 10 grams, coffee 1 cup.
- Dinner. Boiled sweet potato 150 grams, butter 10 grams, bread 75 grams, thick pea soup 200 grams, boiled onions 150 grams, coffee 1 cup, apple pie 150 grams.
- Supper. Celery-lettuce-apple salad 150 grams, crackers 32 grams, cheese (American) 20 grams, Saratoga chips 75 grams, tea 1 cup.

#### Wednesday, February 10, 1904.

- Breakfast. Wheat griddle cakes 200 grams, syrup 50 grams, butter 10 grams, coffee 1 cup, banana 90 grams.
- Dinner. Boiled salt mackerel 25 grams, boiled potatoes 200 grams, boiled turnips 200 grams, bread 75 grams, coffee 1 cup, apple 140 grams.
- Supper. Chocolate cake 150 grams, cranberry sauce 100 grams, chopped fresh cabbage with salt, pepper, and vinegar 100 grams, bread 75 grams, butter 20 grams, tea 1 cup.

#### Thursday, February 11, 1904.

- Breakfast. Breakfast food 40 grams, milk 125 grams, sugar 25 grams, baked potato 150 grams, butter 10 grams, coffee 1 cup, apple 130 grams.
- Dinner. Baked beans 70 grams, salt pork 30 grams, bread 75 grams, boiled cabbage 200 grams, boiled potato 150 grams, coffee 1 cup.
- Supper. Tapioca-peach pudding 250 grams, bread 75 grams, butter 20 grams, tea 1 cup, cranberry sauce 100 grams.

### Friday, February 12, 1904.

- Breakfast. Breakfast food 40 grams, milk 125 grams, banana 90 grams, French fried potatoes 200 grams, sugar 25 grams, coffee 1 cup.
- Dinner. Clam chowder with onions, potatoes, and tomatoes 200 grams, bread 75 grams, mashed potato 200 grams, boiled turnips 150 grams, pie 125 grams, coffee 1 cup.
- Supper. Apple fritters 200 grams, jam 75 grams, tea 1 cup, gingerbread 30 grams.

### Saturday, February 13, 1904.

- Breakfast. Boiled rice 150 grams, milk 125 grams, sugar 25 grams, coffee 1 cup, baked potato 100 grams, butter 10 grams.
- Dinner. Meat pie with meat, potatoes, and onions 200 grams, boiled cabbage 200 grams, boiled potatoes 200 grams, bread 50 grams, coffee 1 cup.
- Supper. Lettuce-apple-celery salad 150 grams, biscuit 150 grams, butter 20 grams, tea 1 cup, stewed prunes 125 grams.

#### Sunday, February 14, 1904.

- Breakfast. Apple 140 grams, fried rice 125 grams, syrup 50 grams, baked sweet potato 142 grams, butter 10 grams, coffee 1 cup.
- Dinner.—Tomato soup with potatoes and onions boiled together, thickened with corn starch 350 grams, bread 75 grams, canned string beans 100 grams, baked apple with sugar 140 grams, coffee 1 cup.
- Supper. Small fried sausage 50 grams, French fried potatoes 200 grams, bread 50 grams, butter 10 grams, stewed prunes 100 grams, tea 1 cup.

#### Monday, February 15, 1904.

- Breakfast. Wheat griddle cakes 200 grams, syrup 50 grams, Johnny cake 50 grams, butter 10 grams, coffee 1 cup.
- Dinner. Hamburg steak, with bread, fat, and onions 150 grams, boiled potato 200 grams, bread 75 grams, butter 10 grams, coffee 1 cup, pickles 25 grams.
- Supper. Boiled Lima beans thoroughly cooked 75 grams, mashed potato 160 grams, bread 75 grams, butter 10 grams, tea 1 cup, stewed peaches 125 grams.

#### Tuesday, February 16, 1904.

- Breakfast.—Brown bread 50 grams, baked potato 230 grams, butter 20 grams, coffee 1 cup, apple 140 grams.
- Dinner. Boiled macaroni 250 grams, stewed tomatoes 250 grams, French fried potatoes 160 grams, bread 75 grams, coffee 1 cup, pie 130 grams.
- Supper. Fried bacon 25 grams, potatoes stewed in cream 250 grams, rice pudding 200 grams, bread 50 grams, tea 1 cup.

### Wednesday, February 17, 1904.

- Breakfast. Fried rice 125 grams, syrup 50 grams, baked sweet potato 190 grams, butter 10 grams, bread 50 grams, coffee 1 cup.
- Dinner. Corned beef 40 grams, cabbage 200 grams, mashed potato 200 grams, bread 75 grams, coffee 1 cup, tapioca-peach pudding 200 grams.
- Supper. Lettuce-apple-celery salad 150 grams, bread 100 grams, butter 20 grams, cranberry sauce 125 grams, tea 1 cup, baked potato 100 grams.

### Thursday, February 18, 1904.

- Breakfast. Boiled hominy 175 grams, milk 125 grams, sugar 25 grams, coffee 1 cup, banana 110 grams.
- Dinner. Split-pea soup (thick) 200 grams, bread 100 grams, mashed potatoes 200 grams, boiled onions 200 grams, coffee 1 cup.
- Supper. Corned-beef hash (mostly potato) 125 grams, bread 50 grams, fried sweet potato 150 grams, butter 20 grams, tea 1 cup, jam 75 grams.

### Friday, February, 19, 1904.

- Breakfast. Fried hominy 125 grams, syrup 50 grams, baked potato 150 grams, butter 10 grams, apple 140 grams, coffee 1 cup.
- Dinner. Boiled salt mackerel 25 grams, boiled potatoes 250 grams, boiled turnips 200 grams, bread 75 grams, coffee 1 cup, apple pie 100 grams.
- Supper. Chocolate cake 150 grams, cranberry sauce 125 grams, chopped fresh cabbage with salt, pepper, and vinegar 100 grams, bread 75 grams, butter 20 grams, tea 1 cup.

#### Saturday, February 20, 1904.

- Breakfast. Breakfast food 40 grams, milk 125 grams, sugar 25 grams, French fried potatoes 200 grams, coffee 1 cup.
- Dinner. Barley broth with potatoes and onions 250 grams, wheat flour dumplings 150 grams, boiled carrots 150 grams, bread 75 grams, boiled sweet potato 180 grams, coffee 1 cup.
- Supper. Suet pudding 150 grams, baked apple with sugar 150 grams, crackers 25 grams, stewed prunes 150 grams, tea 1 cup.

#### Sunday, February 21, 1904.

- Breakfast. Boiled oatmeal (thin) 125 grams, milk 100 grams, sugar 25 grams, baked potato 150 grams, butter 10 grams, coffee 1 cup.
- Dinner. Roast beef 40 grams, mashed potato 250 grams, bread 75 grams, apple sauce 150 grams, stewed tomatoes 150 grams, coffee 1 cup.
- Supper. Bread pudding 200 grams, cranberry sauce 200 grams, potato chips 100 grams, tea 1 cup.

### Monday, February 22, 1904.

- Breakfast. Wheat griddle cakes 200 grams, syrup 50 grams, butter 10 grams, baked potato 180 grams, coffee 1 cup.
- Dinner. Boiled macaroni 200 grams, stewed tomatoes 250 grams, French fried potatoes 200 grams, bread 35 grams, coffee 1 cup, apple sauce 150 grams.
- Supper. Lettuce-apple-celery salad 150 grams, biscuit 150 grams, butter 20 grams, stewed prunes 125 grams, tea 1 cup.

### Tuesday, February 23, 1904.

- Breakfast. Boiled rice 175 grams, milk 100 grams, sugar 25 grams, coffee 1 cup, banana 90 grams.
- Dinner. Bean soup (thick) 200 grams, mashed potato 250 grams, bread 35 grams, butter 10 grams, boiled onions 150 grams, coffee 1 cup, apple pie 100 grams.
- Supper. French fried potatoes 100 grams, crackers 25 grams, butter 15 grams, tea 1 cup, stewed peaches 150 grams.

### Wednesday, February 24, 1904.

- Breakfast. Fried rice 125 grams, syrup 50 grams, baked potato 150 grams, butter 10 grams, bread 35 grams, coffee 1 cup.
- Dinner. Boiled codfish 50 grams, mashed potato 250 grams, boiled carrots 150 grams, bread 50 grams, coffee 1 cup.
- Supper. Bread pudding with raisins 250 grams, jam 75 grams, crackers 25 grams, butter 15 grams, tea 1 cup.

#### Thursday, February 25, 1904.

- Breakfast. Boiled hominy 175 grams, milk 100 grams, sugar 25 grams, coffee 1 cup, baked sweet potato 125 grams, butter 10 grams.
- Dinner. Hamburg steak with much bread, fat, and onions 150 grams, boiled potatoes 200 grams, butter 15 grams, bread 50 grams, coffee 1 cup, pickles 35 grams.
- Supper. Tapioca-apple pudding 250 grams, sponge cake 75 grams, cranberry sauce 100 grams, tea 1 cup.

#### Friday, February 26, 1904.

- Breakfast. Fried hominy 100 grams, syrup 50 grams, bread 50 grams, butter 10 grams, coffee 1 cup, apple 150 grams.
- Dinner. Oyster chowder with tomatoes, potatoes, and onions 200 grams, bread 75 grams, string beans 150 grams, coffee 1 cup.
- Supper. Suet pudding with plums 125 grams, apple sauce 150 grams, baked sweet potato 200 grams, crackers 25 grams, butter 10 grams, tea 1 cup.

### Saturday, February 27, 1904.

- Breakfast. Soft boiled oatmeal 125 grams, milk 100 grams, sugar 25 grams, coffee 1 cup, butter 10 grams, baked potato 140 grams.
- Dinner. Lyonnaise potato 175 grams, bacon 25 grams, boiled turnips 200 grams, coffee 1 cup, rice pudding 150 grams.
- Supper. Banana fritters 200 grams, stewed prunes 150 grams, crackers 25 grams, butter 10 grams, tea 1 cup.

### Sunday, February 28, 1904.

- Breakfast. Stewed peaches 150 grams, wheat griddle cakes 200 grams, syrup 50 grams, coffee 1 cup, baked potato 150 grams, butter 10 grams.
- Dinner. Barley broth with potatoes and onions 250 grams, wheat flour dumplings 150 grams, French fried potatoes 150 grams, bread 35 grams, coffee 1 cup, ice cream 100 grams.
- Supper. Saratoga chips 75 grams, fried sausage 40 grams, butter 15 grams, bread 50 grams, tea 1 cup.

#### Monday, February 29, 1904.

- Breakfast. Boiled rice 175 grams, milk 125 grams, sugar 25 grams, baked potato 150 grams, coffee 1 cup, 350 grams, butter 10 grams.
- Dinner. Baked spaghetti 250 grams, mashed potato 250 grams, bread 75 grams, boiled tomatoes 150 grams, apple pie 112 grams, coffee 1 cup, 350 grams.
- Supper. Biscuit 175 grams, fried bacon 20 grams, fried sweet potatoes 150 grams, butter 20 grams, tea 1 cup, 350 grams.

Total nitrogen, 10.466 grams.

Fuel value, 2670 calories.

#### Tuesday, March 1, 1904.

- Breakfast. Fried rice 150 grams, syrup 50 grams, baked potato 150 grams, butter 10 grams, coffee 1 cup, 350 grams.
- Dinner. Thick pea-soup 250 grams, boiled onions 150 grams, boiled sweet potato 150 grams, bread 75 grams, butter 20 grams, coffee 1 cup, 350 grams.
- Supper. Celery-lettuce-apple salad 120 grams, crackers 82 grams, American cheese 20 grams, Saratoga chips 79 grams, tea 1 cup, 350 grams, rice custard 100 grams.

Total nitrogen, 7.825 grams.

Fuel value, 2279 calories.

#### Wednesday, March 2, 1904.

- Breakfast. Wheat griddle cakes 200 grams, syrup 50 grams, butter 10 grams, coffee 1 cup, 350 grams, banana 75 grams.
- Dinner. Boiled salt mackerel 25 grams, boiled potatoes 250 grams, boiled turnips 150 grams, bread 75 grams, coffee 1 cup, 350 grams, apple sauce 150 grams.
- Supper. Chopped fresh cabbage with salt, pepper, and vinegar, 100 grams, bread 75 grams, butter 20 grams, chocolate cake 150 grams, cranberry sauce 100 grams, tea 1 cup, 350 grams.

Total nitrogen, 8.487 grams.

Fuel value, 2391 calories.

### Thursday, March 3, 1904.

- Breakfast. Boiled hominy 175 grams, milk 125 grams, sugar 25 grams, baked potato 150 grams, butter 10 grams, coffee 1 cup, 350 grams.
- Dinner. Hamburg steak with much bread, fat and onions 150 grams, boiled potato 250 grams, bread 75 grams, butter 10 grams, coffee 1 cup, 850 grams.
- Supper. Tapioca-peach pudding 250 grams, bread 75 grams, jam 75 grams, butter 20 grams, tea 1 cup, 350 grams.

Total nitrogen, 8.750 grams.

Fuel value, 2375 calories.

#### Friday, March 4, 1904.

- Breakfast. Fried hominy 150 grams, syrup 50 grams, baked potato 150 grams, coffee 1 cup, 350 grams, butter 10 grams.
- Dinner. Codfish-balls (1 part fish 4 parts potato) fried in pork fat 150 grams, stewed tomatoes 200 grams, stewed potatoes 150 grams, bread 75 grams, coffee 1 cnp, 350 grams, apple pie 180 grams.
- Supper. French fried potatoes 200 grams, fried sausage 50 grams, butter 10 grams, stewed prunes 125 grams, sponge cake 35 grams, bread 50 grams, tea 1 cup, 350 grams.

Total nitrogen, 10.427 grams.

Fuel value, 2374 calories.

#### Saturday, March 5, 1904.

- Breakfast. Boiled Indian-meal 200 grams, milk 125 grams, sugar 25 grams, coffee 1 cup, 350 grams, fried sweet potato 150 grams, butter 10 grams.
- Dinner. Tomato soup thick, with potatoes and onions boiled together 325 grams, bread 100 grams, scrambled egg 50 grams, mashed potato 150 grams, coffee 1 cup, 350 grams.
- Supper. Bread pudding with raisins 250 grams, stewed peaches 150 grams, bacon 20 grams, French fried potatoes 150 grams, bread 50 grams, butter 10 grams, tea 1 cup, 350 grams.

Total nitrogen, 10.483 grams.

Fuel value, 2802 calories.

#### Sunday, March 6, 1904.

- Breakfast. Fried Indian-meal 150 grams, syrup 50 grams, sliced banana 100 grams, baked potato 150 grams, butter 10 grams, coffee 1 cup, 350 grams.
- Dinner. Corned beef 50 grams, boiled cabbage 200 grams, mashed potato 250 grams, bread 75 grams, fried rice 100 grams, jam 75 grams, coffee 1 cup, 350 grams.
- Supper. Sponge cake 150 grams, apple sauce 150 grams, crackers 82 grams, butter 10 grams, sardine 14 grams, tea 1 cup, 350 grams.

Total Nitrogen, 10.265 grams.

Fuel value, 3173 calories.

#### Monday, March 7, 1904.

- Breakfast. Boiled rice 175 grams, milk 125 grams, sugar 25 grams, baked potato 150 grams, butter 10 grams, coffee 1 cup.
- Dinner. Baked spaghetti 300 grams, mashed potato 250 grams, bread 75 grams, boiled tomatoes 200 grams, apple pie 125 grams, coffee 1 cup.
- Supper. Biscuit 175 grams, fried bacon 80 grams, fried sweet potato 200 grams, butter 20 grams, tea 1 cup.

#### Tuesday, March 8, 1904.

- Breakfast.—Fried rice 150 grams, syrup 50 grams, baked potato 200 grams, butter 10 grams, coffee 1 cup.
- Dinner. Thick pea-soup 300 grams, boiled sweet potato 250 grams, boiled onions 150 grams, bread 75 grams, butter 20 grams, pickles 30 grams, coffee 1 cup.
- Supper. French fried potatoes 150 grams, fried bacon 20 grams, crackers 32 grams, apple sauce 200 grams, rice custard 100 grams, tea 1 cup.

#### Wednesday, March 9, 1904.

- Breakfast. Wheat griddle cakes 200 grams, syrup 50 grams, butter 10 grams, banana 90 grams, coffee 1 cup.
- Dinner. Boiled salt mackerel 25 grams, boiled potato 250 grams, boiled turnips 200 grams, bread 75 grams, apple sauce 200 grams, coffee 1 cup.
- Supper. Chopped fresh cabbage with salt, pepper and vinegar, 75 grams, bread 75 grams, butter 20 grams, chocolate cake 150 grams, cranberry sauce 100 grams, tea 1 cup.

#### Thursday, March 10, 1904.

- Breakfast.—Boiled hominy 175 grams, milk 125 grams, sugar 25 grams, baked potato 250 grams, butter 10 grams, coffee 1 cup.
- Dinner. Hamburg steak with much bread, fat, and onions 150 grams, boiled potato, 250 grams, bread 75 grams, butter 10 grams, coffee 1 cup, pickles 80 grams.
- Supper. Tapioca-peach pudding 250 grams, bread 75 grams, jam 100 grams, butter 20 grams, tea 1 cup.

#### Friday, March 11, 1904.

- Breakfast. Fried hominy 150 grams, syrup 50 grams, baked potato 250 grams, butter 10 grams, apple sauce 150 grams, coffee 1 cup.
- Dinner. Codfish-balls (1 part fish, 4 parts potato, fried in pork fat) 150 grams, stewed tomatoes 200 grams, stewed potatoes 250 grams, bread 75 grams, apple pie 130 grams, coffee 1 cup.
- Supper. French fried potatoes 200 grams, fried sausage 50 grams, bread 50 grams, butter 10 grams, stewed prunes 125 grams, sponge cake 35 grams, tea 1 cup.

#### Saturday, March 12, 1904.

- Breakfast. Boiled Indian-meal 200 grams, milk 125 grams, sugar 25 grams, coffee 1 cup, fried sweet potato 150 grams, butter 10 grams.
- Dinner. Tomato soup thick, with potatoes and onions 325 grams, bread 100 grams, fried egg 30 grams, mashed potato 250 grams, coffee 1 cup, pickles 30 grams.
- Supper. Fried bacon 20 grams, French fried potatoes 150 grams, bread 50 grams, butter 10 grams, bread pudding with raisins 250 grams, stewed peaches 200 grams, tea 1 cup.

#### Sunday, March 13, 1904.

- Breakfast. Fried Indian-meal 150 grams, syrup 50 grams, apple sauce 200 grams, baked potato 250 grams, butter 10 grams, coffee 1 cup.
- Dinner. Hamburg steak with much bread, fat, and onions 150 grams, boiled cabbage 200 grams, boiled potatoes 250 grams, bread 75 grams, butter 10 grams, coffee 1 cup.
- Supper. Suet pudding 150 grams, stewed peaches 200 grams, crackers 32 grams, butter 10 grams, sardine 30 grams, tea 1 cup.

#### Monday, March 14, 1904.

- Breakfast. Wheat griddle cakes 200 grams, syrup 50 grams, butter 10 grams, coffee 1 cup, banana 80 grams.
- Dinner. Baked spaghetti 300 grams, mashed potato 250 grams, bread 75 grams, stewed tomatoes 200 grams, coffee 1 cup, baked apple with sugar 150 grams.
- Supper. Biscuit 175 grams, fried bacon 20 grams, fried sweet potato 200 grams, butter 20 grams, tea 1 cup.

#### Tuesday, March 15, 1904.

- Breakfast.—Boiled rice 175 grams, milk 125 grams, sugar 25 grams, baked potato 200 grams, butter 10 grams, coffee 1 cup.
- Dinner. Thick pea-soup 300 grams, boiled potatoes 250 grams, boiled onions 150 grams, pickles 30 grams, bread 75 grams, butter 20 grams, coffee 1 cup, rice custard 100 grams.
- Supper. Fried bacon 20 grams, French fried potatoes 200 grams, bread 50 grams, apple sauce 200 grams, tea 1 cup.

#### Wednesday, March 16, 1904.

- Breakfast. Fried rice 150 grams, syrup 50 grams, baked potato 200 grams, butter 10 grams, coffee 1 cup.
- Dinner. Codfish-balls (1 part fish, 4 parts potato, fried in pork fat) 150 grams, stewed potatoes 250 grams, stewed tomatoes 200 grams, bread 75 grams, coffee 1 cup, apple pie 125 grams.
- Supper. Apple-lettuce-celery salad 100 grams, bread 100 grams, butter 20 grams, stewed peaches 200 grams, tea 1 cup, sponge cake 50 grams.

#### Thursday, March 17, 1904.

- Breakfast. Boiled hominy 175 grams, milk 125 grams, sugar 25 grams, baked potato 250 grams, butter 10 grams, coffee 1 cup.
- Dinner. Hamburg steak with much bread, fat, and onions 150 grams, mashed potato 250 grams, pickles 30 grams, bread 50 grams, butter 10 grams, boiled turnips 150 grams, coffee 1 cup.
- Supper. Tapioca-peach pudding 250 grams, bread 75 grams, jam 100 grams, butter 20 grams, tea 1 cup.

#### Friday, March 18, 1904.

- Breakfast. Fried hominy 150 grams, syrup 50 grams, baked potato 250 grams, butter 10 grams, apple sauce 150 grams, coffee 1 cup.
- Dinner.—Clam chowder with much potato, tomato and onions 250 grams, bread 75 grams, mashed potato 150 grams, boiled sweet potato 150 grams, coffee 1 cup, pickles 30 grams.
- Supper. Bread pudding with raisins 250 grams, stewed peaches 200 grams, fried bacon 20 grams, French fried potatoes 150 grams, crackers 24 grams, butter 10 grams, tea 1 cup.

#### Saturday, March 19, 1904.

- Breakfast. Boiled Indian-meal 200 grams, milk 125 grams, sugar 25 grams, fried sweet potato 150 grams, butter 10 grams, coffee 1 cup.
- Dinner. Tomato soup thick, with potatoes and onions 325 grams, bread 100 grams, mashed potato 250 grams, coffee 1 cup, pickles 30 grams.
- Supper. Fried egg 30 grams, baked potato 250 grams, butter 20 grams, biscuit 175 grams, tea 1 cup, stewed prunes 125 grams.

#### Sunday, March 20, 1904.

- Breakfast. Fried Indian-meal 150 grams, syrup 50 grams, baked potato 250 grams, butter 20 grams, coffee 1 cup, apple 150 grams.
- Dinner. Roast beef 50 grams, mashed potato 250 grams, boiled carrots 100 grams, bread 50 grams, apple pie 130 grams, coffee 1 cup, pickles 80 grams.
- Supper.—Apple fritters 200 grams, cranberry sauce 150 grams, gingerbread 30 grams, tea 1 cup.

#### Monday, March 21, 1904.

- Breakfast. Wheat griddle cakes 200 grams, syrup 50 grams, butter 10 grams, baked potato 200 grams, coffee 1 cup.
- Dinner. Baked spaghetti 250 grams, mashed potato 250 grams, stewed tomatoes 150 grams, bread 75 grams, apple pie 125 grams, coffee 1 cup.
- Supper. Biscuit 150 grams, fried bacon 20 grams, fried sweet potatoes 150 grams, butter 20 grams, tea 1 cup.

#### Tuesday, March 22, 1904.

- Breakfast. Boiled rice 175 grams, milk 125 grams, sugar 25 grams, baked potatoes 150 grams, coffee 1 cup, butter 10 grams.
- Dinner. Thick pea-soup 250 grams, boiled onions 150 grams, boiled sweet potato 150 grams, bread 75 grams, butter 20 grams, coffee 1 cup, pickles 25 grams.
- Supper. Banana with sugar 75 grams, crackers 32 grams, American cheese 20 grams, Saratoga chips 75 grams, tea 1 cup, rice custard 100 grams.

#### Wednesday, March 23, 1904.

- Breakfast. Fried rice 150 grams, syrup 50 grams, baked potato 150 grams, butter 10 grams, coffee 1 cup, apple 150 grams.
- Dinner. Boiled salt mackerel 25 grams, boiled potato 250 grams, boiled turnips 150 grams, pickle 20 grams, bread 75 grams, coffee 1 cup, apple sauce 100 grams.
- Supper. Chopped fresh cabbage with salt, pepper, and vinegar 100 grams, bread 75 grams, butter 20 grams, chocolate cake 100 grams, cranberry sauce 100 grams, tea 1 cup.

#### Thursday, March 24, 1904.

- Breakfast. Boiled hominy 175 grams, milk 125 grams, sugar 25 grams, baked potato 150 grams, butter 10 grams, coffee 1 cup.
- Dinner. Hamburg steak with much bread, fat, and onions 150 grams, boiled potatoes 250 grams, bread 75 grams, butter 10 grams, coffee 1 cup.
- Supper. Tapioca-peach pudding 250 grams, bread 75 grams, jam 75 grams, butter 20 grams, tea 1 cup.

#### Friday, March 25, 1904.

- Breakfast. Fried hominy 150 grams, syrup 50 grams, baked potato 150 grams, coffee 1 cup, apple 150 grams.
- Dinner. Codfish-balls (1 part fish, 4 parts potato, fried in pork fat) 150 grams, stewed tomatoes 200 grams, stewed potatoes 150 grams, bread 75 grams, coffee 1 cup, apple pie 130 grams.
- Supper. French fried potatoes 200 grams, fried bacon 20 grams, butter 10 grams, bread 50 grams, stewed prunes 125 grams, tea 1 cup.

#### Saturday, March 26, 1904.

- Breakfast. Boiled Indian-meal 175 grams, milk 125 grams, sugar 25 grams, butter 10 grams, fried sweet potato 150 grams, coffee 1 cup.
- Dinner. Tomato soup with potato and onions boiled together 325 grams, bread 100 grams, fried egg 40 grams, mashed potato 150 grams, coffee 1 cup.
- Supper. Saratoga chips 50 grams, fried bacon 20 grams, bread 50 grams, butter 10 grams, bread pudding with raisins 250 grams, stewed peaches 150 grams, tea 1 cup.

#### Sunday, March 27, 1904.

- Breakfast. Fried Indian-meal 150 grams, syrup 50 grams, baked potato 150 grams, coffee 1 cup, butter 10 grams, banana 90 grams.
- Dinner. Corned beef 40 grams, boiled cabbage 200 grams, mashed potato 250 grams, bread 75 grams, coffee 1 cup, squash pie 130 grams.
- Supper. Suet pudding 150 grams, apple sauce 150 grams, crackers 32 grams, butter 10 grams, chopped cabbage with salt, pepper, and vinegar 60 grams, tea 1 cup.

#### Monday, March 28, 1904.

- Breakfast. Fried rice 150 grams, syrup 75 grams, baked potato 250 grams, butter 20 grams, coffee, 1 cup, 850 grams.
- Dinner. Thick pea-soup 200 grams, boiled onions 100 grams, boiled sweet potato 250 grams, bread 50 grams, mashed potato 200 grams, butter 20 grams, coffee 1 cup, 350 grams.
- Supper. Biscuit 125 grams, fried bacon 20 grams, French fried potatoes 200 grams, butter 25 grams, banana 150 grams, tea 1 cup, 850 grams.

Total nitrogen, 9.027 grams.

Fuel value, 2935 calories.

#### Tuesday, March 29, 1904.

- Breakfast. Boiled hominy 175 grams, milk 75 grams, sugar 25 grams, baked potato 250 grams, butter 20 grams, coffee 1 cup, 350 grams.
- Dinner.—Hamburg steak with much bread, fat, and onions 125 grams, boiled potatoes 300 grams, butter 10 grams, bread 35 grams, boiled carrots 125 grams, coffee 1 cup, 350 grams.
- Supper.—Tapioca-peach pudding 300 grams, bread 35 grams, Saratoga chips 75 grams, butter 20 grams, jam 75 grams, tea 1 cup, 350 grams.

Total nitrogen, 8.972 grams.

Fuel value, 2840 calories.

#### Wednesday, March 30, 1904.

- Breakfast. Fried hominy 150 grams, syrup 75 grams, butter 10 grams, banana 250 grams, coffee 1 cup, 850 grams.
- Dinner. Codfish-balls (1 part fish, 5 parts potato, fried in pork fat) 125 grams, bread 35 grams, mashed potatoes 250 grams, stewed tomatoes 200 grams, apple sauce 200 grams, coffee 1 cup, 350 grams.
- Supper. Chopped fresh cabbage with salt, pepper, and vinegar, 75 grams, bread 50 grams, butter 20 grams, fried sweet potato 250 grams, cranberry sauce 200 grams, sponge cake 50 grams, tea 1 cup, 350 grams.

Total nitrogen, 9.856 grams.

Fuel value, 2657 calories.

#### Thursday, March 31, 1904.

- Breakfast.—Fried Indian-meal 100 grams, syrup 75 grams, baked potato 250 grams, butter 20 grams, coffee, 1 cup, 850 grams.
- Dinner. Tomato soup thick, with potato and onions boiled together 800 grams, mashed potato 200 grams, scrambled egg 50 grams, bread 50 grams, butter 10 grams, coffee 1 cup, 850 grams.
- Supper. Fried bacon 20 grams, boiled potato 200 grams, butter 10 grams, bread pudding 150 grams, banana 200 grams, tea 1 cup, 350 grams.

Total nitrogen, 8.420 grams.

Fuel value, 2466 calories.

#### Friday, April 1, 1904.

- Breakfast. Fried hominy 150 grams, syrup 75 grams, baked potato 200 grams, butter 20 grams, coffee 1 cup, 350 grams.
- Dinner. Baked spaghetti 250 grams, mashed potato 250 grams, boiled turnips 150 grams, bread 85 grams, butter 10 grams, apple sauce 200 grams, coffee 1 cup, 850 grams.
- Supper. Fried sweet potato 200 grams, butter 20 grams, jam 100 grams, fried bacon 25 grams, bread 35 grams, apple-tapioca pudding 300 grams, tea 1 cup, 350 grams.

Total nitrogen, 7.342 grams.

Fuel value, 8248 calories.

#### Saturday, April 2, 1904.

- Breakfast. Griddle cakes 200 grams, syrup 50 grams, baked potato 200 grams, butter 10 grams, coffee 1 cup, apple 140 grams.
- Dinner. Boiled mackerel 25 grams, boiled potato 250 grams, boiled turnips 150 grams, pickles 35 grams, bread 50 grams, pie 130 grams, butter 10 grams, coffee 1 cup.
- Supper. Suet pudding 125 grams, cranberry sauce 150 grams, baked sweet potato 200 grams, crackers 25 grams, butter 10 grams, tea 1 cup.

#### Sunday April 3, 1904.

- Breakfast. Banana 90 grams, boiled rice 175 grams, milk 125 grams, sugar 25 grams, baked potato 200 grams, butter 10 grams, coffee 1 cup.
- Dinner. Corned beef 40 grams, boiled cabbage 200 grams, boiled potatoes 200 grams, bread 75 grams, butter 10 grams, coffee 1 cup, stewed prunes 150 grams.
- Supper. Apple-tapioca pudding 250 grams, jam 75 grams, crackers 25 grams, butter 10 grams, pickles 25 grams, tea 1 cup.

#### Monday, April 4, 1904.

- Breakfast. Fried rice 150 grams, syrup 50 grams, apple 130 grams, baked sweet potato 200 grams, butter 10 grams, coffee 1 cup.
- Dinner. Lyonnaise potatoes 175 grams, bacon 25 grams, boiled turnips 200 grams, pie 130 grams, coffee 1 cup, bread 75 grams.
- Supper. Biscuit 175 grams, butter 25 grams, egg 40 grams, stewed apples 150 grams, tea 1 cup.

## III. EXPERIMENTS WITH UNIVERSITY STUDENTS, TRAINED IN ATHLETICS.

Men in training for athletic events deem it necessary to consume large amounts of proteid food. Great muscular activity, it is true, calls for the expenditure of corresponding amounts of energy, but it is by no means clear that the energy so liberated comes from the breaking down of proteid material. Indeed, there is more reason for believing that the energy of muscular contraction comes primarily from the oxidation of non-nitrogenous matter. Nevertheless, custom and long experience sanction a high proteid diet, composed largely of meat or of other foodstuffs rich in nitrogen, for the development of that vigor and strength that go to make the accomplished athlete. For the development of new muscle tissue, to make two muscle fibres where before only one existed, to increase the number of available fibres, thereby adding to the bulk of the active tissue, a certain amount of proteid food is absolutely necessary, just as it is for all active tissues and organs of the body. But that this fact constitutes a satisfactory reason for the daily use of such quantities of proteid food as usually enter into the diet of the average athlete is very questionable.

As an illustration of the character of the diet frequently made use of by men doing prolonged muscular work, I may quote a few figures from an article by Professor W. O. Atwater \* and H. C. Sherman on "The effect of severe and prolonged muscular work upon Food consumption, Digestion, and Metabolism," in which are recorded observations made upon several professional bicycle riders during a six-day race. On one day subject "M" rode 334.1 miles, consuming for food on

<sup>\*</sup> U. S. Department of Agriculture, Office of Experiment Stations, Bulletin 98.

See also "Investigations on the Nutrition of Man in the United States." By C. F. Langworthy and R. D. Milner. U. S. Department of Agriculture, Office of Experiment Stations. Washington, 1904, p. 14.

that day meat extract, 311 grams; milk, 4937 grams; bread, 35 grams; boiled cereals, 877 grams; pastry, 142 grams; sugar, 53 grams; fruit, 2003 grams. His urine for the day contained 46.2 grams of nitrogen, corresponding to the metabolism of 288.7 grams of proteid matter. Subject "A" on one day rode 352.7 miles, taking as food, meat, 149 grams; meat extract, 24 grams; broth, 283 grams; eggs, 369 grams; butter, 78 grams; milk, 142 grams; malted milk, 78 grams; jelly, 213 grams; soup, 191 grams; bread, 361 grams; boiled cereal, 532 grams; sugar, about 400 grams; fruit, 933 grams; cocoa wine, 198 grams. His day's urine contained 39.0 grams of nitrogen, corresponding to the metabolism of 243.7 grams of proteid material.


Obviously, if such high proteid metabolism as these figures imply is a necessary concomitant of vigorous or excessive muscular activity, then a rich proteid diet is needed to make good the loss of nitrogen to the body; but we are more inclined to believe that the large quantity of nitrogen excreted was the result chiefly of the high proteid ration, and only in small measure connected with the work done. With a sufficiency of non-nitrogenous food, the energy of muscular contraction does not come in any large degree from the breaking down of proteid matter, and there would seem to be no adequate reason for assuming a necessity for such rich and excessive proteid diet as athletes in training ordinarily adopt. Yet the contrary view is almost universally held and followed. As a prominent trainer said to the writer not long ago, "if the men are not fed on a rich meat diet and plenty of it, they will grow soft and lose their strength."

With a view to testing some of these points and thereby broaden the scope of the investigation and enhance the value of the study, a group of eight students in the University, all trained athletes, was secured. These men volunteered to aid in the study, and at considerable self-sacrifice gave intelligent and hearty co-operation in all ways possible. The men were under observation from January 15, 1904, to the middle of June, 1904, a period of five months. From January 15 the



STAPLETON

Photograph taken in the middle of the experiment, in April.



urine was collected daily, and for a period of ten days the regular diet of the men was adhered to without any deviation whatever, with a view to ascertaining the extent of the proteid metabolism characteristic of each individual. strictions whatever in diet were suggested, but the ordinary food that the men were accustomed to eat while in training was taken. After this period the men were instructed to diminish somewhat the intake of proteid food, and in following out this plan most of the men diminished quite appreciably the quantity of food consumed at breakfast time, and in some cases stopped taking food of any kind at breakfast other than a cereal, with possibly coffee. No specific diet was imposed, but the men, being willing collaborators in the experiment, gradually cut down the intake of proteid food, diminishing likewise in considerable measure the total volume of food for the twenty-four hours.

The following brief description of the men, taken from Dr. Anderson's report of their physical condition, to be referred to later, will suffice to make clear the characteristics (as athletes) of these subjects of study:

- Mr. G. W. Anderson is a foot-ball, base-ball, and basket-ball player, as well as a crew man (not Varsity). Well built and an all round athlete. 26.5 years of age.
- Mr. W. L. Anderson, a "Y" athlete (hurdler), the captain of the Yale Gymnastic Team, University Gymnastic Champion, and American Collegiate Gymnastic Champion. 21.5 years of age.
- Mr. H. S. Bellis, a member of the Y. G. A., a gymnast and acrobat and in constant training. 26 years of age.
- Dr. W. H. Callahan, Medical Assistant at the Gymnasium, in daily practice in the gymnasium; bowling, hand-ball, and running. 27 years of age.
- Mr. M. Donahue, a very muscular and versatile athlete, a foot-ball player and a Varsity basket-ball player. 25 years of age.
- Mr. C. S. Jacobus, a "Y" athlete, a noted long-distance man, and one of the best University runners, 22.3 years of age.

Mr. H. R. Schenker, an active member of the Y. G. A., a point winner and intercollegiate competitor in gymnastics. 22 years of age.

Mr. John Stapleton, a wrestler and gymnast. A professional, a man of large body and great strength. 24 years of age.

The results of the daily study of the twenty-four hours' urine of each man for the entire period of five months are to be found in the accompanying tables. By a careful inspection of these data many interesting facts are brought to light. First it is to be observed, in harmony with what has already been stated regarding athletes, that the men, as a rule, were accustomed to the taking of large amounts of proteid food daily. Thus, during the preliminary period of ten days, Dr. Callahan excreted through the kidneys 22.79 grams of nitrogen as the daily average, corresponding to the metabolism of 142 grams of proteid matter per day, while on one day the nitrogen excretion reached the high figure of 31.99 grams, corresponding practically to the metabolism of 200 grams of proteid material.

G. W. Anderson, during the preliminary period of ten days—on his ordinary diet—showed an average daily excretion of 17.18 grams of nitrogen. W. L. Anderson in this same period excreted on an average 18.22 grams of nitrogen per day, while on one day the excretion reached 23.42 grams of nitrogen. Bellis showed an average daily excretion of 17.64 grams of nitrogen during this same fore period, while Stapleton excreted nitrogen at an average rate of 19.7 grams per day, thus indicating an average daily metabolism of 123 grams of proteid matter on his ordinary diet. With such data before us it is quite clear that we have here, as expected, a group of sturdy men accustomed to the taking of large amounts of proteid food daily; men who clearly believed that their strength and bodily vigor depended in large measure upon the ingestion and utilization of these quantities of proteid food.

After the termination of this preliminary period, the men began to diminish the amount of albuminons food. In doing

this they were at first given perfect freedom, each man following his own preferences, and making such alterations in his diet as he saw fit. The men were instructed as to the chemical composition of the various foodstuffs, so that they had a general idea of what foods were rich in nitrogen and could act accordingly. Some of the men cut down their intake of proteid food very rapidly, others made more gradual reduction. Some soon eliminated breakfast almost entirely. Others reduced the amount of food at each meal. Some of the men eliminated meat almost completely from their diet, and increased largely the intake of vegetable food. They were all advised, however, not to make too sudden a change in their diet, but to adopt a gradual reduction in the daily food as being less liable to disturb their physiological equilibrium. Naturally, the men knew exactly how much of a reduction in proteid food they were accomplishing each week, from the results of the chemical analysis of the urine. Further, the record of bodyweight furnished evidence of how far the changes in diet were affecting body equilibrium, while their own feelings and ability to do their daily work constituted a check upon a too radical change in living. Careful perusal of the tables will show how far body-weight was changed, and to how great an extent proteid metabolism was reduced during the five months of the experiment.

ANDERSON, G. W.

	Rode			Urine.	<del></del>	
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> .
1904	kilos	c.c.		grams	gram	grams
Jan. 16	75.0	1800	1022	18.02	0.470	
17		1300	1027	18.88	1.860	
18		1600	1025	17.95	1.008	
19		1085	1080	16.27	1.139	
20		890	1031	17.19	1.008	
21		1080	1030	17.69	0.940	
22	75.0	1340	1027	16.32	0.814	
23	75.0	1210	1028	17.42	1.020	• • •
24	75.0	990	1081	14.85	0.850	• • •
25	75.0	1020	1028	]]		
26	• • •	1085	1029	11		
27	• • •	865	1029	11		
28	• • •	800	1029	14.19	0.768	2.70
29	• • •	690	1026	daily	daily	daily
80	• • •	765	1026	average	average	average
31 Feb. 1	75.0	800	1024 1022	K		
reb. 1 2	75.0	970 1070	1022	11		
8	• • •	740	1024	11		
4	• • • •	870	1029	1105	0.633	1 77
5	• • •	2660	1010	11.85	0.055	1.77
8	• • •	860	1025	11		
7		865	1025	H		
8	75.0	890	1027	K		
9	75.0	780	1028	11		
10	75.0	1025	1025	11		
11	75.0	1240	1024	10.90	0.820	1.76
12	75.0	815	1027			1.10
13	75.0	940	1027			
14	75.0	1800	1017			
15	• • •	945	1025	lí		
16	73.5	1090	1029	[]		
17	78.0	720	1031	[]		
18	78.0	1000	1028	10.88	0.792	
19	73.0	1000	1028	11		
20		1800	1024	11		
21		1110	1025	IJ		
22		780	1026	h		
28	78.0	730	1030	10.00	0.699	1.72
24	• • •	730	1031	IJ		

ANDERSON, G. W.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P2O2-
1904	kilos	0.0.		grams	gram	grams
Feb. 25		910	1028	h		
26		780	1030	10.00	0.699	1.72
27	78.0	1 <b>4</b> 70	1025	daily	daily	daily
28		1010	1021	average	average	average
29	72.0	910	1029	וו		
Mar. 1		775	1031	[[		
2	• • •	1555	1018	11.70	0.740	104
8	• • •	885	1028	110	0.749	1.84
4		1060	1029			
6	• • •	910	1029	IJ		
7		770	1030	1)		
8	72.0	850	1029			
9	• • •	780	1030	! [		
10		740	1029	11.11	0.706	
11	72.0	770	1080	H		
12	• • •	920	1028	[[		
13	• • •	1265	1018	ין		
14	• • •	745	1027	[]		
15	71.0	645	1028	11		
16	71.8	700	1029			
17 18	• • •	660 640	1028 1029	7.84	0.630	• • •
19	72.0	885	1029	!		
20		885	1023	l i		
20 21	• • •	1180	1027	R I		
21 22	• • •	660	1022	[ [		
28	71.0	720	1029			
24	11.0	840	1020	7.41	0.628	
25		870	1025	1.41	0.628	• • • •
26		770	1029			
27		800	1031			
28		800	1028	K I		
29		810	1030			
80	71.0	820	1031			
31	• • •	780	1082	9.66	0.597	
Apr. 1		680	1084	5.55	0.001	•••
2		730	1031	11		
3		625	1033			
4		700	1021	l'		
5		740	1031	9.75	0.687	1.68

# 934 PHYSIOLOGICAL ECONOMY IN NUTRITION ANDERSON, G. W.

	Pode			Urine.		
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>8</sub> .
1904	kilos.	0.0.		grams	gram	grams
Apr. 6		695	1033	1)		
7		1220	1020	11		
8		670	1084	9.75	0.687	1.68
9		590	103 <b>4</b>	daily av.	daily av.	daily av.
10		510	1080	lí		
11	71.0	560	1083	9.23	}	
12		640	1030	l)	ŀ	
13		580	1030	8.11		
14		1195	1019	10.32	0.671	• • •
15		1005	1024	9.48		
16		1140	1021	10.60		
17		1050	1023	9.58	Į	
18	71.0	870	1026	9.03		
19		1110	1025	11.12	ļ ļ	
20		1170	1023	9.80		
21		810	1080	7.15	<b>0.580</b>	• • •
22		870	1029	9.55	1	
23	• • • •	660	1029	8.84		
24	70.0	765	1029	8.49	Į į	
25		640	1028	8.06	]	
26	• • • •	950	1026	9.80	[	
27		1120	1028	9.34		
28		1150	1021	9.04	0.522	• • •
29		860	1027	8.98		
80	71.0	870	1028	9.71	i i	
May 1		930	1025	10.16	ĺį –	
2	70.0	925	1021	8.88		
8	70.0	935	1023	9.20		
4		1140	1024	9.10		
5		990	1026	8.32	0.451	• • •
6		960	1026	8.06		
7	• • • •	585	1031	7.55		
8		620	1031	7.48	lí l	
9		660	1029	8.20		
10		740	1024	8.66		
11	70.0	1000	1023	8.10	0.500	
12	70.0	725	1025	8.09	0.590	• • •
13	71.0	650	1029	7.83		
14		780	1031	9.73		
15		670	1028	7.40	J :	
15	• • •	670	1028	7.40	J	

ANDERSON, G. W.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> .
1904	kilos	0.0.		grams	gram	grams
May 16		730	1030	8.50		
17		1025	1025	8.98		
18		930	1028	8.87	1	
19	71.1	865	1025	8.36	:	
20	71.0	1030	1027	9.95		
21	70.2	735	1030	8.51	0.523	
22	70.0	570	1029	6.50	daily	
23		625	1026	6.90	average	
24	70.5	870	1027	8.45		
25		665	1026	10.25	1	
26	70.0	640	1028	6.84	1	
27	70.5	780	1024	7.58	0.586	
28	70.0	660	1032	7.58	0.000	• • •
29		650	1027	6.91		
30	70.0	700	1028	6.97	[]	
June 1		940	1026	8.40		
2		726	1028	7.49		• • •
8	70.9	980	1020	8.65		
4		946	1026	8.80		
5		582	1029	6.32		
6	70.5	718	1027	7.67		• • •
7		670	1028	8.40		• • •
8		1160	1021	10.44		• • •
9		1010	1025	9.09		• • •
10	70.2	1510	1020	11.88		
11		725	1029	9.70		•••
12		1020	1028	10.10		
18		920	1024	10.82		• • •
14	70.0	740	1080	10.58		• • •
15	70.9	1425	1023	18.84	• • •	• • •
Feb. 1			1097		0.683	1.75
Daily aver Feb. 1 to	Apr. 12			9.94		•••
Daily aver Apr. 13 to	age from June 15	•••	•••	8.81	•••	• • •

ANDERSON, W. L.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric <b>∆</b> cid.	P3O8.
1904.	kilos	0.0.		grams	gram	grams
<b>Ja</b> n. 15	68.1	1240	1026	16.44	0.733	
16	63.1	1720	1020	18.68	0.755	
17	63.1	1470	1025	18.85	1.528	
18	61.3	1940	1018	19.09	0.866	
19	68.1	1850	1020	23.42	0.784	
20	63.1	600	1026	լ 16.85		
21	63.1	1720	1021	daily av.	• • •	• • •
22	62.7	1470	1017	18.41	0.584	
28		1210	1026	19.58	0.858	
24	63.0	1180	1025		0.598	
25	62.9	1060	1022	lı l		
26	63.1	1025	1024	11		
27	63.1	1560	1014	11 1		
28	63.6	890	1024	12.47	0.497	1.95
29	63.6	1850	1019	11	daily	daily
80	63.6	1070	1020	11	average	average
81	63.8	1410	1018	J J		_
Feb. 1	63.8	1425	1019	lı l		
2	62.8	940	1028			
3	62.0	865	1019			
4	62.2	1280	1020	10.65	0.498	1.99
5	62.2	1120	1017	11	•	
6	62.3	1390	1020	11		
7	62.3	1050	1019	Į) į		
8	62.8	1150	1021	h		
9	62.3	1430	1018	11		
10	60.0	910	1021			
11	60.0	1095	1018	10.82	0.505	2.07
12	60.0	865	1028	11		
18	62.1	865	1014	11		
14	62.1	1580	1015	D		
15	62.2	1270	1025	h		
16	61.2	690	1027	11		
17		1070	1020	11		
18	61.9	990	1025	12.58	0.507	
19		1480	1019	11		
20	60.9	1450	1019	11		
21	62.0	2582	1011	þ		
22	62.0	1811	1026	11.70	0.846	2.40
23	62.2	1460	1018	J 11.70	U.OEU	4.70

ANDERSON, W. L.

	Po-			Urine.		
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>3</sub> ,
1904	kilos	c.c.		grams	gram	grams
Feb. 24	60.0	1085	1022	11		
25	60.0	1215	1020	11		
26	62.2	835	1020	11.70	0.846	2.40
27	62.2	1590	1021	daily	daily	daily
28	62.2	1790	1012	average	average	average
29	62.3	820	1025	li -	_	
Mar. 1	61.1	770	1029	H		
2	62.2	670	1029	11		
8	62.2	1295	1023	} 12.91	0.669	2.32
4	• • •	1530	1020			
5	62.2	1260	1017			
6	62.3	1840	1024	IJ		
7	62.3	920	1025	)		
8	62.8	1200	1018	11		
<b>,</b> 9		880	1024	9.21	0.475	
11		1670	1019	8.21	0.410	• • • •
12	62.0	1865	1022			
18		1640	1030	}		
14	62.2	790	1026	1)		
15	62.2	715	1023	11		
16	62.2	1850	1016	11		
17	62.2	1140	1012	9.66	0.475	
18	61. <b>4</b>	980	1021			
19	62.9	1050	1024			
20	• • •	1012	1023	<b>[</b> ]		
21	62.0	910	1026	1)		
22	• • •	1460	1021	11		
23	61.9	1810	1020	11		
24	61.2	1050	1022	12.25	0.718	• • •
25	62.0	860	1031			
. 26	61.0	975	1028			
27	• • •	1080	1024	ļ,		
28	•••	1120	1022			
29	• • •	1690	1020			
30		1580	1017	11.19	0.551	
Apr. 1	62.3	1325	1018			
2	62.9	1075	1018			
8	62.9	1500	1015	K		
4	68.1	800	1025	10.21	0.625	1.92
5	62.9	1000	1022	1		

ANDERSON, W. L.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>2</sub> .
1904	kilos	c.c.		grams	gram	grams
Apr. 7	62.3	750	1027	la l		
8	62.3	600	1026	10.21	0.625	1.92
9	62.3	500	1028	daily av.	daily av.	daily av.
10	62.8	830	1028	اربت وست را		
11	61.3	600	1024	7.47	1	
12	62.2	950	1023	17		
13		800	1028	6.77		
14		1090	1018	8.88	0.454	
15		1250	1025	6.58	1	
16		1480	1014	7.81		
17	61.3	1130	1024	10.98	J	
18		590	1021	5.56	1	
19		840	1024	9.63		
20		980	1024	12.24		
21	• • •	1220	1022	10.69	0.431	• • •
22		1190	1019	9.50		
23		1390	1020	10.48		
24	62.2	1160	1017	7.52	J	
25		1010	1014	5.94	]	
26		1200	1019	10.14	l	
27		1660	1015	9.66		
28	61.8	1085	1021	8.59	0.459	• • •
29	61.8	1020	1022	10.88		
80		1180	1022	10.28	1	
May 1	• • • •	1695	1016	9.55	,	
2	• • • •	1530	1018	6.98	)	
8	• • • •	1280	1023	9.67	1	
4	• • •	1200	1020	8.50	0.237	
5		1000	1024	9.80	0.287	• • •
6		1890	1022	11.08		
7	• • • •	1025	1017	6.46	i	
8		1520	1017	8.66	,	
9	62.8	970	1017	6.40		
10	61.5	1140	1000	10.00		
11 12	600	1140	1022	10.38	0.451	
12	60.9	745	1017	6.39	0.401	• • •
18		850 1140	1022	8.31 10.12		
15		1865	1022	1		
16		1120	1017 1017	8.76 6.58	,	:
				0.00		

ANDERSON, W. L.

Date.	Body-			Urine.		
	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>8</sub> .
1904	kilos	0.0.		grams	gram	grams
May 17		1940	1014	10.83		
18		1370	1021	10.03	lı l	
19	61.8	1120	1022	10.15	11 1	
20		1170	1018	8.64	[ ]	
21		1530	1015	8.46	0.439	
22		890	1024	7.96	daily	
28		1200	. 1017	7.49	average	
24	61.1	770	1024	6.60		
25		690	1024	7.04	l) l	
26	60.2	745	1025	8.09		
27		970	1028	10.40	11	
28		1160	1022	10.87	0.448	
29					!!	
30		920	1025	8.94		
31	'				[]	•
June 1		1400	1012	5.54		
2	59.5	1070	1020	8.48		
8		1190	1018	8.78		
4	59.7	1020	1022	9.00		
5		870	1023	8.67		
6		870	1025	8.72		
7		1350	1017	9.31		
8	60.4	1880	1017	11.55		• • • •
9	60.4	1520	1017	9.58		
10		1590	1013	7.35		
11	60.4	1425	1017	7.74		
12		870	1021	7.67		
18		1410	1017	10.32	· · · · ·	• • •
14		1180	1022	10.84		• • •
15	61,0	875	1019	5.67		• • •
Feb. 1			1020	• • •	0.516	8.14
Daily aver	April 12		•••	10.78	•••	•••
15			• • •	10.05		

BELLIS.

	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P <sub>5</sub> O <sub>2</sub> .
1904	kilos	c.c.		grams	grams	grams
Jan. 15	81.8	1180	1026	18.24	0.778	
16	82.2	1915	1019	16.09	0.862	
17	82.2	1740	1024	16.18	0.881	• • •
18	82.2	2080	1021	20.10	0.868	• • •
19	82.2	1700	1026	17.34	0.771	
20	82.2	1800	1022	17.82	0.650	
21	82.2	1565	1027	26.18	1.020	
22	82.2	1435	1026	22.64	1.010	• • •
23	82.2	1285	1017	10.49	0.467	
24	'	1070	1026	16.37	0.786	
25	80.7	590	1029	l)		
26	81. <b>4</b>	1300	1020			
27	82.1	1020	1026	11		
28	81.2	1190	1027	<b>13.85</b>	0.693	2.82
29	81.2	1440	1018	daily	daily	daily
80	81.8	1450	1023	average	average	average
81	81.2	1000	1026	ן ו		
Feb. 1	81.2	1230	1025	h		
2	81.2	1145	1024	11		
8	81.2	1000	1030			
4	81.2	910	1080	12.98	0.674	2.29
5	81.2	1860	1023			
6	81.2	1160	1022			
7	81.2	1210	1027	ען		
8	81.2	1375	1016	h		
9	81.2	920	1031			
10	81.2	1775	1016			
11	81.2	1130	1026	<b>13.08</b>	0.659	2.44
12	81.2	1075	1026			
13	81.2	1405	1021	]]		
14	81.2	1055	1024			
15	81.0	1470	1025	h		
· 16	81.1	920	1029			
17	81.1	1530	1019			
18	81.1	1880	1023	14.15	0.750	• • •
19	81.1	1680	1020			
20	81.1	950	1029	[[		
21	81.1	2050	1018	ען		
22	81.1	1450	1030	14.79	0.850	2.60
23	81.1	1540	1027	)	0.500	

BELLIS.

<b>D</b>	Body-			Urine.		
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P2O2-
1904	kilos	c.c.		grams	gram	grams
Feb. 24	79.4	800	1030	h -		Ŭ
25	80.0	785	1033	] <b>]</b>		
26	80.4	1420	1018	14.79	0.850	2.60
27	80.1	1250	1026	daily av.	daily av.	daily av.
28	80.1	1070	1025	J		
29	80.1	1270	1024	h I		
Mar. 1	80.0	1150	1024			
2	80.0	1580	1018	11		
. 8	80.0	1820	1018	10.77	0.550	2.84
4	80.0	1685	1018			
5	80.0	1220	1026			
6	80.1	1550	1015			
7	80.0	1170	1020	li l		
8	80.0	1120	1017	11		
9	80.0	1220	1020			
10	80.0	1210	1020	} 10.29	0.628	
11	80.0	1650	1019			1
12	80.0	1855	1026			
18	80.0	1040	1025			ŀ
14	80.0	670	1027	1)		Ì
15	80.0	1380	1017			
16	80.0	915	1024			
17	80.0	1600	1016	10.05	0.622	
18	80.0	1510	1018	1		
19	80.0	1490	1018	[]		
20	79.9	920	1026	IJ		
21	79.8	710	1029	l)		1
22	79.8	800	1026			
23	79.8	800	1030			
24	79.8	1100	1025	10.15	0.728	
25	79.4	1060	1017			
26	79.4	1580	1020			
27	79.2	1040	1028	[]		
28	79.3	1860	1019	]}		
29	79.2	1335	1017			
80		1030	1017			
81		905	1019	8.52	• • •	•••
Apr. 1		870	• • •			
2		810	1029			
8	78.6	785	1028	ו		
	<u> </u>	L				1

# 342 PHYSIOLOGICAL ECONOMY IN NUTRITION BELLIS.

	Body-			Urine.		
Date.	Body- weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P3O8.
1904	kilos	c.c.		grams	gram	grams
Apr. 4		720	1029	h		
5		730	1019	ł 1		
6		790	1022	11		
7		880	1026	7.27	0.665	1.62
8		790	1025	daily	daily	daily
9		860	1027	average	average	average
10		890	1025	<b>[</b> ]		
11		890	1025	8.18	1	
12		590	1025	IJ		
13	77.2	960	1025	7.95		
14		930	1019	6.59	<b>0.496</b>	• • •
15		1090	1018	6.82		
16		1250	1022	10.35		
17		650	1025	5.97	)	
18		970	1020	7.92	)	
19	• • • •	1040	1020	6.55		
20		1150	1019	8.49		
21	• • •	1225	1019	8.75	0.522	• • •
22	78.0	1880	1026	10.60		
23		1150	1022	10.21		
24		1130	1018	9.28	,	
25	• • • •	1170	1021	11.30	]	
26	• • • •	1080	1028	11.79	i	
27	• • • •	1460	1017	10.60		
28	• • •	1360	1019	10.53	0.495	• • •
29	• • •	1360	1015	10.28		
30	• • • •	1450	1014	9.87		
May 1		985	1020 1022	8.76 8.86	K	
2 3		1200	1022	8.88		
8 4	77.0	1475 1150	1016			
5	77.9	1400	1023	7.78 9.57	0.271	
6	• • •	1840	1018	7.64	0.271	• • •
7	• • • •	1040	1020	8.73		
8	78.1	895	1021	6.89	l J	
ŷ	10.1	1480	1024	6.13		
10	l	1125	1024	7.60		
11		1290	1020	8.59	0.401	
12	78.1	845	1023	8.57	V. 201	
18		1170	1023	8.92	Į)	

BELLIS.

		Body-		Urine.					
Dat	<b>6.</b>	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>8</sub> .		
190	4	kilos	c.c.		grams	gram	grams		
Мау	14		1005	1020	6.09	h l			
·	15	77.1	855	1023	7.75	0.403			
	16		1210	1014	5.81	0.401	• • •		
•	17		1520	1014	7.98	daily av.			
	18		1150	1028	8.35	h			
	19	77.2	1905	1015	9.60	11			
ł	20	77.2	1210	1020	10.67	[ ]			
	21	76.8	815	1027	8.46	0.461			
l	22	76.5	1020	1019	7.71	11 1			
l	23	76.8	890	1023	5.98	11 .			
ł	24	77.1	1030	1022	6.61	[J			
		·							
Daily Feb.		age from	1127	1021		0.531	1.98		
			1121	1021		0.001	1.00		
	Daily average from Feb. 28 to April 12				9.30	1 1			
		age from	• • •	• • • •	0.00		•••		
		to May 24	• • •	•••	8.45		•••		

CALLAHAN.

	Body-	Urine.					
Date.	Body- weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P3O3.	
1904	kilos	0.0.		grams	gram	grams	
Jan. 15	92.2	1665	1023	19.48	1.008		
16	92.7	1800	1024	22.90	1.235		
17	92.7	1500	1028	18.81	0.996		
18	92.8	1490	1027	19.58	1.398		
19	93.6	1600	1029	22.18	1.070		
20	94.0	1610	1024	28.76	0.710		
21	94.0	1870	1025	24.68	1.040		
22	95.0	1730	1025	22.73	1.160		
28	95.0	2150	1025	31.99	1.470		
24	95.0	1450		21.84	0.952		
25	94.0	1220	1028	1			
26	94.0	1810	1027				
27	93.0	1310	1025	11			
28	94.0	1400	1027	17.90	0.988	3.26	
29	94.0	1220	1027	daily	daily	daily	
30	93.0	1115	1030	average	average	average	
31	98.0	870	1030	J			
Feb. 1	98.0	1170	1028	h			
2	98.0	1080	1029				
8	93.0	1850	1027		,	,	
4	92.0	1150	1028	16.92	0.976	2.86	
5	92.0	950	1029			1	
6	92.0	1160	1027				
7	92.0	1100	1030	J			
8	91.0	850	1027	1			
9	91.0	995	1023	11			
10	90.0	670	1027	11			
11	89.5	615	1029	12.28	0.780	2.14	
12	89.5	865	1028				
13	89.5	740	1031				
14	90.0	1020	1024	IJ			
15	90.0	980	1027	11			
16	89.5	715	1029		,		
17	89.0	565	1030				
18	89.0	535	1031	9.88	0.755		
19	89.0	540	1084	11			
20	89.5	880	1027				
21	89.0	1170	1025	IJ			
22	89.0	820	1030	10.26	0.641	1.52	
28	88.0	610	1031	J 10.20	V.UZ1	1.02	
	L			L	L		

CALLAHAN.

	Rody.	Urine.					
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> .	
1904	kilos	c.c.		grams	gram	grams	
Feb. 24	87.0	520	1034	h			
25	88.0	1160	1017	]			
26	87.5	885	1038	10.26	0.641	1.52	
27	88.0	610	1031	daily	daily	daily	
28	87.0	610	1080	average	average	average	
29	86.0	1095	1026	1			
Mar. 1	86.0	780	1027	11			
2	86.0	725	1027	11			
8	85.0	630	1026	11.07	0.796	1.74	
4	85.0	850	1080				
5	85.0	780	1031				
6	85.5	925	1028	l)			
7	85.0	610	1029	1)			
8	85.0	720	1027	11			
9	85.0	565	1032	11 1			
10	84.5	520	1082	7.27	0.598		
11	84.0	570	1082	11 1			
12	84.0	510	1033	! [ ]			
18	84.0	450	1031	l)			
14	88.5	470	1025	l) l			
15	84.0	1030	1022	-			
16	84.0	825	1026				
17	84.0	690	1026	} 8.09	0.645	• • •	
18	84.0	1160	1025				
19	83.5	1820	1010				
20	84.0	695	1029	ן יו			
21	83.5	760	1027	)	j		
22	88.0	610	1029				
23	83.0	1180	1020				
24	88.0	620	1082	8.18	0.711	• • •	
25	83.0	560	1032				
26	83.0	820	1020				
27	83.0	1000	1027	ן יו			
28	82.0	485	1033	)			
29	82.0	590	1029				
80	82.0	1300	1020				
81	82.5	1590	1020	7.60	0.587	• • •	
Apr. 1	82.0	1600	1018				
2	82.0	610	1026				
8	81.5	430	1026			J	

CALLAHAN.

Date.	Body-	Urine.					
Date.	weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P2O3.	
1904	kilos	c.c.		grams	gram	grams	
Apr. 4	81.0	290	1030	]			
5	82.0	550	1083				
6	82.0	860	1023	[]			
7	82.0	880	1021	6.04	0.458	1.55	
8	82.0	820	1015	daily	daily	daily	
9	82.0	720	1028	average	average	average	
10	82.0	490	1030	ן ון			
11	82.0	540	1022	4.98	)	ļ	
12	82.0	680	1026	ויון	1		
13	82.0	670	1030	8.64			
14	82.0	755	1026	10.24	0.528	•••	
15	82.0	555	1026	5.29			
16	82.0	1790	1017	14.18	1 '	1	
17	83.0	870	1029	9.97	J	1	
18	88.0	990	1020	6.95	)	l	
19	83.0	735	1022	7.82	1	1	
20	88.0	930	1023	8.81	1		
21	83.0	700	1032	10.16	} 0.585		
22	83.0	1220	1024	11.13	1	ł	
23	83.0	860	1027	7.89	İ	ł	
24	83.0	1140	1015	7.59	j	l	
25	83.0	520	1026	8.02	0.544	l	
26	83.5	670	1028	8.08	5 0.011		
27	88.0	1200	1019	7.92	1		
28	83.0	1340	1020	8.84		i	
29	83.5	900	1032	13.28	}		
30	88.5	1250	1022	13.50		ļ	
May 1	84.0	1160	1030	16.77		1	
2	84.0	1080	1026	11.78		l	
8	84.0	960	1029	16.13	0.453	•••	
4	88.5	1140	1035	19.86			
5	83.5	1290	1081	18.35			
6		780	1040	16.43			
7	88.0	870	1033	14.67			
8	83.0	930	1030	14.68	J		
9	83.0	1050	1022	11.86	1		
10	83.0	960	1021	8.18	1		
11	88.0	1020	1024	9.21	0.666		
12	83.0	840	1024	9.08	V.000	•••	
13	83.5	930	1025	10.05	1		
14	83.5	1200	1016	5.55	j		

CALLAHAN.

Date. Body- weight.				Urine.		
Date.	weight.	Volume. 24 hours.	8p. Gr.	Nitrogen.	Uric Acid.	P308.
1904	kilos	c.c.		grams	gram	grams
May 15	88.5	1110	1018	8.99	li l	
16	83.5	1050	1018	9.64	il I	
17	84.0	900	1024	8.37		
18	83.5	1020	1018	8.20	0.666	
19	84.0	945	1022	9.88	11	• • • •
20	84.0	870	1023	8.72	daily	
21	83.0	1160	1014	7.24	average	
22	82.0	420	1017	5.72	j	
23	82.5	480	1026	7.78		
24	82.5	720	1024	10.07		
25	82.0	360	1080	5.64		
26	82.0	510	1029	7.68		
27	82.0	576	1028	6.60		
28	81.5	900	1016	6.59		
29	81.5	570	1023	6.60		
30	81.0	945	1022	6.47		
81	81.5	870	1026	10.44	l l	
June 1	81.5	840	1024	9.02	,	
2	81.5	810	1021	7.78	l	
3	82.0	540	1021	5.67		
4	82.0	450	1025	6.10	l	
5	82.0	480	1034	8.21		
6	82.5	480	1030	6.91		
7	82.5	570	1029	9.51		
8	83.0	650	1028	10.49		
9	83.0	570	1024	7.46		
10	88.0	620	1026	8.40		
11	88.0	690	1027	9.11		
12	88.0	590	1025	7.90		
13	83.0	840	1025	10.74		
14	83.5	780	1021	7.86		·
15	83.0	650	1029	10.37		
Daily ave		809	1026		0.624	1.74
Daily aver Feb. 8 to	April 12			8.56		
Daily aver April 13	rage from to June 15			9.52		• • •

848 PHYSIOLOGICAL ECONOMY IN NUTRITION DONAHUE.

		Urine.						
Date.	Body- weight.							
	weißtre	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>2</sub> .		
1904	kilos	c.c.		grams	gram	grams		
Jan. 19	64.5	1500	1017	17.46	0.872			
20		1060	1027	14.88	0.756			
21	64.5	910	1088	14.41	0.525			
22	• • •	1150	1028	13.25	0.599			
28	• • •	810	1026	11.76	0.428			
24	• • •	790	1030	17.91	0.672			
25		820	1028	1)				
26		960	1024	11				
27	• • •	800	1026			4.5.		
28		870	1025	10.67	0.498	1.90		
29	• • •	740	1025	daily	daily	daily		
80	63.5	655	1029	average	average	average		
81	• • •	790	1026	12				
Feb. 1	• • •	570	1029	11				
2	• • •	780	1027	11				
8	• • •	760	1029		0.450	1.01		
4	• • •	680 930	1030 1025	10.34	0.458	1.81		
5 6	•••	930 820	1026	li :				
7	• • •	600	1020	11				
8	68.0	740	1025	K				
9	00.0	565	1020					
10	• • •	710	1027					
11	•••	870	1025	9.24	0.470	1.76		
12		705	1025	0.22	0.210	1.,0		
13		740	1026	[]				
14		595	1027	11				
15		860	1026	1				
16	63.2	830	1024					
17		780	1026	11				
18		735	1026	11				
19		720	1028	7.47	0.465	• • • •		
20	68.0	690	1027					
21	• • •	900	1025	11				
22	62.7	775	1027	IJ				
23		600	1028	1)				
24		920	1027					
25		700	1027	7.38	0.552	1.56		
26		1040	1023	11				
27		600	1031	IJ.				
		·		1				

DONAHUE.

Date	Body-	Urine.					
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>8</sub> .	
1904	kilos	c.c.		grams	gram	grams	
Feb. 28		710	1024	7.38	0.552	1.56	
29		850	1026	1			
Mar. 1		910	1028	11			
2	62.0	740	1027			!	
8		710	1027	7.79	0.448	1.91	
4		910	1027	daily	daily	daily	
5		725		average	average	average	
6		940	1025				
7	62.7	700	1025	1)			
8		720	1024				
9		1020	1022				
10	• • •	700	1026	6.72	0.863		
11		1180	1021			•	
12		780	1022				
13	• • •	940	1018	}			
14	• • •	930	1018	1			
15	62.7	985	1018				
16	• • •	875	1019	11			
17		515	1024	6.33	0.504		
18	• • •	960	1019				
19	• • •	880	1022				
20	62.7	720	1025	1			
21	• • •	780	1023	]			
22	• • •	920	1024	11	'		
23	• • •	850	1020	5.61	0.863		
24		720	1026		1 0.000		
26	62.7	780	1024				
27	• • •	970	1026	Į.	1		
28	• • •	860	1021	ו			
29	• • •	990	1022				
80	<b>62</b> .8	1250	1023				
81	• • •	990	1020	7.17	0.408	1.90	
Apr. 1	• • •	910	1022	11		i i	
2	***	980	1028				
8 4	62.8	870	1026	K			
5	62.8	1080 740	1025 1028				
6	• • •	740 850	1028	9.91	0.593	1 20	
7	• • •	740	1027	8.81	0.588	1.79	
8	• • •	1010	1028				
°	•••	1010	1020	,			

#### DONAHUE.

	Rody.			Urine.		
Date.	Body- Weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>8</sub> .
1904	kilos	0.0.		grams	gram	grams
Apr. 9		700	1029	) 9.91	0.598	1.79
10		900	1025	daily av.	daily av.	daily av.
11		1150	1020	ا محم	1	
12		1230	1021	9.59		
13		1000	1021	7.38	ł	
14		710	1025	6.18	0.348	
15		920	1022	8.78		
16	63.0	700	1025	6.89	1	
17		775	1024	8.28	J	
18		1050	1020	7.69	)	
19		910	1021	6.55		
20		1140	1020	6.64		
21		1055	1022	6.71	0.899	• • •
22		1020	1020	6.79		
23		700	1025	6.34	1	
24	<b></b>	825	1022	7.18	}	
25		850	1022	6.27	í	
26	l	1010	1021	7.09		
27		1150	1021	7.31	. 1	
28		950	1021	6.38	0.411	
29		950	1023	8.61		
80		525	1028	6.96	1	
May 1	68.0	750	1022	7.65		
2		775	1022	6.88	ń	
8		710	1022	6.01		
4		1075	1021	7.22	1	
5		700	1021	5.34	0.120	
6		950	1024	7.41		
7		950	1028	6.72		
8	l	745	1024	5.90	J	
9		910	1020	6.61	1	
10		720	1023	6.91		
11	63.0	750	1024	7.11		
12	63.0	700	1022	6.51	0.865	
18		710	1024	6.94		
14		825	1025	8.68		
15		1100	1020	7.79	J	
16		1000	1021	6.12		
17		1010	1020	5.94		
18		950	1020	5.75	0.848	

DONAHUE.

	Body-		Urine.					
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P308.		
1904	kilos	c.c.		grams	gram	grams		
May 19		1025	1020	6.64	h -			
20		1100	1020	8.45	11	1		
21		900	1027	8.64				
22		750	1025	8.53	0.848	• • • •		
23		750	1022	7.69	daily			
24		920	1023	7.34	average			
25		750 ·	1022	10.22	li .			
26	62.2	775	1021	6.51	'			
27		880	1028	8.18	11 0 400			
28		800	1028	7.06	0.406	• • •		
29		790	1023	8.49	11			
30		850	1022	7.91				
81		1030	1021	7.60		• • •		
June 1		800	1022	6.53				
2		1000	1021	8.16				
3		850	1022	7.40				
4								
5	62.8	650	1027	8.49				
6		850	1023	7.33				
7		910	1022	7.48				
8		770	1022	8.87	l			
9		910	1025	9.50				
10		850	1025	8.01				
11		650	1025	7.72				
12		700	1025	8.36		<b>.</b>		
13		930	1022	9.15				
14		800	1026	8.45				
15	62.2	700	1027	7.90				
Daily average from Feb. 14		857	1023		0.395	1.79		
Feb. 14	Daily average from Feb. 14 to Apr. 12 Daily average from			7.55				
•	o June 15	• • •	•••	7.89	•••	•••		

352 PHYSIOLOGICAL ECONOMY IN NUTRITION

JACOBUS.

Date.	Body-	Urine.				
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P2O8.
1904	kilos	c.c.		grams	gram	grams
Jan. 15	57.8	630	1032	10.70	0.542	
16		810	1024	11.76	0.960	
17		880	1029	11.09	0.780	
18		780	1080	10.67	0.600	
19		810	1029	12.94	0.595	
20		670	1080	11.98	0.558	
21		730	1029	40.91	0.586	
22		990	1025	12.12	0.594	
28	58.7	825	1021	9.55	0.486	
24		680	1027	9.10	0.449	
25		1260	1018	la l		
26		875	1022	11		
27		680	1028	11 1		
28	57.0	850	1022	9.58	0.442	1.72
29	58.0	760	1020	daily	daily	daily
80	57. <b>7</b>	560	1029	average	average	average
31		660	1028	}		
Feb. 1		700	1023	h		i
2	57.2	567	1030	11		
3		600	1030	11		Ì
4	57.5	582	1026	8.80	、0.548	1.60
5		740	1024			
6	56.8	470	1028	ll i		
7		405	1082	IJ		•
8	56.0	575	1028	lı l		
9	57.2	540	1028			1
10	<b>5</b> 6. <b>6</b>	520	1029	11		
11	57.0	1000	1020	7.73	0.460	1.74
12		920	1015	]]		] '
13	• • •	845	1020			
14		480	1028	[]		
15		750	1025	h l		
16	56.8	810	1022	]]		
17	56.4	760	1020			
18		1100	1017	6.27	0.399	
. 19	56.6	950	1020			
20	56.3	620	1025			
21		580	1027	JJ I	!	
22	57.8	770	1020	1 000	0.501	1 477
28	56.7	795	1024	6.96	0.521	1.47

JACOBUS.

Date.	Body-	Urine.					
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>5</sub> .	
1904	kilos	6.6.		grams	gram	grams	
Feb. 24	56.6	890	1016	l)			
25	56.7	880	1022				
26	56.8	690	1027	6.96	0.521	1.47	
. 27		885	1024	daily	daily	daily	
28		970	1019	Javerage	average	average	
29	56.8	815	1024	]]			
Mar. 1	57.0	760	1027				
2	57.0	690	1025	11			
8	56.8	1720	1012	8.43	0.315	1.75	
4		1010	1019	11			
5		790		11			
6		440	1038	ן ען			
7	56.6	970	1019	]]			
8		1130	1018				
9		440	1081	7.00	0.505		
10	•••	1480	1016	7.88	0.535	• • •	
11		1800	1015				
12	• • •	1165	1018	11			
13		1580	1011	را		,	
14 15	56.8	1200	1015	]]			
16	• • • •	1830 1065	1010 1015	11			
17	56.8	1005	1018	5.98	0.454		
18		1030	1013	0.80	0.303	• • • •	
19	56.8	1445	1018	11			
20	56.8	975	1012				
21		870	1018	15			
22		1400	1014				
23	57.3	840	1020				
24		1020	1021	8.30	0.418	• • •	
25	::: ,		1024				
26	::: '	1196	1017	IJ			
28	:::	1170	1015	1			
29	57.3	1080	1028				
80		1225	1021	[]			
81		1175	1020	9.59	0.471	• • • •	
Apr. 1		1380 ·	1016				
2		775	1023	11			
8		615	1026	IJ			
4		695	1029	8.71	0.810	1.82	
	1	L	L				

354 PHYSIOLOGICAL ECONOMY IN NUTRITION JACOBUS.

	D. 4-		-	Urine.		
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>8</sub> .
1904	kilos	C.4.	C.4.		gram	grams
Apr. 5		830	1026	<u>ا</u> ا		_
6		762	1019			
7	57.0	1205	1026	8.71	0.010	100
8		1180	1015	11	0.310	1.82
9		1035	1017	daily	daily	daily
10		1300	1015	average	average	average
11		1045	1026	7 00	1	
12		610	1024	7.88		
18		410	- 1027	6.81	1	
14		600	1025	6.84	0.460	
15		1160	1017	8.49		
16	56.8	820	1021	7.72		
17		555	1029	6.96	J	
18	• • •	610	1026	7.36	1	,
19		515	1028	6.67	1	
20		710	1028	7.71	1	
21		480	1026	6.94	0.441	
22	55.0	565	1026	7.19		
· 23	• • •	460	1031	7.16		
24		550	1027	6.27	j	
25		750	1020	6.30	1	
26		1170	1017	7.98		
28		1010	1019	6.06	0.505	
29		1110	1018	8.59	0.507	• • •
80		1050	1020	8.44	1	
May 1		715	1030	9.44	}	
2	56.6	1460	1014	10.50	1	
8		1185	1019	9.45		
4		1270	1019	10.52		
5	56.6	1010	1019	9.70	0.382	
6		1090	1021	9.94		
7		1100	1016	8.65		
8		765	1024	9.78	j	
10		555	1026	7.96	)	
11		640	1027	6.60		
12		1160	1016	7.03	0000	
18		665	1025	7.66	0.298	• • •
14		1240	1014	6.61		
15	56.6	595	1027	7.89	J	
16		1460	1012	7.45		• • •
<u></u>				·		

JACOBUS.

	Body-			Urine.		
Date.	Body- weight.	Volume. 24 hours,	Sp. Gr.	Nitrogen.	Uric Acid.	P3O9-
1904	kilos	c.c.		grams	gram	grams
May 17	• • •	1620	1012	6.80		
18		2010	1012	6.75	h l	
19		2050	1010	6.27		
20		1180	1018	7.29		
21		880	1020	7.07	0.876	
22		730	1025	7.62	daily	
28		450	1029	6.48	average	
24		540	1028	7.64	IJ	
25		950	1017	7.53	1)	
26	56.8	555	1023	5.88		
27		1060	1014	7.25	11 1	
28		1020	1019	6.37	} 0.308	
29		690	1022	6.84	11	
30		590	1025	6.69		
<b>3</b> 1		1290	1015	8.86	[].	
June 1		1685	1010	7.85		
2		1475	1012	8.23		
8		400	1024	5.21		
4		705	1019	6.73		
5		1010	1018	6.80		
6		600	1024	5.69		
7		925	1016	6.72		
8	<b> </b>	1015	1014	7.80		• • • •
9		925	1019	6.05		
10	57.0	1285	1012	5.25		
18		855	1018	8.86		
14		735	1019	6.97		
15	57.0	495	1028	8.82		•••
Feb. 1	Daily average from Feb. 1 Daily average from		1031		0.433	1.67
Feb. 1 t	o April 12			7.74		
Daily ave April 13	rage from to June 15			7.43		

SCHENKER.

	Rody.	Urine.									
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>2</sub> O <sub>8</sub> .					
1904	kilos	c.o.	0.0.		gram	grams					
Jan. 16	73.5	1140	1026	16.21	0.688						
17		2180	1020	18. <del>44</del>	0.990						
18		1155	1024	14.97	0.745						
20		1090	1027	19.82	0.797	• • •					
21		1515	1027	20.63	0.966						
22		1675	1019	15.38	0.653						
28	71.7	1100	1022	13.40	0.681						
24		1890	1021	14.18	0.719	• • • •					
<b>2</b> 5		1410	1019	1							
26		840	1027	11							
27		925	• • •	11							
28		1140	1028	13.28	0.698	2.08					
29	72.8	1025	1024	daily	dail <b>y</b>	daily					
80		1900	1015	average	average	average					
81		1105		[2							
Feb. 1	• • •	1320	1028	}							
2	71.6	885	1029	11	1						
8	• • • •	1260	1028			0.00					
4		1480	1020	13.86	0.698	2.16					
5	:::	970	1027	11							
6	71.4	1150	1026	11							
7	1 :::	1270	1022	J	l						
8	71.5	1270	1022	11	ł						
9	71.2	940	1028	11							
10	:	980	1027			000					
11	71.4	845	1029	12.84	0.654	2.00					
12		800 620	1027 1031	11		]					
18		890	1031	11							
14	700	1020	1025	ľ.							
15	70.9	1490	1029								
16	• • • •	1180	1019	11							
17	• • • •	1230	1024	6.34	0.638						
18 19	• • • •	950	1022	0.04	V.038						
20		1510	1029	11							
20 21	• • •	1325	1024								
21 23		510	1020	K							
23 24	71.9	1140	1024								
24 25	11.8	1730	1016	8.85	0.624	2.17					
26	• • •	1145	1022								
20	• • •	1140	IVAL	l'							

#### SCHENKER.

	Body-	Urine.									
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>6</sub> .					
1904	kilos	c.c.		grams	gram	grams					
Feb. 27		2015	2015 1018		0.624	2.17					
28		1150	1028	daily av.	daily av.	daily av.					
29		1280	1028	h l							
Mar. 1	71.8	1245	1026	11							
2		770	1030	11							
8	70.8	1480	1016	11.49	0.799	2.43					
4		1325	1028	]] ]							
5		1960	1014	11 1							
6		1280	1021	ز	-						
7	• • •	1900	1016	1)							
8	72.4	1760	1016	11 1							
9	• • •	910	1026	11 1		1					
10		2050	1015	10.11	0.784						
11	72.5	940	1020	11 1		1					
12		2200	1002	11	-						
18		2790	1011	I) I		ì					
14		1880	1014	lı l		1					
15	72.8	1630	1012			İ					
16		1070	1022	11 1		1					
17		2040	1012	11.08	0.797						
18		1655	1021	11 1							
19	• • •	1485	101 <b>4</b>	H							
20	• • •	2550	1013	[) [							
21	71.9	2000	1014	la l							
22	• • •	2170	1012	10.14	0.551	• • • •					
28	• • •	1670	1014		0.001						
24	72.3	2020	1015	ו יו							
26		875			• • •	• • • •					
27	• • •	1520	1018	n I							
28		1675	1017	]] ]							
29		1175	1025								
30	• • •	1110	1021	10.85	0.660	• • •					
81	• • •	2340	1012								
Apr. 1	• • •	1440	1021								
2	• • •	1200	1021	ן ין							
8	•••	:::		1) 1							
4	• • •	1110	1022	11	0 200						
. 5	• • •	785		11.64	0.782	• • • •					
6	:::	1180	1028	11							
7	74.1	945	1027	'							

#### SCHENKER.

	Body-	Urine.									
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Urio Acid.	P308.					
1904	kilos	c.c.		grams	gram	grams					
Apr. 8		1050	1024	h	İ						
9		925	1022	11.64	0.782						
10		1555	1020	J daily av.	daily av.						
11	73.8	1440	1015	8.44	ì						
18	• • •	1250	1023	12.23							
14		1330	1026	10.06	0.620						
15	• • • •	1880	1018	12.41	11 0.020						
16		2440	1013	11.27							
17		1680	1018	10.78	J.						
18	78.3	1640	1018	11.22	}						
19	1 :::	1420	1020	12.95							
20	74.2	1440	1019	9.85		l i					
21		1670	1018	10.22	0.746	• • • •					
22	1 :::	1220	1019	9.52	l						
23	78.3	1640	1021	9.05							
24	• • • •	2200	1016	9.50	J						
25	• • •	1400	1020	9.07	}						
26		1400	1020	8.15							
27	• • • •	1710	1017	9.18							
28	• • • •	2220	1011	9.59	0.594	• • •					
29	78.4	1940	1015	11.29							
80	• • • •	980	1025	10.76							
May 1	••,•	1625	1015	10.05	,						
2	• • •	1950	1014	8.89	1						
8	• • •	1795	1014	8.62	1						
4	• • •	2400	1010	8.50	0000						
5		1895	1019	11.03	0.298	• • •					
6	78.7	1920 2300	1019	11.75	1						
7	• • •		1014	11.87	}						
8 9	• • •	1880	1018	11.51	(						
10	• • •	2095 1360	1014	11.46 9.96	1						
10	• • •	1040	1016 1022	9.90 8.74	0.229						
12	•••	1250	1022 1024	11.02	0.228	• • •					
15	• • •	1250 1215	1024 1020	8.68	1						
, 16	• • •	1450	1020	8.96	,						
17	• • •	1725	1018 1014	8.29	• • •	• • • •					
18	• • •	1420	1014	8.29 8.77	$\cdot \cdot \cdot$	• • •					
19	72.7	990	1020	8.78	0.427						
20		2390	1014	12.48	[ V.321	• • •					
	•••	2080	101.2	19.40	,						

#### SCHENKER.

Date.	Body-	. Urine.									
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P2O5.					
1904	kilos	0,0.		grams	grams	grams					
May 21	72.7	1000	1025	8.76	1)						
22		1570	1018	9.98	0.427						
23		1780	1015	10.04	1 1	• • •					
24		1210	1018	8.71	daily av.						
25		1390	1016	9.09	l)						
26	72.5	1380	1015	9.11							
27	72.1	1920	1018	9.56	1						
28		1870	1016	11.22	0.842	• • •					
29	• • •	1635	1015	8.88	1 1						
80	• • •	1670	1017	8.52	[]						
31	• • •	2190	1018	9.46		• • •					
June 1	• • •	1140	1016	6.02		• • •					
2		1520	1018	8.94		• • •					
8		1610	1017	8.50							
4	72.2	1880	1020	8.94		• • •					
5		795	1022	8.11		• • •					
6	• • •	1590	1018	9.83							
7	• • •	2000	1012	9.48							
8		1800	1016	12.10		• • •					
9	71.9	1500	1018	9.68	• • •	• • •					
Daily aver	rage from	1500	1019		0.694	2.20					
Daily aver	age from			l							
-	April 10			10.37							
Daily aver	rage from										
	to June 9	• • •	• • •	9.82	• • •	• • •					

STAPLETON.

Urine. Body-weight. Date. Volume. Uric Acid. Sp. Gr. Mitrogen. P<sub>1</sub>O<sub>3</sub>. 24 hours. 1904 kilos c.c. grams gram grams 77.2 1350 1026 16.20 0.941 Jan. 15 16 77.2 2200 1020 20.83 1.163 2000 1025 21.00 1.095 17 77.2 1650 1026 21.78 0.897 18 78.1 19 77.1 1585 1029 20.64 1.139 20 77.0 1890 1028 19.26 0.480 1026 21 77.1 1560 20.81 0.971 22 1620 1028 0.785 77.1 18.87 23 78.0 1810 1025 21.72 0.624 1024 17.42 0.890 24 77.0 1480 76.0 1029 25 990 26 76.0 1280 1026 77.0 885 1080 27 28 77.1 1660 1021 14.95 0.841 2.83 29 76.1 1005 1029 daily daily daily 80 77.1 1810 1027 average average average 81 76.1 800 1031 Feb. 1028 76.1 1090 1 76.1 1025 2 1175 8 76.0 1120 1030 4 76.0 950 1031 13.48 0.743 2.34 5 76.0 1150 1023 6 76.0 1165 1029 7 76.0 770 1033 8 76.0 825 1030 9 76.0 1160 1026 10 76.0 1140 1023 11 76.0 1160 1028 12.72 0.712 2.74 970 1026 12 76.0 18 76.0 1115 1024 14 76.0 1860 1023 15 76.0 1040 1026 16 76.0 1020 1023 76.0 1380 1021 17 1029 12.36 0.755 18 76.0 880 1030 76.0 19 945 76.0 1940 1017 20 21 76.0 2670 1016 22 76.0 1590 1023 13.03 0.967 2.56 1870 28 76.0 . . .

STAPLETON.

Body-		Urine.									
Date.	weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P <sub>3</sub> O <sub>8</sub> .					
1904	kilos	c.c.		grams	gram	grams					
Feb. 24	76.0	870	1026	1		1					
25	76.0	1275	1024								
26	76.0	1140	1025	} 18.03	0.967	2.56					
27	76.0	1930	1018	daily	daily	daily					
28	76.0	1120	1026	average	average	average					
29	76.0	1140	1028	h							
Mar. 1	76.0	1800	1022	[]							
2	76.0	1295	1028	11							
3	77.0	825	1030	12.91	0.803	2.59					
4	76.0	1860	1021								
5	76.0	1280	1026								
6	76.0	1155	1029	l)							
7	76.0	880	1031	1							
8	76.0	860	1081								
9	76.0	940	1029								
10	76.0	550	1034	11.02	0.707						
11	76.0	780	1028								
12	77.0	790	1026	11							
13	76.0	700	1030	IJ							
14	76.0	830	1027	1							
15	76.0	1650	1014	11	·						
16	76.0	1120	1019	11							
17	76.0	690	1027	10.26	0.757	• • • •					
18	76.0	1170	1024	11							
19	76.0	2230	1010	11							
20	77.2	1180	1025	ļ J							
21	76.0	540	1032	)							
22	76.0	880	1030								
23	76.0	990	1028								
24	76.0	1180	1024	11.55	0.794	• • •					
25	76.0	1470	1025								
26	76.0	1280	1025								
27	76.0	1240	1021	<u> </u>							
28	76.0	1020	1028	[]							
29	77.0	1440	1018								
80	76.0	1595	1018		0.00=						
81	76.0	1210	1017	11.14	0.667	• • •					
Apr. 1	76.0	1940	1018								
2	76.0	1110	1028								
8		1185	1026	ין							

	Body-		Urine.									
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P2O3.						
1904	kilos	0.0.		grams	gram	grams						
Apr. 4		1420	1027	h								
5	• • •	1260	1029									
6	• • •	1095	1026	17.56	0.929	2.77						
7	• • •	1815	1030	daily	daily	daily						
8	• • •	1180	1029	average	average	average						
9	:::	1140	1030	) average	average	a.c.ago						
11	75.0	1285	1031	16.43	1							
12	• • •	940	1028	ו								
13	• • • •	750	1027	9.59								
14		720	1026 1028	10.45	0.550							
15	• • • •	910 815	1028	10.70								
16 17	• • • •	1110	1021	6.32 5.38								
	• • • •	820	1012	4.58	,							
18 19		750	1024	7.88	1							
20	74.0	880	1025	7.18								
20 21	12.0	1005	1024	9.17	0.418							
22	• • •	1090	1027	9.74	0.220	• • •						
28		1085	1019	8.17								
24	• • • • • • • • • • • • • • • • • • • •	990	1020	8.26	. 1							
25		1150	1021	9.21	ń							
26	l :::	980	1025	9.23								
27	75.0	1570	1019	8.29								
28		1040	1025	9.05	0.599							
29		1020	1022	9.98								
80		1000	1026	9.40								
May 1		1025	1025	10.56	J							
2		1070	1027	11.25	1							
8		1585	1018	10.87								
4	75.0	1805	1023	8.70								
5		1350	1022	9.88	0.877							
6		920	1024	7.84								
7		900	1025	9.28								
8		930	1025	11.50	J							
9		875	1027	11.81	h							
10	• • • •	1010	1022	8.85								
11	• • • •	1010	1023	9.26		ł						
12	74.0	600	1030	9.29	0.625	•••						
13		940	1028	10.60								
14	• • •	975	1028	7.87	J							

STAPLETON.

	Pode	Urine.									
Date.	Body- weight.	Volume. 24 hours.	Sp. Gr.	Nitrogen.	Uric Acid.	P2O3.					
1904	kilos	<b>c.c.</b>		grams	gram	grams					
May 15		800	1025	6.68	0.625						
16		1200	1019	8.14							
17		515	1030	6.58							
18		790	1029	9.67	l) l						
19	74.1	745	1029	8.58	li i						
20	74.1	880	1022	9.51							
21	74.1	1100	1027	9.64	0.691						
22		890	1028	9.56	daily						
28		690	1020	8.08	verage						
24	76.0	905	1025	9.04	IJ						
25		665	1031	8.46	h .						
26	74.5	630	1026	7.45	0.668	2.27					
27		950	1024	8.88	0.008						
28	78.0	850	1027	10,60	1'	daily av.					
80		1060	1020	10.88							
81		1640	1015	8.56							
June 1		1280	1013	9.37							
2	l	1180	1019	10.84							
8		620	1024	6.96							
4	l l	910	1020	11.47							
5		880	1019	11.09							
6		855	1025	11.14							
7		1250	1017	8.78							
8	l	885	1025	11.10							
9	74.0	780	1020	6.35							
10		1540	1017	8.87		`					
11		750	1015	5.85							
12	78.4	1265	1015	9.71		• • •					
Jan. 25			1024		0.699	3.64					
Jan. 25 t	Daily average from Jan. 25 to April 12			13.19							
Daily ave April 18	rage from to June 12	·	• • • •	9.00		•••					

It is plain from the foregoing results, that all the men of this group, like the members of the professional group, experienced no difficulty in reducing in large measure their rate of proteid metabolism. The intake of proteid food was steadily diminished, with a corresponding diminution in the extent of nitrogen metabolism. Take as an illustration the average daily output of nitrogen from April 13 to June 15, a period of sixty-three consecutive days:

AVERAGE DAILY EXCRETION OF METABOLIZED NITROGEN'
FOR THE LAST TWO MONTHS OF THE EXPERIMENT.

			grams	
G. W. Anderson	•	•	8.81	
W. L. Anderson	•		10.07	
H. S. Bellis			8.45*	Grand average for this
W. H. Callahan			9.52	$rac{1}{2}$ period = 8.81 grams of $rac{1}{2}$
M. Donahue			7.39	nitrogen per day.
C. S. Jacobus .			7.43	ntuogen per day.
H. R. Schenker		•	9.82	
John Stapleton.		•	9.00	

An excretion of 8.81 grams of nitrogen through the kidneys corresponds to the metabolism of 55 grams of proteid matter. Compare this average amount of proteid matter metabolized each day with the figures obtained during the preliminary period of ten days, when the men were living on their ordinary diet. Then, many of the men were excreting nitrogen at the rate of 17 to 22 grams per day. In a general way, we may safely say that all these men during the last two months of the experiment were living on about one-half the proteid food they were formerly accustomed to take.

Further, the average daily excretion of nitrogen for the preceding seventy-three days, i. e., from February 1 to April 13, was in most instances nearly, if not quite, as low as during the last two months of the experiment, so that we are certainly justified in the statement that these men—trained

<sup>\*</sup> This average covers the period from April 13 to May 24 only, as Mr. Bellis was compelled to withdraw from the experiment on the latter date, owing to an accident in the gymnasium.

athletes, doing athletic work more or less strenuous — were able to practise during this long period marked physiological economy in the use of proteid food, equal approximately at least to a saving of full fifty per cent in proteid matter.

The individual tables must be carefully studied, however, in order to trace out the changes in detail in the rate of nitrogen metabolism, and in so doing much information will be obtained regarding modification in the excretion of uric acid. a matter to be discussed in another connection, later on. Further, it is interesting to note in the tables the changes in bodyweight of the men. Some of the men, like Dr. Callahan, who were abundantly supplied with adipose tissue, lost very considerably in body-weight, but eventually came to a standstill, with establishment of body equilibrium, under the changed dietary habits. Some of the men reached this condition of equilibrium much more quickly than others. Dr. Callahan who suffered a large loss in body-weight — to his great gain, as he expressed it -dropped from 92.2 kilos to 83 kilos in two months, but from March 22 to June 15 his bodyweight, while naturally showing fluctuation, did not fall again permanently.

What now was the amount of metabolized nitrogen per kilo of body-weight in these men toward the close of the experiment? Taking the average daily nitrogen excretion for the period from April 13 to June 15, and the body-weights of the men at this same period, as indicated in the accompanying table, we have the following figures:

					Body- weight.		Average daily Nitrogen excreted.	Metabolized Nitro- gen per kilo of body-weight.
						kilos	grams	gram
Bellis	•	•	•			<b>7</b> 8	8.45	0.108
Callahan.						83	<b>9.52</b>	0.114
Donahue.		•			•	62	7.39	0.119
Stapleton						75	9.00	0.120
Anderson,	G.	W.				71	8.81	0.124
Jacobus .						56	7.43	0.132
Schenker.	•	•				73	9.82	0.134
Anderson,	$\mathbf{w}$	. L.		•		61	10.07	0.165

These figures, with one exception, show as low a proteid metabolism per kilo of body-weight as was obtained with the soldiers on a prescribed diet, yet these men were athletes accustomed to vigorous muscular exercise, and likewise accustomed to the eating of relatively large amounts of proteid food. Theoretically, it might not be expected that these men would drop to as low a level as men who were not addicted to the consumption of excessive amounts of proteid foods, yet for two months, and practically for a period of four months, these University students easily maintained themselves at this lower level of nitrogen metabolism.

From April 26 to June 13, at the request of the students themselves, the daily diet was prescribed; not, however, as regards the quantity of food to be eaten, but merely as to its character. The men ate at the University Dining Hall, and it was a simple matter to have their table supplied with a special dietary. The following dietary was therefore prepared for their use.

It is not to be understood that the men took all that the daily list provided, but they made their selections from the menu, and in quantity took what they deemed necessary, or what satisfied their appetites. It may be added that the men were all well content with the variety provided and expressed themselves, many times, as better pleased with a simple dietary of this kind than with the heavier proteid foods of earlier days. It should be added that Dr. Callahan was compelled to be absent from New Haven during a large portion of the period covered by this dietary, hence his nitrogen excretions do not correspond in quantity with the nitrogen of the above rations.

## Tuesday, April 26, 1904.

Breakfast. — Banana, boiled hominy with sugar and cream, coffee, rolls, butter.
Lunch. — Spaghetti, stewed tomatoes, potatoes, boiled onions, bread, butter, coffee, fried hominy with syrup.

Dinner. — Split-pea soup, fried bacon with French fried potato, spinach, bread, butter, stewed prunes, lettuce-celery-apple salad, cream puffs, coffee.



STAPLETON

Photograph taken in the middle of the experiment, in April.



## Wednesday, April 27, 1904.

Breakfast. — Fruit, farina with cream, coffee, rolls, butter, baked potato.

Lunch.—Fried oysters, mashed potato, bread, butter, coffee, string beans, aliced banana with cream.

Dinner. — Cream of celery soup, codfish-cakes, boiled potato, boiled lima beans lettuce-orange salad, ice cream, coffee.

## Thursday, April 28, 1904.

Breakfast. — Banana, coffee, rolla, cream, butter, fried hominy and syrup. Lunch. — Fried sweet potato, cold tongue, baked potato, bread, butter, coffee, baked apple with cream.

Dinner. — Vegetable soup, Hamburg steak made with plenty of bread, etc., Lyonnaise potato, baked potato, spinach, pie, coffee.

### Friday, April 29, 1904.

Breakfast. — Fruit, Indian meal, boiled, with sugar and cream, baked potato, rolls, coffee.

Lunch. — Clam chowder with crackers, farina croquettes, stewed tomato, griddle cakes with syrup, coffee, bread, butter.

Dinner. — Soup, fish, mashed potato, radishes, string beans, cranberry sauce, bread, butter, lettuce salad, lemon pie, coffee.

## Saturday, April 30, 1904.

Breakfast. — Banana, fried Indian-meal, syrup, baked potato, coffee, rolls.

Lunch. — Tomato purée, baked macaroni, baked potato, sliced ham, baked apple and cream, bread, butter, coffee.

Dinner. — Soup, small sausage fried, mashed potato, boiled sweet potato, spinach, stewed tomato, strawberry short cake, coffee.

## Sunday, May 1, 1904.

Breakfast. — Fruit, boiled rice, sugar, cream, coffee, rolls, butter.

Dinner. — Vegetable soup, stewed chicken, cranberry sauce, boiled potato, boiled onion, stewed corn, water ice, coffee, bread, butter.

Supper. - French fried potato, bacon, Waldorf-salad, bread, butter, pie.

## Monday, May 2, 1904.

Breakfast. — Banana, malto-rice, cream, coffee, rolls.

Lunch. - Baked beans, catsup, baked potato, stewed prunes, apple pudding.

Dinner. — Barley soup, string beans, boiled onion, mashed potato, fried bacon, cranberry sauce, bread, butter, sliced banana, cream, coffee.

### Tuesday, May 3, 1904.

- Breakfast. Fruit, boiled Indian-meal, sugar, cream, baked potato, butter, rolls, coffee
- Lunch. Baked macaroni, French fried potato, boiled spinach, stewed prunes, coffee, bread, butter, fried rice with syrup.
- Dinner. Split-pea soup, fried sausage with Lyonnaise potato, boiled sweet potato, butter, lettuce-orange salad, baked apple with cream, coffee.

### Wednesday May 4, 1904.

- Breakfast.—Banana, farina with sugar and cream, baked potato, coffee, rolls, butter.
- Lunch. One egg on toast, string beans, boiled potato, bread, butter, coffee, pie.
  Dinner. Tomato purée, codfish-cakes, baked potato, boiled lima beans, lettuce-apple salad, bread, butter, cream puffs, coffee.

### Thursday, May 5, 1904.

- Breakfast. Fruit, boiled hominy, sugar, cream, coffee, rolls, butter.
- Lunch. Cold ham, fried sweet potato, baked potato, cream, butter, coffee, cabinet pudding, vanilla sauce.
- Dinner. Scotch broth, one lamb chop, mashed potato, fried sweet potato, spinach, bread and butter, baked apple and cream, coffee.

## Friday, May 6, 1904.

- Breakfast. One apple, fried hominy with syrup, baked potato, rolls, butter,
- Lunch. Clam chowder with crackers, potato croquettes, sliced tomatoes, bread, butter, griddle cakes and syrup, coffee.
- Dinner. Bean soup aux croutons, boiled halibut, mashed potato, boiled onions with cream, radishes, cranberry sauce, lettuce salad, bread, butter, coffee, one orange.

## Saturday, May 7, 1904.

- Breakfast. Banana, boiled rice, sugar, cream, baked potato, coffee, rolls, butter.
- Lunch. Consommé, spaghetti à l'italienne, boiled sweet potato, bread, butter, cakes, preserves, coffee.
- Dinner. Split-pea soup, roast beef, mashed potato, boiled spinach, boiled sweet potato, bread and butter, strawberries and cream, coffee.

#### Sunday, May 8, 1904.

Breakfast. — Banana, fried rice, syrup, coffee, corn bread and butter, baked potato.

Dinner. — Vegetable soup, stewed chicken, cranberry sauce, string beans, boiled or mashed potato, water ice, bread and butter, cakes, coffee.

Supper. — French fried potato, bacon, lettuce-orange salad, bread and butter, cake, preserves, and coffee.

### Monday, May 9, 1904.

Breakfast. — Fruit, boiled Indian-meal, sugar and cream, coffee, rolls, butter.

I.unch.—Calf's liver, baked potato, stewed prunes, boiled onion, bread and butter, coffee, apple pudding.

Dinner. — Consommé, baked macaroni, fried sweet potato, stewed tomato, preserves, lettuce, bread, butter, cakes, coffee.

## Tuesday, May 10, 1904.

Breakfast. - Fruit, farina, milk, sugar, baked potato coffee, bread, butter.

Lunch. — Egg omelette with jelly, fried bacon, mashed potato, spinach, apple pudding, bread, butter, coffee.

Dinner. — Small fried sausage, boiled potato, rice custard, lettuce-orange salad, bread, butter, coffee, baked apples with cream.

#### Wednesday, May 11, 1904,

Breakfast. — Banana and cream, fried hominy, syrup, baked potato, coffee, bread, butter.

Lunch. — Cream of tomato soup, mashed potato, lima beans, bread, butter, prune souffié, coffee.

Dinner. — Hamburg steak made with much bread, boiled sweet potato, boiled onions, lettuce, lemon pie, bread, butter, coffee.

## Thursday, May 12, 1904.

Breakfast. — Fruit, baked potato, boiled Indian-meal, sugar, cream, coffee, rolls, butter.

Lunch. — Consommé, French fried potato, one egg on toast, rice pudding, apple sauce, coffee, bread, butter.

Dinner. — One chop, boiled or mashed potato, string beans, apple-lettuce salad, lemon pie, bread, butter, coffee.

## Friday, May 13, 1904.

Breakfast. — Banana and cream, fried rice with syrup, rolls, butter, coffee.

Lunch. — Clam chowder, boiled potato, boiled onions, fried bacon, carrots, apple dumpling, bread, butter, coffee.

Dinner. — Split-pea soup (thick), frizzled beef, fried sweet potato, spinach, cranberry tart, bread, butter, coffee, cakes.

#### 110

### Saturday, May 14, 1904.

Breakfast. — Baked apple and cream, boiled hominy, with sugar and cream, baked potato, coffee, rolls, butter.

Lunch. — Cream of celery soup, farina croquettes, with tomato sauce, stewed corn, mashed potato, bread, butter, coffee, fruit.

Dinner. — Fish, boiled potato, boiled onions, bread pudding, preserves, lettuce-tomato salad, small cakes, bread, butter, coffee.

### Sunday, May 15, 1904.

- Breakfast. Fruit, baked potato, boiled oatmeal with sugar and cream, coffee, rolls, butter.
- Lunch. Consommé with croutons, fried rice with syrup, French fried potato, strawberry short-cake with whipped cream, bread, coffee.
- Dinner. Stewed chicken, fried sweet potato, cranberry sauce, celery, string beans, bread, butter, coffee, ice cream, cakes.

#### Monday, May 16, 1904.

- Breakfast.—Banana, griddle cakes and syrup, baked potato, coffee, rolls, butter.
- Lunch. Fried bacon, mashed potato, spinach, bread, butter, rice croquettes with preserves, apple pie, coffee.
- Dinner. Consommé, one lamb chop, mashed potato, string beans, boiled onions, orange-lettuce salad, bread, butter, tapioca pudding, coffee.

## Tuesday, May 17, 1904.

Breakfast. - Banana, farina, cream, sugar, baked potato, rolls, butter, coffee.

Lunch. — Vegetable soup, French fried potato, one egg on toast, rice pudding, apple sauce, bread, butter, coffee.

Dinner. — Small fried sausage, boiled potato, lima beans, lettuce salad, bread, butter, baked apples with cream, rice custard, coffee.

### Wednesday, May 18, 1904.

- Breakfast.—Sliced banana, fried rice, syrup, baked potato, bread, butter, coffee.
- Lunch. Cream of celery soup, farina croquettes, tomato sauce, fried sweet potato, string beans, bread, butter, prune souffié, coffee.
- Dinner. Split-pea soup, Hamburg steak made with much bread, mashed potato, spinach, bread, butter, lemon pie, coffee.

### Thursday, May 19, 1904.

- Breakfast. Sliced banana, boiled hominy, cream, sugar, baked potato, bread, butter, coffee.
- unch. Egg omelette, jelly, French fried potato, boiled onions, fried hominy, syrup, bread, butter, apple pudding, coffee.
- Dinner. Tomato purée, baked macaroni, fried bacon, fried sweet potato, spinach, bread, butter, Indian-meal pudding, coffee.

### Friday, May 20, 1904.

- Breakfast. Sliced orange, fried hominy, syrup, baked potato, bread, butter, coffee.
- Lunch. Fish cakes, boiled sweet potato, mashed potato, lima beans (boiled), bread, butter, bread pudding, coffee.
- Dinner. Consommé, boiled halibut, mashed potato, string beans, bread, butter, rice croquettes, cranberry jam, coffee.

#### Saturday, May 21, 1904.

- Breakfast. Sliced banana, cream, sugar, boiled Indian-meal, baked potato, bread, butter, coffee.
- Lunch. One lamb chop, potato croquettes, fried Indian-meal, syrup, stewed tomatoes, bread, butter, coffee, water ice.
- Dinner. Bean purée, scrambled egg, bacon, French fried potato, lettuceorange salad, farina pudding, prunes, bread, butter, coffee.

## Sunday, May 22, 1904.

- Breakfast. Sliced orange, sugar, boiled oatmeal, cream, baked potato, bread, butter, coffee.
- Lunch. Boiled macaroni, fried rice, syrup, mashed potato, boiled onions, bread, butter, ice cream, cake, coffee.
- Dinner. Cream of celery soup, stewed chicken, French fried potato, mashed potato, spinach, bread, butter, cranberry sauce, strawberry short-cake, cream, coffee.

## Monday, May 23, 1904.

- Breakfast.—Sliced banana, griddle cakes, syrup, baked potato, bread, butter, coffee.
- Lunch. One egg on toast, consommé, French fried potato, lettuce, rice croquettes, syrup, apple sauce, bread, butter, coffee.
- Dinner. Vegetable soup, baked macaroni, fried bacon, potato croquettes, string beans, bread, butter, water ice, coffee.

### Tuesday, May 24, 1904.

- Breakfast. Banana, boiled rice, cream, sugar, baked potato, bread, butter, coffee.
- Lunch. Cream of celery soup, farina croquettes, tomato sauce, boiled onions, mashed potato, bread, butter, stewed prunes, coffee.
- Dinner. Tomato purée, Hamburg steak made with much bread, French fried potato, spinach, farina croquettes, bread, butter, lemon pie, coffee.

### Wednesday, May 25, 1904.

- Breakfast. Banana, boiled hominy, cream, sugar, baked potato, bread, butter, coffee.
- Lunch. Small fried sausage, boiled potato, lettuce salad, fried hominy, syrup, bread, butter, apple sauce, coffee.
- Dinner. Consommé, scrambled eggs, French fried potato, lettuce-orange salad, lima beans, bread, butter, bread pudding, coffee.

#### Thursday, May 26, 1904.

- Breakfast. Banana, boiled oatmeal, sugar, cream, baked potato, rolls, butter, coffee.
- Lunch. One egg on toast, spinach, mashed potato, apple sauce, bread, butter, rice pudding, coffee.
- Dinner. Tomato purée, boiled macaroni, boiled onions, French fried potato, lettuce, bread, butter, tapioca pudding, coffee.

## Friday, May 27, 1904.

- Breakfast. Orange, boiled Indian-meal, sugar, cream, rolls, coffee.
- Lunch.—Clam chowder, potato croquettes, lima beans, bread, butter, strawberries, cream, coffee.
- Dinner. Bean soup, boiled halibut, mashed potato, string beans, rice croquettes, cranberry jam, bread, butter, water ice, coffee.

## Saturday, May 28, 1904.

- Breakfast. Banana, breakfast flakes, sugar, cream, baked potato, rolls, butter, coffee.
- Lunch. One boiled egg, French fried potato, stewed tomatoes, fried Indianmeal, syrup, bread, butter, coffee.
- Dinner. Split-pea soup, baked beans, Boston brown bread, lettuce-orange salad, stewed prunes, bread, butter, ice cream, coffee.

#### Sunday, May 29, 1904.

- Breakfast. Orange, boiled oatmeal, sugar, cream, baked potato, rolls, butter, coffee.
- Lunch. Boiled spaghetti, mashed potato, boiled onions, fried rice, syrup, bread, butter, strawberries, cake, coffee.
- Dinner. Cream of celery soup, stewed chicken, boiled potato, spinach, bread, butter, cranberry sauce, custard pie, coffee.

#### Monday, May 30, 1904.

- Breakfast. -- Banana, boiled rice, sugar, cream, baked potato, rolls, butter,
- Lunch. Consommé, French fried potato, bacon, lettuce-orange salad, wheat griddle cakes, syrup, bread, butter, coffee.
- Dinner. Barley broth, one lamb chop, mashed potato, fried sweet potato, apple sauce, bread, butter, bread pudding, coffee.

### Tuesday, June 7, 1904.

- Breakfast. Banana, boiled rice, cream, sugar, baked potato, rolls, butter,
- Lunch. Vegetable soup, French fried potato, one egg on toast, apple sauce, rice pudding, bread, butter, tea.
- Dinner. One small fried sausage, boiled potato, lima beans, lettuce-orange salad, bread, butter, ice cream, cake, coffee.

## Wednesday, June 8, 1904.

- Breakfast. Banana, breakfast flakes, sugar, cream, baked potato, rolls, butter, coffee.
- Lunch. Cream of celery soup, potato croquettes, string beans, two alices bacon, bread, butter, bread pudding, coffee.
- Dinner. Split-pea soup, boiled halibut, mashed potato, asparagus, bread, butter, cream pie, coffee.

## Thursday, June 9, 1904.

- Breakfast. Orange, boiled hominy, cream, sugar, baked potato, rolls, butter, coffee.
- Lunch. Egg omelette, jelly, French fried potato, boiled onions, bread, butter, wheat griddle cakes, syrup, coffee.
- Dinner. Tomato purée, baked macaroni, dried beef stewed with milk, boiled potato, spinach, bread, butter, Indian-meal pudding, coffee.

### Friday, June 10, 1904.

Breakfast. - Banana, fried hominy, syrup, rolls, butter, coffee.

Lunch. — Clam chowder, mashed potato, boiled lima beans, bread, butter, tapioca pudding, coffee.

Dinner. — Consommé, bluefish (broiled), Lyonnaise potato, asparagus, bread, butter, cranberry jam, lemon pie, coffee.

#### Saturday, June 11, 1904.

Breakfast. — Orange, boiled Indian-meal, cream, sugar, baked potato, rolls, butter, coffee.

Lunch. — Barley broth, corn fritters, mashed potato, two slices bacon, bread, butter, stewed tomatoes, strawberries, cream, coffee.

Dinner. — Bean pureé, scrambled egg, rice croquettes, French fried potato, lettuce-orange salad, stewed prunes, bread, butter, farina pudding, coffee.

#### Sunday, June 12, 1904.

Breakfast. — Banana, boiled oatmeal, sugar, cream, baked potato, rolls, butter, coffee.

Lunch. — Boiled macaroni, string beans, mashed potato, farina croquettes, water ice, cake, coffee.

Dinner. — Cream of celery soup, stewed chicken, French fried potato, lettuce salad, bread, butter, strawberry short-cake, cream, coffee.

## Monday, June 13, 1904.

Breakfast. — Orange, breakfast flakes, sugar, cream, baked potato, rolls, butter, coffee.

Lunch. — Vegetable soup, one egg on toast, fried potatoes, apple sauce, rice croquettes, bread, butter, cottage pudding, tea.

Dinner. — Clam broth, beef stew, with potatoes, carrots and onions, boiled potatoes, bread, butter, apple-lettuce salad, apple pie, cheese, coffee.

#### NITROGEN BALANCE.

We may now consider whether these men, who had so greatly reduced the extent of their proteid metabolism, and who had apparently attained a condition of body equilibrium, were truly in nitrogenous equilibrium, or free from any daily loss of nitrogen. To test this point, a careful and exact comparison of the nitrogen intake and output was made for a period of seven days, commencing on May 18. This was not easy to accomplish, as the men were still allowed freedom in the quantity of food eaten, and also in the choice of food, within the limits of the menu provided, so that a large number of food analyses had to be made. A greater difficulty, however, in obtaining proof of equilibrium lay in the laudable ambition of the men to make a good showing, - for they had all become interested in the main problem, and had acquired great faith in the efficiency of a low proteid ration, - which led them to great caution in the matter of eating during the balance trial, thereby running the risk of diminishing in too great degree the fuel value of the food.

The following data obtained with Donahue are self-explanatory. The figures showing the quantities of food at each meal are instructive, as indicating the general makeup of the daily dietary, both as to quality and quantity.

### Wednesday, May 18 1904.

Breakfast. — Banana 141 grams, bread 60 grams, butter 15 grams, coffee 150 grams, cream 80 grams, sugar 31 grams.

Lunch. — Soup 247 grams, string beans 65 grams, bread 21 grams, butter 80 grams, coffee 150 grams, sugar 21 grams, fried potato 222 grams.

Dinner. — Consommé 150 grams, bread 45 grams, butter 10 grams, mashed potato 150 grams, spinach 200 grams, apple pie 103 grams, coffee 150 grams, cream 75 grams, sugar 28 grams.

Food.	Grams.	Per	cent Nit	rogen.	Total Nitrogen.
Banana	. 141	×	0.23	=	0.324 grams.
Butter $15 + 30 + 10 =$	= 55	×	0.15	=	0.088
<b>Sugar.</b> $31 + 21 + 28 =$	= 80	×	0.00	=	0.000
Cream $80 + 75 =$	= 155	×	0.46	=	0.718
Bread	. 60	×	1.66	=	0.996
Coffee $. 150 + 150 + 150 =$	= 450	×	0.06	=	0.270
Bread	. 21	×	1.60	=	0.336
Soup	. 247	×	0.41	=	1.013
Fried potato	. 222	×	0.32	=	0.710
String beans	. 65	×	0.84	=	0.221
Consommé	. 150	×	0.38	=	0.570
Bread	. 45	×	1.80	=	0.810
Spinach	. 200	×	0.58	=	1.060
Potato	. 150	×	0.38	=	0.570
Pie	. 103	×	0.48	=	0.443
Total nitroger	n in food				8.119 grams.
Total nitroger	a in urine	•			5.750

Fuel value of the food . . . . 2676 calories.

### Thursday, May 19, 1904.

Breakfast. — Banana 98 grams, boiled hominy 150 grams, bread 60 grams, butter 10 grams, coffee 150 grams, cream 125 grams, sugar 45 grams.

Lunch. — Bread 61 grams, butter 19 grams, potato 100 grams, fried hominy 114 grams, syrup 48 grams, boiled onion 139 grams, coffee 150 grams, sugar 21 grams.

Dinner. — Tomato pureé 200 grams, bread 40 grams, fried sweet potato 77 grams, macaroni 100 grams, spinach 100 grams, bacon 21 grams, bread pudding 100 grams, cream 50 grams, sugar 7 grams.

Food.	Grams. I	er cent Nitrogen.	Total Nitrogen.
Banana	98 ×	0.23 =	0.225 grams.
Bread	60 ×	1.54 =	0.924
Butter 10 + 19	= 29 ×	0.15 =	0.044
Sugar $45 + 21 + 7$	= 78 ×	0.00 =	0.000
Coffee $150 + 150$	= 800 ×	0.06 =	0.180
Cream $125 + 50$	= 175 ×	0.47 =	0.828
Hominy	150 🗙	0.20 =	0.800
Bread	61 ×	1.60 =	0.976
Potato	100 ×	0.49 =	0.490
Fried hominy	114 ×	0.67 =	0.764
Syrup	48 ×	0.024 =	0.012
Onion	189 ×	. 0.27 =	0.375
Bread	40 ×	1.74 =	0.696
Tomato purée	200 ×	0.53 =	1.060
Fried sweet potato	77 ×	0.88 =	0.298
Macaroni	100 ×	0.98 =	0.980
Spinach	100 ×	0.56 =	0.560
Bacon	21 ×	3.00 =	0.680
Pudding	100 ×	0.20 =	0.200
Total nitro	gen in food		9.482 grams.
Total nitro	gen in urine .		A A40

Fuel value of the food . . . . 2758 calories.

## Friday, May 20, 1904.

Breakfast. — Orange 70 grams, baked potato 87 grams, roll 59 grams, butter 32 grams, sugar 7 grams.

Lunch. — Bread 67 grams, butter 15 grams, fish cake 88 grams, potato 150 grams, bread pudding 150 grams, cream 50 grams.

Dinner. — Consommé 150 grams, fish 70 grams, string beans 70 grams, potato 155 grams, cranberry sauce 102 grams, bread 82 grams, coffee 100 grams, sugar 14 grams.

Food.		Grams.	Per cent Nitrogen.	Total Nitrogen.
Orange		70	× 0.20 =	= 0.140 grams.
Butter	32 + 15 =	47	× 0.15 =	
Roll		59	× 1.72 =	= 1.015
Sugar	7 + 14 =	21	× 0.00 =	= 0.000
Potato		87	× 0.40 =	= 0.348
Bread		67	× 1.71 =	= 1.146
Fish-cake		88	× 1.22 =	= 1.074
Potato		150	× 0.30 =	= 0.450
Bread pudding		150	× 0.99 =	= 1.485
Cream		50	× 0.44 =	= 0.220
Potato		155	× 0.34 =	= 0.527
Consommé		150	× 0.59 =	= 0.885
Bread		82	× 1.97 =	= 0.630
String beans .		70	× 0.36 =	= 0.252
Cranberry sauce		102	× 0.08 =	= 0.081
Fish		70	× 3.18 =	2.226
Coffee		100	× 0.06 =	= 0.060
	Total nitrogen	in food .	. <b></b> .	. 10.560 grams.
	Total nitrogen	in urine		. 8.450

Fuel value of the food . . . 1911 calories.

## Saturday, May 21, 1904.

Breakfast. — Banana 106 grams, boiled Indian-meal 150 grams, sugar 21 grams, cream 50 grams, bread 59 grams, butter 16 grams.

Lunch. — Bread 55 grams, butter 13 grams, lamb chop 37 grams, potato croquette 105 grams, tomato 216 grams, sugar 14 grams, water ice 148 grams.

Dinner.—Bean soup 100 grams, fried egg 22 grams, bacon 10 grams, lettuce salad 68 grams, fried potato 100 grams, coffee 100 grams, cream 50 grams, sugar 21 grams, stewed prunes 247 grams.

Food.	Grams.	Per o	ent Nitrog	en.	Total Nitrogen.
Bread	59	×	1.65	=	0.974 grams.
Butter 16 + 13 =	29	×	0.15	=	0.044
Banana	106	×	0.23	=	0.244
Boiled Indian-meal	150	×	0.17	=	0.255
Sugar $21 + 14 + 21 =$	56	×	0.00	=	0.000
Cream $50 + 50 =$	100	×	0.48	=	0.430
Bread	55	×	1.82	=	1.001
Potato croquette	105	×	0.71	=	0.746
Lamb chop	87	×	4.63	=	1.718
Tomato	216	×	0.17	=	0.367
Water ice	143	×	0.012	=	0.017
Prunes	247	×	0.16	=	0.895
Bean soup	100	×	1.21	=	1.210
Fried potato	100	×	0.60	=	0.600
Egg	22	×	2.27	=	0.499
Bacon	10	×	3.05	=	0.306
Salad	63	×	0.21	=	0.132
Coffee	100	×	0.06	=	0.060
Total nitrogen	in food				. 8.992 grams.
Total nitrogen					. 8.640

Fuel value of the food . . . . 2294 calories.

## Sunday, May 22, 1904.

- Breakfast. Orange 60 grams, oatmeal 207 grams, roll 46 grams, butter 14 grams, coffee 150 grams, cream 150 grams, sugar 35 grams.
- Lunch. Potato 150 grams, boiled onions 145 grams, macaroni 130 grams, fried rice 138 grams, syrup 48 grams, ice cream 160 grams, cake 26 grams.
- Dinner. Celery soup 150 grams, spinach 100 grams, mashed potato 100 grams, bread 19 grams, coffee 100 grams, cream 50 grams, sugar 7 grams, strawberry short-cake 169 grams.

Food.		Grams.	Per cent Nitrogen	. Total Nitrogen.							
Orange		60	× 0.20 =	0.120 grams.							
Oatmeal		207	× 0.43 =	0.890							
Sugar	35 + 7 =	42	× 0.00 =	0.000							
Cream	150 + 50 =	200	× 0.45 =	. 0.900							
Roll		46	× 1.67 =	0.768							
Coffee	150 + 100 =	250	× 0.06 =	0.150							
Butter		14	× 0.15 =	0.021							
Potato		150	× 0.30 =	0.450							
Onions		145	× 0.25 =	0.863							
Macaroni		180	× 0.46 =	0.598							
Fried rice		138	× 0.75 =	1.085							
Syrup	<i>.</i>	48	× 0.024 =	0.012							
Ice cream		160	× 0.58 =	0.848							
Cake		26	× 1.20 =	0.812}							
Bread		19	× 1.57 =	0.298							
Celery soup .		150	× 0.33 =	0.495							
Spinach		100	× 0.55 =	0.550							
Short-cake		169	× 0.50 =	0.845							
Mashed potato .		100	× 0.87 =	0.370							
	Total nitrogen	in food .		. 9.025 grams.							
	Total nitrogen in urine 8.530										

Fuel value of the food . . . . 2781 calories.

## Monday, May 23, 1904.

- Breakfast. Banana 201 grams, cream 100 grams, sugar 28 grams, griddle cakes 108 grams, syrup 48 grams.
- Lunch. Consommé 150 grams, rice croquette 140 grams, syrup 48 grams, fried potato 100 grams, bread 36 grams, butter 15 grams, apple sauce 90 grams, coffee 75 grams, sugar 7 grams.
- Dinner. Vegetable soup 100 grams, potato croquette 50 grams, string beans 120 grams, macaroni 104 grams, bacon 20 grams, bread 26 grams, water ice 184 grams.

Food.		Grams. Per c	ent Nitrogen.	Total Nitrogen.
Griddle cakes .		108 ×	0.91 =	0.937 grams
Banana		201 ×	0.23 =	0.462
Cream		100 ×	0.45 =	0.450
Sugar	28 + 7 =	85 ×	0.00 =	0.000
Syrup	48 + 48 =	96 ×	0.024 =	0.023
Consommé		150 ×	0.65 =	0.975
Rice croquette .		140 ×	0.61 =	0.854
Butter		15 ×	0.15 =	0.023
Bread		36 ×	1.75 =	0.630
Apple sauce .		90 ×	0.020 =	0.018
Fried potato .		100 ×	0.60 =	0.600
Coffee		75 ×	0.06 =	0.045
Vegetable soup		100 ×	0.70 =	0.700
Bread		26 ×	1.75 =	0.455
Macaroni		104 ×	0.87 =	0.905
String beans .		120 ×	0.22 =	0.264
Water ice		184 ×	= 800.0	0.011
Bacon	. ,	20 ×	8.28 =	0.656
Potato croquette		50 ×	0.77 =	0.385
	Total nitrogen i			8.393 grams.
	Total nitrogen i	n urine		7.690

Fuel value of the food . . . . 2819 calories.

## Tuesday, May 24, 1904.

Breakfast.—Orange 80 grams, fried rice 186 grams, syrup 72 grams, coffee 100 grams, cream 50 grams, sugar 21 grams.

Lunch.—Celery soup 125 grams, bread 34 grams, butter 19 grams, boiled onion 127 grams, potato 150 grams, tomato sauce 50 grams, stewed prunes 189 grams, cream 50 grams.

Dinner. — Tomato soup 125 grams, bread 20.5 grams, fried potato 100 grams, spinach 180 grams, coffee 100 grams, cream 50 grams, sugar 14 grams, cream pie 158 grams.

Evening. — Ginger ale 250 grams.

Food.						Grams.	Pe	r cent Nitro	gen.	Total Nitrogen.
Rice						186	×	0.36	=	0.670 grams.
Syrup						72	×	0.024	=	0.017
Coffee		. 100	+	100 :	=	200	X	0.06	=	0.120
Sugar		. 21	+	14	=	35	×	0.00	=	0.000
Orange						80	×	0.20	=	0.160
Cream	5	+50	+	50	=	150	×	0.45	=	0.675
Bread						84	×	1.66	=	0.564
Butter						19	×	0.15	=	0.029
Celery soup .			•			125	×	0.48	=	0.600
Onion	•					127	×	0.80	=	0.881
Prunes						189	×	0.17	=	0.321
Potato				•		150	×	0.26	=	0.890
Tomato sauce						50	×	0.23	=	0.115
Tomato soup						125	×	0.19	=	0.288
Bread						20.5	×	1.82	=	0.378
Fried potato						100	×	0.46	=	0.460
Spinach						180	X	0.54	=	0.702
Cream pie .			•			158	×	0.93	=	1.469
Ginger ale .				٠.		250	×	0.00	=	0.000
		Total	nit	roge	n i	in food				. 7.284 grams.
		Total	nit	roge	n i	in urine				. 7.840

Fuel value of the food . . . . 2422 calories.

#### NITROGEN BALANCE - Donahus.

		Nitrogen	Output.						
		Taken in	Nitrogen in Urine.	Weight of Fraces* (dry).					
May	18	8.119 grams.	5.75 grams.						
	19	9.482	6.64	15 grams.					
	20	10.560	8.45	• • •					
	21	8.992	8.6 <del>4</del>	• • •					
	22	9.025	8,53	• • •					
	28	8.393	7.69	89					
	24	7.284	7.34	24					
				128 grams contain					
				6.40 % N.					
		61.855	58.04 +	8.192 grams nitrogen.					
		61.855 grams	nitrogen.	61.232 grams nitrogen.					
		Nitrogen balance fo	r seven days =	+0.628 grams.					
		Nitrogen balance pe	er day =	+0.089 grams.					
			Average Intake.						
		Calories per day .		. 2450.					
		Nitrogen per day .		. 8.83 grams.					

<sup>\*</sup> The fæces of the period were separated as customary by the ingestion of lampblack.

Examination of these data shows that the total amount of nitrogen ingested for the seven days was 61.855 grams, while there were eliminated in the urine 53.04 grams and through the fæces 8.192 grams of nitrogen, thus showing a plus balance for the period of 0.623 gram of nitrogen. In other words, with an average daily intake of 8.83 grams of nitrogen and with an average fuel value of the food amounting to only 2450 calories per day, the body was not only kept from loss, but was able to store up a little nitrogen for future needs. Surely, one could not ask for any better demonstration of physiological economy in nutrition than these data, for this seven days' period, afford.

Further, it should be mentioned, as confirmatory of the view that this subject had long been in a condition of nitrogenous equilibrium on about this quantity of food, that the average daily excretion of metabolized nitrogen during this seven days' period was 7.57 grams, while the average daily excretion from April 13 to June 15 was 7.89 grams of nitrogen. Finally, attention may be called to the fact that the ingestion of 8.83 grams of nitrogen corresponds to 55.18 grams of proteid food, while an excretion of 7.57 grams of nitrogen means the metabolism of 47.3 grams of proteid matter. A saving of more than fifty per cent in proteid food and proteid metabolism, with maintenance of body and nitrogen equilibrium with its possible physiological gains is not to be ignored.

With Jacobus, a similar trial for nitrogen balance gave the following results:

#### JACOBUS.

#### Wednesday, May 18, 1904.

- Breakfast. Banana 78 grams, fried rice 100 grams, syrup 58 grams, bread 48 grams, butter 15 grams, coffee 150 grams, cream 118 grams, sugar 85 grams.
- Lunch. Tomato soup 105 grams, sweet potato 61.5 grams, farina croquette 91 grams, syrup 49 grams, bread 35 grams, butter 23 grams, coffee 150 grams, cream 46 grams, sugar 21 grams.
- Dinner. Consommé 155 grams, Hamburg steak 67 grams, spinach 30 grams, potato 150 grams, bread 27 grams, butter 8 grams, pie 110 grams, cream 61 grams, sugar 10 grams.

Food.	Grams.	Per	cent Nitroge	n. 7	lotal Nitrogen.
Butter $15 + 28 + 8 =$	46	×	0.15 =	=	0.069 grams.
Banana	73	×	0.23 =	=	0.168
Fried rice	100	×	0.75 =	=	0.750
Bread	48	×	1.66 =	=	0.797
Cream . $.118 + 46 + 61 =$	225	×	0.46 =	=	1.085
Sugar $35 + 21 + 10 =$	66	×	0.00 =	=	0.000
Coffee $150 + 150 =$	300	×	0.06 =	=	0.180
Syrup $58 + 49 =$	107	×	0.024 =	=	0.026
Tomato soup	105	×	0.41 =	=	0.481
Bread	35	×	1.60 =	=	0.560
Farina croquette	91	×	1.09 =	=	0.991
Sweet potato	61.5	×	0.32 =	=	0.197
Consommé	155	×	0.88 =	=	0.589
Potato	150	×	0.88 =	=	0.570
Hamburg steak	67	×	8.64 =	=	2.439
Bread	27	×	1.80 =	=	0.486
Spinach	80	×	0.53 =	=	0.159
Pie	110	×	0.43 =	=	0.478
Total nitrogen	in food				9.920 grams.
Total nitrogen	in urine				6.750

Fuel value of the food . . . . 2846 calories.

#### JACOBUS.

## Thursday, May 19, 1904.

Breakfast. — Banana 105 grams, baked potato 79 grams, bread 40 grams, butter 15 grams, coffee 225 grams, cream 85 grams, sugar 24 grams.

Lunch. — Omelette (plain) 60 grams, fried hominy 68 grams, syrup 48 grams, potato 100 grams, boiled onion 81 grams, bread 45 grams, butter 14.5 grams, apple pudding 117 grams, sugar 10 grams, cream 40 grams.

Dinner. — Soup 100 grams, bacon 31 grams, baked macaroni 53 grams, fried sweet potato 55 grams, boiled Indian-meal 50 grams, bread 35 grams, butter 15 grams, coffee 155 grams, cream 96 grams, sugar 31 grams, chocolate 40 grams.

Food.		Grams.	Per cent	Nitrogen.	Total Nitrogen.
Banana		105	× 0.5	23 =	0.242 grams
Bread		40	× 1.5	54 =	0.616
Cream	85 + 40 + 96 =	221	× 0.	47 =	1.038
Coffee	225 + 155 =	380	× 0.0	06 ==	0.228
Sugar	10 + 24 + 31 =	65	× 0.9	00 =	0.000
Potato		79	× 0.	<b>1</b> 9 =	0.387
Butter 15	5 + 14.5 + 15 =	44.5	× 0.3	15 =	0.067
Bread		45	× 1.0	30 =	0.720
Onion		81	× 0.5	27 =	0.219
Fried hominy .		68	× 0.0	37 =	0.456
Eggs (omelette)		60	× 1.	58 =	0.948
Potato		100	× 0.4	19 =	0.490
Syrup		48	× 0.0	)24 =	0.012
Apple pudding		117	× 0.5	28 =	0.828
Soup		100	× 0.8	i3 =	0.530
Bread		35	× 1.3	74 =	0.609
Fried sweet potat	to	55	× 0.8	38 =	0.209
Bacon		81	× 3.0	00 =	0.930
Baked macaroni		53	× 0.9	33 =	0.498
Boiled Indian-me	al	50	× 0.2	<b>20</b> =	0.100
Chocolate		40	× 0.7	<b>'8</b> =	0.292
	Total nitrogen i				8.914 grams. 6.270

Fuel value of the food . . . . 2831 calories.

#### JACOBUS.

#### Friday, May 20, 1904.

- Breakfast.—Orange 70 grams, fried hominy 57 grams, syrup 48 grams, baked potato 118 grams, bread 34 grams, butter 16 grams, coffee 75 grams, cream 33 grams, sugar 10 grams.
- Lunch. Fish-cake 88 grams, fried hominy 61 grams, syrup 32 grams, potato 100 grams, bread 45 grams, butter 22.5 grams, coffee 75 grams, cream 35 grams, sugar 10 grams, bread pudding 81 grams.
- Dinner. Consommé 75 grams, boiled fish 99 grams, potato 132 grams, rice croquette 83 grams, syrup 50 grams, bread 49 grams, butter 19 grams, coffee 150 grams, cream 48 grams, sugar 20 grams.

Evening. — Beer 875 grams.

Food.	Grams.	Per	cent Mitre	ogen.	Total Nitrogen.
Butter $16 + 22.5 + 19$	= 57.5	×	0.15	=	0.086 grams.
Orange	. 70	×	0.20	=	0.140
Coffee $75 + 75 + 150$	= 800	×	0.06	=	0.180
Cream $83 + 85 + 48$	= 116	×	0.44	=	0.510
Sugar $10 + 10 + 20$	<b>= 4</b> 0	×	0.00	=	0.000
Potato	. 118	×	0.40	=	0.452
Fried hominy	. 57	×	0.74	=	0.422
Bread	. 84	×	1.72	=	0.585
Syrup $48 + 82 + 50$	= 130	×	0.024	=	0.081
Potato	. 100	×	0.80	=	0.800
Bread	. 45	×	1.71	=	0.770
Fried hominy	. 61	×	0.57	=	0.348
Fish-cake	. 88	×	1.22	=	1.074
Bread pudding	. 81	×	0.99	=	0.802
Bread	. 49	×	1.97	=	0.965
Fish	. 99	×	8.18	=	3.1 <b>4</b> 8
Potato	. 132	×	0.34	=	0.449
Rice croquette	. 88	×	1.06	=	0.880
Beer	. 875	×	0.069	=	0.259
Consommé	. 75	×	0.59	=	0.448
Total nitroge	n in food				. 11.844 grams.
Total nitroge	n in urine				. 7.290

Fuel value of the food . . . . 2914 calories.

## Saturday, May 21, 1904.

- Breakfast. Banana 72 grams, roll 48 grams, butter 7 grams, boiled Indianmeal 100 grams, cream 90 grams, sugar 17 grams.
- Lunch.—Lamb chop 49 grams, potato 95 grams, tomato 91 grams, beans 45 grams, bread 49 grams, butter 13.5 grams, coffee 150 grams, sugar 28 grams, water ice 153 grams.
- Dinner. Soup 75 grams, bacon 12 grams, fried egg 50 grams, potato 100 grams, bread 40 grams, butter 9 grams, prunes 175 grams, coffee 75 grams.

Food.										Grams.	Pe	er cent Nitr	ogen.	Total Nitrogen.
Butter				7 -	<b>+ 1</b> 8	3.5	<b>i</b> +	- 9 :	=	29.5	×	0.15	=	0.044 grams.
Banana										72	. ×	0.23	=	0.167
Boiled L	ndi	an	m	eal						100	X_	0.17	=	0.170
Cream					. 8	0	+	45	=	135	×	0.43	=	0.581
Sugar.			•		. 1	7	+	28	=	45	×	0.00	=	0.000
Roll .										48	×	1.65	=	0.792
Bread .										49	×	1.82	=	0.892
Lamb ch	юр	٠.								49	×	4.63	=	2.269
Potato										95	×	0.71	=	0.675
Coffee.					15	0	+	75 :	=	225	×	0.06	=	0.135
Tomato				•.						91	×	0.17	=	0.155
Water I	ce									158	×	0.012	=	0.018
Bread.	•				•					40	×	1.62	=	0.648
Soup .										75	, ×	1.21	=	0.908
Prunes										175	'×	0.16	=	0.280
Potato					•					100	×	0.60	=	0.600
Egg .	•				•					50	×	2.27	=	1.135
Bacon.										12	×	3.05	=	0.366
				T	ota	l n	ita	roge	n	in foo	d			. 9.835 grams.
				T	ota	lr	it	roge	en.	in uri:	ne .			. 7.070

Fuel value of the food . . . . 2157 calories.

# Sunday, May 22, 1904.

- Breakfast. Orange 44 grams, catmeal 130 grams, roll 52 grams, coffee 150 grams, cream 120 grams, sugar 30 grams.
- Lunch. Fried rice 72 grams, syrup 48 grams, boiled onions 70 grams, potato 100 grams, bread 33 grams, butter 14.5 grams, coffee 150 grams, cream 45 grams, sugar 20 grams, ice cream 147 grams.
- Dinner. Soup 100 grams, chicken 75 grams, fried potato 50 grams, spinach 15 grams, bread 48 grams, butter 12 grams, coffee 75 grams, cream 50 grams, sugar 14 grams, strawberry short-cake 201 grams.

Food.		Grams.	Per	cent Mitro	gen.	Total Nitrogen.
Orange		44	X	0.20	=	0.088
Oatmeal		180	X	0.43	=	0.559
Roll		52	X	1.67	=	0.868
Cream 45 + 120 -	- <b>50</b> =	215	X	0.45	=	0.968
Sugar 30 + 20 +	<b>⊦ 14 =</b>	64	X	0.00	=	0.000
Coffee 150 + 150 +	<b>- 75</b> =	375	X	0.06	=	0.225
Bread		83	X	1.57	=	0.518
Butter 14.5	<b>- 12 =</b>	26.5	X	0.15	=	0.040
Onions		70	×	0.25	=	0.175
Potato		100	X	0.80	=	0.300
Ice cream		147	×	0.58	=	0.779
Fried rice		72	X	0.75	=	0.540
Syrup		48	×	0.024	=	0.012
Bread		48	×	1.91	=	0.917
Soup		100	X	0.30	=	0.800
Chicken		75	X	3.02	=	2.265
Fried potato		. 50	X	0.37	=	0.185
Spinach		15	X	0.55	=	0.083
Strawberry short-cake .		210	×	0.50	=	1.005
Total ni	trogen	in food				. 9.827 grams
Total ni	trogen	in urine				. 7.620

Fuel value of the food . . . . 2836 calories.

## Monday, May 23, 1904.

Breakfast. — Banana 70 grams, roll 64 grams, butter 9 grams, coffee 75 grams, cream 85 grams, sugar 10 grams.

Lunch. — Boiled eggs 73 grams, fried potato 75 grams, bread 58 grams, butter 11.5 grams, apple sauce 90 grams, coffee 75 grams, cream 35 grams, sugar 21 grams.

Dinner. — Bacon 35 grams, potato croquette 47 grams, bread 39 grams, butter 9.5 grams, chocolate 45 grams, coffee 100 grams, cream 50 grams, sugar 15 grams, water ice 151 grams.

Grams.	Pe	r cent Nitro	gen.	Total Nitrogen.			
70	×	0.28	=	0.161 grams.			
30	×	0.15	=	0.045			
170	×	0.45	=	0.765			
46	×	0.00	=	0.000			
64	×	1.68	=	1.043			
250	×	0.06	=	0.150			
75	×	0.60	=	0.450			
58	×	1.64	=	0.951			
78	×	2.07	=	1.511			
90	×	0.02	=	0.018			
47	×	0.77	=	0.362			
35	×	3.28	=	1.148			
39	×	1.75	=	0.683			
45	×	0.78	=	0.329			
151	×	0.006	=	0.009			
Total nitrogen in food							
	70 30 170 46 64 250 75 58 78 90 47 35 39 45 151	70 × 30 × 170 × 46 × 64 × 250 × 75 × 58 × 78 × 90 × 47 × 35 × 39 × 45 × 151 ×	70 × 0.28 30 × 0.15 170 × 0.45 46 × 0.00 64 × 1.68 250 × 0.06 75 × 0.60 58 × 1.64 78 × 2.07 90 × 0.02 47 × 0.77 35 × 3.28 39 × 1.75 45 × 0.78 151 × 0.006	70 × 0.28 = 30 × 0.15 = 170 × 0.45 = 46 × 0.00 = 64 × 1.68 = 250 × 0.06 = 75 × 0.60 = 58 × 1.64 = 78 × 2.07 = 90 × 0.02 = 47 × 0.77 = 35 × 3.28 = 39 × 1.75 = 45 × 0.78 = 151 × 0.006 = 1 in food			

Fuel value of the food . . . . 2041 calories.

# Tuesday, May 24, 1904.

- Breakfast. Orange 80 grams, boiled rice 105 grams, roll 55 grams, butter 9 grams, coffee 100 grams, cream 50 grams, sugar 30 grams.
- Lunch. Soup 100 grams, fried potato 80 grams, boiled onions 130 grams, bread 44.5 grams, butter 12.5 grams, stewed prunes 108 grams, coffee 100 grams, cream 50 grams, sugar 21 grams.
- Dinner. Hamburg steak 84 grams, mashed potato 135 grams, bread 14 grams, butter 3 grams, cream pie 153 grams.

Food.	Grams.	Per	oent Nitrog	en. To	tal Nitrogen.
Rice	105	×	0.36	=	0.378 grams.
Coffee $100 + 100 =$	200	×	0.06	=	0.120
Orange	80	×	0.20	=	0.160
Roll	55	×	1.64	=	0.902
Butter $9 + 12.5 + 8 =$	24.5	X	0.15	=	0.087
Sugar $80 + 21 =$	51	×	0.00	=	0.000
Cream $50 + 50 =$	100	×	0.45	=	0.450
Bread	44.5	X	1.66	=	0.739
Prunes	108	×	0.17	=	0.184
Soup (celery)	100	×	0.48	=	0.480
Onions	180	×	0.30	=	0.390
Potato (fried)	80	X	0.25	=	0.200
Potato (mashed)	135	×	0.46	=	0.621
Hamburg steak	84	×	3.96	=	8.3 <b>26</b>
Bread	14	×	1.82	=	0.255
Cream pie	153	×	0.98	=	1.423
Total nitroger	in food				9.665 grams.
Total nitroger					7.640

Fuel value of the food . . . . 2174 calories.

### NITROGEN BALANCE. - Jacobus.

	Nitrogen -	Output.	
	Taken in. Nitro	gen in Urine. Weig	ht of Fæces (dry).
<b>May</b> 18	9.920 grams.	6.75 grams.	5.0 grams.
19	8.914	6.27	10.0
20	11.844	7.29	30.0
21	9.885	7.07	28.3
22	9.827	7.62	20.7
28	7.625	6.48	18.8
24	9.665	7.64	32.0
			144.8 grams contain
			6.58 % N.
	67.680	49.12 +	9.528 grams nitrogen.

67.630 grams nitrogen. 58.648 grams nitrogen.

Nitrogen balance for seven days = +8.982 grams. Nitrogen balance per day = +1.283 grams.

#### Average Intake.

Nitrogen per day . . . . . . . . 9.661 grams.

During this balance period of seven days, 67.630 grams of nitrogen were taken in with the food, while 49.12 grams of nitrogen were excreted through the urine and 9.528 grams were passed out through the fæces. This means a large plus balance of 8.98 grams of nitrogen for the entire period, showing that the body was being supplied with considerably more proteid than was necessary for the establishment of nitrogen equilibrium. The average daily intake of nitrogen was 9.661 grams, whereas this might have been reduced to 8.4 grams per day with perfect assurance of nitrogen equilibrium being maintained. Further, it is to be noticed that the average daily intake of food for this period had a fuel value of only 2542 calories. The average daily excretion of metabolized nitrogen during the balance period was only 7.01 grams, while the average daily excretion for the last two months of the experiment amounted to 7.43 grams.

With Schenker the following results were obtained:

### SCHENKER.

# Wednesday, May 18, 1904.

Breakfast. - None.

Lunch. — Bread 53 grams, butter 22 grams, stewed potato 148 grams, string beans 110 grams.

Dinner. — Consommé 200 grams, bread 84 grams, butter 35 grams, Hamburg steak 119 grams, boiled potato 200 grams, spinach 100 grams, apple pie 138 grams.

Food.	•					Grams.	1	Per	COL	at B	litr	oger	١.	Total Ni	trogen.
Bread						53	×		1.6	30		=		0.848	grams.
Stewed potato						148	×		0.5	32		=		0.474	
Butter	22	+	85	. :	=	57	X		0.1	5		=		0.086	
String beans.						110	×.		0.8	34		=		0.874	
Consommé .						200	X		0.8	38		=		0.760	
Bread						84	X		1.8	30		=		1.512	
Hamburg steak						119	X		8.6	34		=		4.332	
Potato						200	X		0.8	38		=		0.760	
Spinach						100	X		0.4	58		=		0.580	
Apple pie	•					138	×		0.4	18		=		0.593	
	Γot	al	nit	ro	gen	in food								10.269	grams.
						in urine								8.770	_

Fuel value of the food . . . . 2006 calories.

# Thursday, May 19, 1904.

Breakfast. — None.

Lunch. — Bread 82 grams, butter 32 grams, potato 232 grams, omelette 60 grams, apple-tapioca 180 grams.

Dinner. — Tomato soup 200 grams, bread 57 grams, butter 15 grams, macaroni 107 grams, fried sweet potato 100 grams, bacon 28 grams.

Food.			Grams.	Per	ent Nitrogen.	Total Nitrogen.
Bread			82	X	1.60 =	1.312 grams.
Butter		32 + 15 =	47	×	0.15 =	0.071
Potato			232	×	0.49 =	1.137
Omelette .			60	×	1.58 =	0.948
Apple-tapioca			180	×	0.28 =	0.504
Bread			57	×	1.74 =	0.992
Tomato soup			200	×	0.53 =	1.060
Macaroni .			107	×	0.93 =	0.995
Bacon			28	×	8.00 =	0.840
Fried sweet pe	tato		100	X	0.88 =	0.880
		l'otal nitrogen l'otal nitrogen				_

Fuel value of the food . . . . 1900 calories.

# Friday, May 20, 1904.

- Breakfast. Orange 70 grams, baked potato 142 grams, roll 84 grams, butter 32.5 grams, coffee 150 grams, cream 50 grams, sugar 21 grams.
- Lunch. Fish-ball 85 grams, potato 175 grams, lima beans 60 grams, bread 57 grams, butter 17 grams, bread pudding 336 grams.
- Dinner. Consommé 150 grams, halibut 110 grams, potato 186 grams, string beans 50 grams, bread 72 grams, butter 18 grams, cranberry sauce 163 grams, sugar 19 grams.

Food.	Grama.	Per cent Nitrogen.	Total Mitrogen.
Orange	70	× 0.20 =	0.140 grams.
Butter 32	2.5 + 17 + 18 = 67.5	× 0.15 =	0.101
Roll	84	× 1.72 =	1.445
Coffee	150	× 0.06 =	0.090
Cream	50	× 0.44 =	0.220
Sugar	21 + 19 = 40	× 0.00 =	0.000
Potato	142	× 0.40 =	0.568
Potato	175	× 0.30 =	0.525
Fish-ball	85	× 1.22 =	1.037
Bread	57	× 1.71 =	0.975
Bread pudding	836	× 0.99 =	8.826
Lima beans .	60	× 0.76 =	0.456
Bread	72	× 1.97 =	1. <del>4</del> 18
Potato	186	× 0.34 =	0.632
Fish (halibut)	110	× 8.18 =	3.498
String beans .	50	× 0.86 =	0.180
Cranberry sauce	163	× 0.08 =	0.049
Consommé	150	× 0.59 =	0.885
	Total nitrogen in food		15.545 grams.
	Total nitrogen in urine		12.480

Fuel value of the food . . . . 2798 calories.

## Saturday, May 21, 1904.

- Breakfast. Banana 184 grams, cream 80 grams, boiled Indian-meal 155 grams, baked potato 140.5 grams, butter 15 grams, sugar 7 grams.
- Lunch. Lamb chop 25 grams, tomato 148 grams, potato croquette 147 grams, fried Indian-meal 47.5 grams, syrup 48 grams, bread 35 grams, water ice 162 grams.
- Dinner.—Bean soup 150 grams, bread 25 grams, butter 19 grams, bacon 29 grams, fried potato 150 grams, orange salad 67 grams, stewed prunes 208 grams, cream 50 grams.

Food.	Grams.	Per cent	Nitrogen.	Total Nitrogen.
Boiled Indian-meal	155	× 0.	17 =	0.264 grams.
Banana	184	× 0.	23 =	0.423
Cream $80 + 50 =$	130	× 0.	43 =	0.559
Baked potato	140.5	× 0.	40 =	0.562
Sugar	7	·× 0.	00 =	0.000
Butter $15 + 19 =$	34	× 0,	15 =	0.051
Lamb chop	25	× 4.	68 =	1.158
Tomato	148	× 0.	17 =	0.252
Potato croquette	147	× 0.	71 =	1.044
Bread	85	× 1.	82 =	0.637
Fried Indian-meal	47.5	× 1.	09 =	0.518
Syrup	48	× 0.	024 =	0.012
Water ice	162	× 0.	012 =	0.019
Bread	25	× 1.	62 =	0.405
Orange salad	67	× 0.	21 =	0.141
Stewed prunes	208	× 0.	16 =	0.382
Fried potato	150	× 0.	60 =	0.900
Bacon	29	× 3.	05 =	0.885
Bean soup	150	× 1.	21 =	1.815
Total nitrogen	in food			9.977 grams.
Total nitrogen				8.760

Fuel value of the food . . . . 2661 calories.

## Sunday, May 22, 1904.

- Breakfast. Orange 126 grams, baked potato 169 grams, roll 48 grams, butter 15 grams.
- Lunch.—Bread 53 grams, butter 15.5 grams, macaroni 165 grams, potato 150 grams, fried rice 114 grams, syrup 48 grams, ice cream 148 grams, cake 45 grams.
- Dinner. Celery soup 150 grams, fried potato 50 grams, spinach 40 grams, mashed potato 50 grams, chicken 85 grams, strawberry short-cake 218 grams.

Food.	Grams.	Per cent Nitrogen.	Total Nitrogen.
Orange	126	× 0.20 =	0.252 grams.
Roll	43	× 1.67 =	0.718
Butter	15 + 15.5 = 30.5	× 0.15 =	0.046
Potato	169	× 0.40 =	0.676
Bread	58	× 1.57 =	0.832
Macaroni	165	× 0.46 =	0.759
Potato	150	× 0.80 =	0.450
Cake	45	× 1.20 =	0.540
Ice cream	148	× 0.53 =	0.784
Fried rice	114	× 0.75 =	0.855
Syrup	48	× 0.024 =	0.012
Fried potato .	50	× 0.57 =	0.285
Spinach	40	× 0.55 =	0.220
Chicken	85	× 3.02 =	2.567
Strawberry shor	t-cake 218	× 0.50 =	1.065
Celery soup .	150	× 0.83 =	0.495
Mashed potato	50	× 0.87 =	0.185
, -	Total nitrogen in food		10.741 grams.
	Total nitrogen in urine		9.980

Fuel value of the food . . . . 2788 calories.

# Monday, May 23, 1904.

- Breakfast. Banana 225 grams, griddle cakes 127 grams, syrup 96 grams, roll 62 grams, butter 9.5 grams.
- Lunch.—Consommé 100 grams, scrambled eggs 82 grams, fried potato 150 grams, rice croquette 150 grams, syrup 72 grams, bread 24 grams, butter 15 grams, apple sauce 125 grams.
- Dinner. Vegetable soup 100 grams, bread 51 grams, butter 17 grams, bacon 59 grams, potato croquette 80 grams, macaroni 100 grams, water ice 184 grams.

Food.	Grams.	Per	cent Nitr	ogen.	Total Nitrogen
Banana	. 225	X	0.28	=	0.518 grams.
Griddle cakes	. 127	X	0.91	=	1.156
Butter $9.5 + 15 + 17$	= 41.5	×	0.15	=	0.062
Roll	. 62	×	1.63	=	1.011
Syrup 96 + 72 :	= 168	X	0.024	=	0.040
Scrambled eggs	. 82	X	2.07	=	1.697
Fried potato	. 150	X	0.60	=	0.900
Rice croquette	. 150	X	0.61	=	0.915
Apple sauce	. 125	X	0.020	=	0.025
Consommé	. 100	X	0.65	=	0.650
Bread	. 24	X	1.64	=	0.394
Bread	. 51	X	1.75	=	0.893
Bacon	. 59	X	3.28	=	1.935
Potato croquette	. 80	X	0.77	=	0.610
Macaroni	. 100	X	0.87	=	0.870
Water ice	. 184	X	0.006	=	0.011
Vegetable soup	. 100	X	0.48	=	0.480
Total nitroge	n in food				12.167 grams.
Total nitroge					10.040

Fuel value of the food . . . . 3100 calories.

. ....

# Tuesday, May 24, 1904.

### Breakfast. - None.

Lunch. — Cream of celery soup 100 grams, bread 35 grams, butter 10 grams, farina croquette 88 grams, potato 150 grams, stewed prunes 176 grams.
 Dinner. — Tomato soup 150 grams, Hamburg steak 77 grams, potato 150 grams, spinach 75 grams, farina croquette 107 grams, syrup 48 grams, bread 30.5 grams, butter 7.5 grams, cream pie 162 grams.

Food.		Grams.	Per cent Nitrog	en.	Total Nitrogen.
Bread		85	× 1.64	=	0.574 grams.
Butter	. 10 + 7.5 =	17.5	× 0.15	=	0.026
Farina croquette		88	× 0.74	=	0.651
Potato		150	× 0.26	=	0.890
Prunes		176	× 0.17	=	0.299
Cream of celery	soup	100	× 0.48	=	0.480
Tomato soup .		150	× 0.19	=	0.285
Hamburg steak		77	× 3.96	=	8.049
Potato (fried) .		150	× 0.46	=	0.690
Spinach		75	× 0.54	=	0.405
Bread		80.5	× 1.82	=	0.555
Cream pie		162	× 0.98	=	1.507
Farina croquette		107	× 0.76	=	0.813
Syrup		48	× 0.024	=	0.012
	Total nitrogen	in food			9.736 grams.
	Total nitrogen	in urine			8.710

Fuel value of the food . . . . 2151 calories.

### NITROGEN BALANCE. - Schenker.

	Nitrogen	Output.					
	Taken in.	Nitrogen in U	rine. Weight of Faces (dry).				
May 18	10.269 grams.	, 8.77 gran	ne, 8.0 grams.				
19	8.239	8.73	41.0				
20	15.545	12.48	<b>26</b> .5				
21	9.977	8.76	48.4				
22	10.741	9.98	29.0				
23	12.167	10.04	14.7				
24	9.736	8.71	20.5				
			11.0				
			198.4 grams contain				
			6.50 % N.				
•	76.674	67.47	+ 12.571 grams nitrogen.				
	76.674 grams n	itrogen. 80	0.041 grams nitrogen.				
Nitro	gen balance for seve	en days =	-3.867 grams.				
Nitro	gen balance per day	=	-0.481 gram.				
	Ave	rage Intake.					
Calor	ies per day		2486.				

. . . . . . 10.95 grams.

In this seven days' balance trial, there were taken in with the food 76.674 grams of nitrogen, with an output of 67.47 grams of nitrogen through the urine and 12.571 grams through the fæces, thus showing a minus balance of 3.367 grams of nitrogen for the seven days. The fuel value of the food averaged 2486 calories per day, while the average daily excretion of metabolized nitrogen amounted to 9.63 grams. Schenker's daily nitrogen excretion for the last two months of the experiment averaged 9.82 grams. Presumably, the slight minus

Nitrogen per day .

balance of nitrogen was due to the relatively small fuel value of the food, which doubtless was not quite sufficient for the body-weight, and the degree of bodily activity then prevailing.

With G. W. Anderson, a plus balance was obtained as follows:

### G. W. ANDERSON.

# Wednesday May 18, 1904.

Breakfast. — Banana 103 grams, coffee 150 grams, sugar 42 grams, cream 125 grams, fried rice 45 grams, baked potato 92 grams, roll 65 grams, butter 11 grams.

Lunch. — Soup 150 grams, farina croquette 88 grams, sweet potato 206 grams, string beans 75 grams, bread 62 grams, butter 15.5 grams, coffee 150 grams, sugar 14 grams, cream 30 grams.

Dinner. — Consommé 200 grams, bread 59 grams, butter 20 grams, Hamburg steak 109 grams, mashed potato 150 grams, coffee 200 grams, cream 30 grams, sugar 24 grams, spinach 100 grams, apple pie 150 grams.

Food.	Grams.	Po	er cent Nitroge	. Total Nitrogen.
Butter . $.11 + 15.5 + 20 =$	46.5	×	0.15 =	= 0.070 grams.
Sugar . $.42 + 24 + 14 =$	80.0	×	0.00 =	= 0.000
Banana	103.0	X	0.23 =	= 0.237
Cream . $125 + 30 + 80 =$	185.0	×	0.46 =	= 0.851
Fried rice	<b>4</b> 5.0	×	0.75 =	= 0.388
Roll	<b>65.</b> 0	×	1.66 =	1.079
Potato	92.0	×	0.89 =	· 0.859
Coffee $.150 + 150 + 200 =$	500.0	×	0.06 =	: 0.800
Bread	62.0	×	1.60 =	. 0.992
Soup	150.0	×	0.41 =	: 0.615
Farina croquette	88.0	×	1.09 =	= 0.959
Sweet potato	206.0	×	0.82 =	: 0.659
String beans	75.0	×	0.34 =	= 0. <b>255</b>
Bread	59.0	×	1.80 =	1.062
Soup (consommé)	200.0	×	0.88 =	= 0.760
Hamburg steak	109.0	×	3.64 =	<b>3.968</b>
Potato	150.0	×	0.38 =	= 0.570
Spinach	100.0	X	0.53 =	= 0.580
Apple pie	150.0	×	0.48 =	= 0.645
Total nitroger	in food			. 14.249 grams.
Total nitroger	in urine			. 8.870

Fuel value of the food . . . . 3323 calories.

### Thursday, May 19, 1904.

- Breakfast. Banana 170 grams, cream 140 grams, sugar 28 grams, coffee 150 grams, baked potato 127 grams, rolls 47 grams, butter 15 grams.
- Lunch. Bread 50 grams, butter 18 grams, fried potato 150 grams, fried hominy 100 grams, syrup 48 grams, coffee 150 grams, cream 50 grams, sugar 14 grams.
- Dinner. Soup 200 grams, fried sweet potato 70 grams, macaroni 125 grams, spinach 105 grams, bacon 18 grams, butter 15 grams, coffee 150 grams, sugar 14 grams, cream 40 grams.

Food.	Grams.	Per	cent Nitrogen.	Total Nitrogen.
Banana	. 170	×	0.23 =	0.391 grams.
Cream . $.140 + 50 + 40$	= 280	×	0.46 =	1.068
Sugar $28 + 14 + 14$	= 56	×	0.00 =	0.000
Butter $15 + 18 + 15$	= 48	×	0.15 =	0.072
Rolls	. 47	×	1.66 =	0.780
Potato	. 127	×	0.39 =	0.495
Coffee $. 150 + 150 + 150$	= 450	×	0.06 =	0.270
Bread	. 50	×	1.60 =	0.800
Fried potato	. 150	×	0.32 =	0.480
Syrup	. 48	×	0.024 =	0.012
Hominy	. 100	×	0.67 =	0.670
Bread	. 46	×	1.74 =	0.800
Soup	. 200	×	0.53 =	1.060
Fried sweet potato	. 70	×	0.38 =	0.266
Macaroni	. 125	×	0.98 =	1.163
Spinach	. 105	×	0.56 =	0.588
Bacon	. 13	×	3.00 =	0.390
Total nitrog	en in food			. 9.295 grams.
Total nitrog	en in urine			. 8.860

Fuel value of the food . . . . 2932 calories.

### Friday, May 20, 1904.

Breakfast.—Orange 70 grams, roll 120 grams, butter 33 grams, coffee 150 grams, sugar 28 grams, cream 50 grams.

Lunch. — Bread 48 grams, butter 19 grams, fish-cake 81 grams, potato 200 grams, lima beans 50 grams, fried hominy 118 grams, syrup 96 grams, coffee 150 grams, cream 50 grams, sugar 14 grams.

Dinner. — Consommé 100 grams, boiled halibut 148 grams, mashed potato 177 grams, string beans 90 grams, stewed cranberry 76 grams, bread 56 grams, butter 16 grams, coffee 150 grams, cream 50 grams, sugar 14 grams.

					•
Food.	Grams.	Per o	ant Nitrog	en.	Total Nitrogen.
Butter 88 + 19 + 16	= 68	×	0.15	=	0.102 grams.
Orange	. 70	X	0.20	=	0.140
Rolls	. 120	×	1.72	=	2.064
Coffee $. 150 + 150 + 150$	= 450	×	0.06	=	0.270
Cream . $50 + 50 + 50$	= 150	X	0.44	=	0.660
Bread	. <b>4</b> 8	×	1.71	=	0.821
Fish-cake	. 81	X	1.22	=	0.988
Potato	. 200	X	0.30	=	0.600
Lima beans	. 50	X	0.76	=	0.880
Fried hominy	. 118	X	0.57	=	0.678
Syrup	. 96	×	0.024	=	0.023
Bread'	. 56	X	1.97	=	1.104
Consommé	. 100	X	0.59	=	0.590
Halibut	. 143	X	3.18	=	4.547
Mashed potato	. 177	×	0.84	=	0.602
Sugar $28 + 14 + 14$	= 56	X	0.00	=	0.000
String beans	. 90	×	0.36	=	0.824
Cranberry	. 76	×	0.08	=	0.023
Total nitrog	en in food				13.911 grams.
Total nitrog	•				9.960
Town merof	will	- •		• •	

Fuel value of the food . . . . 8052 calories.

# Saturday, May 21, 1904.

Breakfast. — Banana 157 grams, roll 61 grams, butter 15 grams, coffee 150 grams, cream 150 grams, sugar 28 grams.

Lunch. — Bread 58 grams, butter 17 grams, lamb chop 38 grams, potato croquettes 138 grams, tomato 161 grams, water ice 162 grams, coffee 150 grams, cream 50 grams, sugar 21 grams.

Dinner. — Soup 75 grams, bread 59 grams, butter 22.5 grams, bacon 15 grams, lettuce salad 64 grams, fried potato 100 grams, stewed prunes 283 grams, coffee 150 grams, cream 50 grams, sugar 14 grams.

Food.	Grams.	Per c	ent Nitrog	en.	Total Nitrogen.
Banana	157.0	×	0.23	=	0.361 grams.
Roll	61.0	X	1.65	=	1.007
Cream 150 + 50 + 50 =	250.0	X	0.43	=	1.075
Butter $15 + 17 + 22.5 =$	54.5	X	0.15	=	0.082
Sugar $28 + 21 + 14 =$	63.0	×	0.00	=	0.000
Coffee $150 + 150 + 150 =$	450.0	X	0.06	=	0.270
Bread	58.0	×	1.82	=	1.056
Lamb chop	88.0	×	4.68	=	1.528
Potato croquettes	138.0	×	0.71	=	0.980
Tomato	161.0	X	0.17	=	0.274
Water ice	162.0	×	0.012	=	0.019
Bread	59.0	X	1.62	=	0.956
Soup	75.0	×	1.21	=	0.908
Bacon	15.0	X	8.05	=	0.458
Prunes $175 + 108 =$	283.0	X	0.16	=	0.453
Salad	64.0	X	0.21	=	0.134
Fried potato	100.0	×	0.60	=	0.600
Total nitrogen	in food				10.161 grams.
Total nitrogen		• •		•	8.510

Fuel value of the food . . . . 2826 calories.

# Sunday, May 22, 1904.

Breakfast. — Orange 80 grams, oatmeal 100 grams, potato 103 grams, butter 16 grams, roll 50 grams, coffee 150 grams, cream 125 grams, sugar 35 grams.

Lunch.—Bread 47 grams, butter 14.5 grams, macaroni 116 grams, potato 150 grams, fried rice 85 grams, syrup 48 grams, ice cream 169 grams, coffee 150 grams, cream 50 grams, sugar 14 grams.

Dinner. — Soup 100 grams, bread 40.5 grams, butter 16 grams, fried potato 50 grams, mashed potato 100 grams, coffee 150 grams, cream 100 grams, sugar 21 grams, strawberry short-cake 214 grams.

Food.	Grams.	Per o	nt Nitroge	. Total Nitrogen.
Orange	. 80	×	0.20 =	: 0.160 grams.
Butter $16 + 14.5 + 16$	= 46.5	×	0.15 =	= 0.070
Oatmeal	. 100	×	0.43 =	= 0.480
Sugar. $.85 + 14 + 14 + 7$ :	= 70	×	0.00 =	= 0.000
Potato	. 103	×	0.40 =	= 0.412
Cream . $125 + 50 + 100$ :	= 275	×	0.45 =	= 1.288
Roll	. 50	×	1.67 =	- 0.835
Coffee . $150 + 150 + 150$ :	= 450	×	0.06 =	= 0.270
Bread	. 47	×	1.57 =	= 0.788
Macaroni	. 116	×	0.46 =	= 0.534
Potato	. 150	×	0.80 =	= 0.450
Fried rice	. 85	×	0.75 =	= 0.638
Syrup	. 48	×	0.024 =	= 0.012
Ice cream	. 169	×	0.53 =	= 0.896
Bread	. 40.5	×	1.91 =	= 0.774
Celery soup	. 100	×	0.83 =	= 0.330
Fried potato	. 50	×	0.57 =	= 0.285
Mashed potato	. 100	×	0.87 =	= 0.370
Strawberry short-cake	. 214	×	0.50 =	= 1.070
Total nitroge	n in food			9.512 grams.
Total nitroge				6.500
TOWE INTO BE				•

Fuel value of the food . . . . 8429 calories.

# Monday, May 23, 1904.

- Breakfast. Banana 211 grams, roll 59 grams, butter 15 grams, coffee 150 grams, cream 150 grams, sugar 28 grams.
- Lunch. Consommé 100 grams, bread 63 grams, butter 15 grams, rice 113 grams, syrup 48 grams, sugar 14 grams, fried potato 170 grams, apple sauce 125 grams.
- Dinner. Vegetable soup 100 grams, bread 70 grams, butter 16 grams, bacon 42 grams, potato croquettes 49 grams, macaroni 105 grams, string beans 184 grams, water ice 148 grams.

Food.	Grams.	Per cent N	itrogen.	Total Nitrogen.
Banana	211	× 0.28	=	0.485 grams.
Butter $15 + 15 + 16 =$	46	× 0.15	=	0.069
Cream	150	× 0.45	=	0.675
Roll	<b>59</b>	× 1.63	=	0.962
Sugar $28 + 14 =$	42	× 0.00	=	0.000
Coffee	150	× 0.06	=	0.090
Bread	63	× 1.64	=	1.033
Consommé	100	× 0.65	=	0.650
Apple sauce	125	× 0.02	=	0.025
Fried rice	113	× 0.61	=	0.689
Syrup	48	× 0.02	4 =	0.012
Fried potato	170	× 0.60	=	1.020
Bread	70	× 1.75	=	1.225
Vegetable soup	100	× 0.70	=	0.700
Macaroni	105	× 0.87	<b>'</b> =	0.914
Bacon	42	× 8.28	=	1.378
Potato croquettes	149	× 0.77	=	1.147
String beans	184	× 0.22	=	0.405
Water ice	148	× 0.00	6 =	0.009
Total nitrogen	in food			11.488 grams.
Total nitrogen	in urine			6.900

Fuel value of the food . . . 8057 calories.

### Tuesday, May 24, 1904.

- Breakfast. Orange 80 grams, roll 55 grams, butter 16 grams, potato 91 grams, coffee 150 grams, cream 50 grams, sugar 21 grams.
- Lunch. Celery soup 150 grams, bread 62 grams, butter 19 grams, mashed potato 200 grams, farina croquettes 87 grams, syrup 48 grams, stewed prunes 138 grams.
- Dinner. Soup 150 grams, bread 43 grams, butter 15 grams, Hamburg steak 82 grams, fried potato 150 grams, spinach 85 grams, rice croquettes 57 grams, syrup 48 grams, coffee 150 grams, cream 50 grams, sugar 14 grams, lemon pie 125 grams.

Food.	•	Grams.	Per cent Nitrog	gen. Total Nitrogen.
Butter	16 + 19 + 15 =	50	× 0.15	= 0.075 grams.
Orange		80	× 0.20	= 0.160
Roll		55	× 1.64	= 0.902
Potato		91	× 0.25	= 0.228
Coffee	150 + 150 =	800	× 0.06	= 0.180
Cream	50 + 50 =	100	× 0.45	= 0.450
Sugar	21 + 14 =	85	× 0.00	= 0.000
Bread		62	× 1.66	= 1.029
Celery soup		150	× 0.48	= 0.720
Mashed potato.		200	× 0.26	= 0.520
Farina croquette		87	× 0.74	= 0.644
Syrup	48 + 48 =	96	× 0.024	= 0.028
Prunes		188	× 0.17	= 0.285
Soup		150	× 0.19	= 0.285
Bread		43	× 1.82	= 0.783
Hamburg steak		82	× 3.96	= 3.247
Fried potato .		150	× 0.46	= 0.690
Spinach		85	× 0.54	= 0.459
Rice croquettes		57	× 0.76	= 0.438
Lemon pie		125	× 0.98	= 1.163
	Total nitrogen	in food		12.226 grams
	Total nitrogen	in urine		8.450

Fuel value of the food . . . . 3022 calories.

### NITROGEN BALANCE. - G. W. Anderson.

	Nitrogen	Output.				
	Taken in.	Nitrogen in Urine.	Weight of Faces (dry).			
May 18	14.249 grams.	8.87 grams.	• • •			
19	9.295	8. <b>36</b>	••			
20	13.911	9.95	12.0 grams.			
21	10.161	8.51	46.0			
22	9.512	6.50	89.0			
23	11.488	6.90	67.0			
24	12.226	8. <b>4</b> 5				
			164.0 grams contain			
	,		6.92 % N.			
	80.842	<b>57.54</b> +	11.349 grams nitrogen.			
	80.842 grams r	nitrogen. 68	3.889 grams nitrogen.			

Nitrogen balance for seven days = + 11.953 grams. Nitrogen balance per day = + 1.707 grams.

### Average Intake.

With G. W. Anderson, the balance trial was characterized by an intake of 80.842 grams of nitrogen, with an output for the seven days of 57.54 grams through the urine and 11.349 grams through the fæces, thus making a total excretion of 68.889 grams of nitrogen, and showing a plus balance of 11.953 grams. In other words, the body of this subject, under the conditions prevailing, was storing up nitrogen for future use at the rate of 1.7 grams per day. This also means that a daily intake of 9.8 grams of nitrogen would have been quite sufficient to maintain nitrogen equilibrium, certainly with the large fuel value of the food taken, i. e., 3091 calories per day as the average value.

The average daily excretion of metabolized nitrogen during the balance period amounted to 8.22 grams, while the average daily excretion for the last two months of the experiment was 8.81 grams.

With Stapleton, the following results were obtained, showing a distinct positive balance:

### STAPLETON.

# Wednesday, May 18, 1904.

- Breakfast. Banana 118 grams, bread 29 grams, butter 11 grams, sugar 56 grams, cream 125 grams.
- Lunch. Tomato soup 247 grams, bread 37 grams, butter 9 grams, croquettes 97 grams, potato 100 grams, string beans 46 grams, coffee 150 grams, cream 50 grams, sugar 21 grams.
- Dinner. Bread 109 grams, butter 40 grams, Hamburg steak 87 grams, potato 150 grams, spinach 100 grams, coffee 150 grams, cream 100 grams, sugar 30 grams, apple pie 110 grams.

Grams.	Per	cent Nitr	ogen.	Total Nitrogen.
118	×	0.23	=	0.271 grams.
60	×	0.15	=	0.090
29	×	1.66	=	0.481
86	×	0.00	=	0.000
275	×	0.46	=	1.265
247	×	0.41	=	1.013
87	×	1.60	=	0.592
97	×	1.09	=	1.057
100	×	0.82	=	0.320
46	×	0.84	=	0.156
800	×	0.06	=	0.180
109	×	1.80	=	1.962
87	×	3.64	=	3.167
150	×	0.38	=	0.570
100	×	0.53	=	0.530
110	×	0.43	=	0.473
in food				12.127 grams.
•				9.670
	118 60 29 86 275 247 87 97 100 46 800 109 87 150 110 in food	118 × 60 × 29 × 86 × 275 × 247 × 87 × 100 × 46 × 800 × 109 × 150 × 150 ×	118 × 0.23 60 × 0.15 29 × 1.66 86 × 0.00 275 × 0.46 247 × 0.41 87 × 1.60 97 × 1.09 100 × 0.82 46 × 0.84 800 × 0.06 109 × 1.80 87 × 3.64 150 × 0.38 100 × 0.53 110 × 0.43 in food · · · ·	118

Fuel value of the food . . . . 3109 calories.

# Thursday, May 19, 1904.

Breakfast. - None.

Lunch. — Bread 48 grams, butter 14 grams, omelette 125 grams, boiled onion 63 grams, fried sweet potato 100 grams, coffee 300 grams, cream 150 grams, sugar 56 grams, apple pudding 146 grams.

Dinner. — Tomato soup 200 grams, bread 42 grams, butter 9 grams, macaroni 75 grams, potato 36 grams, spinach 70 grams, bacon 16 grams, coffee 150 grams, cream 50 grams, sugar 21 grams.

Food.	Grams.	Per cent Nitrogen.	Total Nitrogen.
Coffee 800 + 150 =	= 450	× 0.06 =	0.270 grams.
Butter 14+9 :	= 28	× 0.15 =	0.035
Bread	. 48	× 1.60 =	0.768
Omelette	. 125	× 1.58 =	1.975
Onions	. 63	× 0.27 =	0.170
Sugar 56 + 21 =	= 77	× 0.00 =	0.000
Cream 150 + 50 :	= 200	× 0.47 =	0.940
Potato	. 100	× 0.49 =	0.490
Apple pudding	. 146	× 0.28 =	0.409
Bread	. 42	× 1.74 =	0.781
Tomato soup	. 200	× 0.58 =	1.060
Macaroni	. 75	× 0.98 =	0.698
Fried sweet potato	. 36	× 0.38 =	0.187
Spinach	. 70	× 0.56 =	0.392
Bacon	. 16	× 3.00 =	0.480
Total nitroge	n in food		. 8.555 grams.
Total nitroge			. 8.580

Fuel value of the food . . . . 2072 calories.

# Friday, May 20, 1904.

- Breakfast. Orange 150 grams, roll 65 grams, coffee 150 grams, cream 50 grams, sugar 35 grams.
- Lunch. Bread 64 grams, butter 18 grams, fish-cake 72 grams, potato 150 grams, lima beans 50 grams, coffee 150 grams, cream 100 grams, sugar 21 grams, bread pudding 150 grams.
- Dinner. Fish 113 grams, string beans 62 grams, potato 150 grams, rice croquettes 102 grams, syrup 48 grams, stewed cranberry 95 grams, bread 33 grams, butter 16 grams, coffee 300 grams, cream 100 grams, sugar 42 grams.

Evening. — Beer 750 grams.

Food.	Grams.	Per	cent Nitro	gen.	Total Nitrogen.
Orange	150	X	0.20	=	0.800 grams.
Sugar $35 + 21 + 42 =$	98	X	0.00	=	0.000
Roll	65	×	1.72	=	1.118
Coffee $. 150 + 150 + 800 =$	600	X	0.06	=	0.360
Cream . $50 + 100 + 100 =$	250	X	0.44	=	1.100
Bread	64	×	1.71	=	1.094
Butter $18 + 16 =$	34	X	0.15	=	0.051
Fish-cake	72	X	1.22	=	0.878
Potato	150	×	0.30	=	0.450
Lima beans	50	×	0.76	=	0.880
Bread pudding	150	X	0.99	=	1.485
String beans	62	X	0.36	=	0.223
Fish	113	X	3.18	=	8.593
Bread	83	X	1.97	=	0.650
Potato	150	X	0.84	=	0.510
Rice croquettes	102	×	1.06	=	1.081
Cranberry	95	×	0.030	=	0.029
Syrup	<b>4</b> 8	×	0.024	=	0.012
Beer	750	×	0.069	=	0.518
Total Nitrogen	in food				18.882 grams.
Total Nitrogen	in urine	٠			9.510

Fuel value of the food . . . . 2999 calories.

### Saturday, May 21, 1904.

- Breakfast. Banana 74 grams, baked potato 95 grams, roll 71 grams, butter 18 grams, coffee 150 grams, cream 100 grams, sugar 30 grams.
- Lunch. Bread 48 grams, butter 14 grams, lamb chop 27 grams, potato croquette 91 grams, tomato 105 grams, coffee 150 grams, cream 50 grams, sugar 21 grams, water ice 185 grams.
- Dinner. Bean soup 150 grams, fried egg 127 grams, bacon 2 grams, fried potato 108 grams, bread 77 grams, butter 18 grams, coffee 150 grams, cream 50 grams, sugar 21 grams, jelly roll 56 grams.
- Evening. Beer 600 grams.

Food.	Grams.	Per	cent Nitr	ogen.	Total Nitrogen.
Banana	74	×	0.23	=	0.170 grams.
Roll	71	×	1.65	=	1.172
Coffee $. 150 + 150 + 150 =$	450	×	0.06	=	0.270
Sugar $30 + 21 + 21 =$	72	×	0.00	=	0.000
Cream . $100 + 50 + 50 =$	200	X	0.43	=	0.860
Potato	95	X	0.40	=	0.380
Butter $18 + 14 + 18 =$	50	×	0.15	=	0.075
Bread	48	×	1.82	=	0.874
Lamb chop	27	×	4.63	=	1.250
Croquette (potato)	91	×	0.71	=	0.646
Tomato	105	×	0.17	=	0.179
Water ice	185	×	0.012	=	0.022
Bread	77	×	1.62	=	1.247
Bean soup	150	×	1.21	=	1.815
Fried egg	127	×	2.27	=	2.883
Bacon	2	×	3.05	=	0.061
Fried potato	108	×	0.60	=	0.648
Jelly roll	56	×	0.86	=	0.482
Beer	600	X	0.069	=	0.414
Total Nitrogen	in food				13.448 grams.
Total Nitrogen	in urine	э			9.640

Fuel value of the food . . . . 2871 calories.

### Sunday, May 22, 1904.

- Breakfast. Orange 60 grams, oatmeal 150 grams, wheat roll 51 grams, butter 17 grams, coffee 150 grams, cream 100 grams, sugar 49 grams.
- Lunch. Bread 37 grams, butter 13 grams, potato 114 grams, macaroni 115 grams, fried rice 92 grams, coffee 300 grams, cream 100 grams, sugar 42 grams, ice cream 104 grams, cake 37.5 grams.
- Dinner. Chicken 89 grams, spinach 100 grams, fried potato 70 grams, bread 46 grams, butter 12 grams, coffee 150 grams, cream 100 grams, sugar 30 grams, strawberry short-cake 195 grams.

Food.	Grams.	Per	cent Nitrogen.	Total Nitrogen.
Roll	51	X	1.67 =	0.852 grams.
Sugar $49 + 42 + 30 =$	121	X	0.00 =	0.000
Orange	60	X	0.20 =	0.120
Oatmeal	150	X	0.48 =	0.645
Coffee $. 150 + 800 + 150 =$	600	X	0.06 =	0.360
Cream $. 100 + 100 + 100 =$	800	X	0.45 =	1.850
Butter $17 + 18 + 12 =$	42	X	0.15 =	0.068
Potato	114	X	0.80 =	0.842
Macaroni	115	×	0.46 =	0.529
Bread	87	X	1.57 =	0.581
Fried rice	92	X	0.75 =	0.690
Ice cream	104	X	0.53 =	0.551
Cake	87.5	X	1.20 =	0.450
Spinach	100	X	0.55 =	0.550
Chicken	89	X	8.02 =	2.688
Fried potato	70	×	0.57 =	0.399
Bread	46	X	1.91 =	0.879
Short-cake	195	×	0.50 =	0.975
Total Nitrogen	in food			12.024 grams.
Total Nitrogen	in urine	٠		9.560

Fuel value of the food . . . 8442 calories.

# Monday, May 23, 1904.

- Breakfast. Banana 98 grams, roll 68 grams, butter 15 grams, coffee 150 grams, cream 100 grams, sugar 40 grams.
- Lunch. Bread 53 grams, butter 17.5 grams, boiled eggs 101 grams, apple sauce 130 grams, coffee 150 grams, cream 50 grams, sugar 17 grams.
- Dinner. Bread 28 grams, butter 7 grams, bacon 40 grams, macaroni 62 grams, potato croquette 69 grams, coffee 150 grams, cream 50 grams, sugar 21 grams, water ice 116 grams.

Food.	Grams.	Per	cent Nitr	ogen.	Total Nitrogen.
Banana	98	×	0.28	=	0.225 grams.
Butter . $15 + 17.5 + 7 =$	39.5	×	0.15	=	0.059
Coffee $150 + 150 + 150 =$	450	×	0.06	=	0.270
Cream $100 + 50 + 50 =$	200	×	0.45	=	0.900
Sugar $.40 + 17 + 21 =$	78	×	0.00	=	0.000
Roll	<b>6</b> 8	×	1.63	=	1.108
Bread	58	×	1.64	=	0.869
Apple sauce	180	×	0.02	=	0.026
Boiled eggs	101	×	2.07	=	2.091
Bread	28	×	1.75	=	0.490
Macaroni	62	×	0.87	=	0.539
Bacon	40	×	8.28	=	1.812
Potato croquette	69	×	0.77	=	0.581
Water ice	116	X	0.006	=	0.007
Total nitrog	en in food				8.427 grams.
Total nitrog	en in urin	е.			8.030

Fuel value of the food . . . . 2346 calories.

# Tuesday, May 24, 1904.

Breakfast. — Orange 80 grams, roll 118 grams, butter 23 grams, coffee 150 grams, cream 50 grams, sugar 25 grams.

Lunch. — Soup 100 grams, bread 59 grams, butter 15.5 grams, potato 100 grams, farina croquettes 109 grams, tomato sauce 75 grams, boiled onions 107 grams, stewed prunes 105 grams, coffee 150 grams, cream 100 grams, sugar 18 grams.

Dinner. — Celery soup 150 grams, Hamburg steak 63 grams, potato 100 grams, spinach 50 grams, bread 32 grams, butter 15 grams, coffee 150 grams, cream 50 grams, sugar 16 grams, cream pie 79 grams.

Evening. — Beer 750 grams.

Food.	Grams.		Per cent Ni	rogen.	Total Nitrogen.
Roll	118	×	1.64	=	1.935 grams.
Orange	80	×	0.20	=	0.160
Coffee $150 + 150 + 150 =$	450	×	0.06	=	0.270
Cream $.50 + 100 + 50 =$	200	×	0.45	=	0.900
Sugar . $25 + 18 + 16 =$	<b>59</b>	×	0.00	=	0.000
Butter $.23 + 15.5 + 15 =$	<b>53.5</b>	×	0.15	=	0.080
Bread	<b>59</b>	×	1.66	=	0.979
Tomato sauce	75	×	0.28	=	0.178
Onions	107	×	0.30	=	0.821
Celery soup	100	×	0.48	=	0.480
Potato	100	×	0.26	=	0.260
Farina croquettes	109	×	0.74	=	0.807
Prunes	105	×	0.17	=	0.179
Hamburg steak	63	×	3.96	=	2.496
Potato	100	×	0.46	=	0.460
Soup	150	×	0.19	=	0.285
Bread	<b>82</b>	X	1.82	=	0.582
Spinach	50	×	0.54	=	0.270
Cream pie	79	×	0.98	=	0.735
Beer	750	×	0.069	=	0.518
Total nitro	gen in fo	bo			11.889 grams.
Total nitro	gen in ur	ine .			9.040

Fuel value of the food . . . . 2822 calories.

NITROGEN BALANCE. - Stapleton.

	Nitrogen	Out	put.
	Taken in. N	itrogen in Urine.	Weight of Freces (dry).
May 18	12.127 grams.	9.67 grams.	39.6 grams.
19	8.555	8.58	<b>84</b> .5
20	13.832	9.51	65.8
21	13. <del>44</del> 8	9.64	17.8
22	12.024	9.56	11.4
23	8.427	8.08	27.7
24	11.889	9.04	• • •
			196.8 grams contain
			7.08% N.
	80.302	64.03 +	13.898 grams nitrogen.
	80.802 grams r	itrogen. 77.92	8 grams nitrogen.

Nitrogen balance for seven days = +2.874 grams. Nitrogen balance per day = +0.339 gram.

#### Average Intake.

With this subject, the total intake of nitrogen for the seven days' period was 80.302 grams. The output of nitrogen through the urine amounted to 64.03 grams, while 13.898 grams were passed out through the fæces, making a total output of 77.928 grams of nitrogen. This shows a plus balance of 2.374 grams of nitrogen for the seven days, indicating a gain to the body of 0.339 gram per day. The fuel value of the food averaged 2809 calories per day, while the daily excretion of metabolized nitrogen averaged 9.14 grams. This is in close agreement with the average daily excretion of nitrogen through the urine of this subject for the last two months of the experiment, viz., 9.00 grams of nitrogen.

# With W. L. Anderson, the following results were obtained:

### W. L. ANDERSON.

## Wednesday, May 18, 1904.

Breakfast. — Banana 90 grams, fried rice 150 grams, syrup 50 grams, wheat roll 64 grams, butter 11 grams, coffee 150 grams, cream 125 grams, sugar 21 grams.

Lunch. — Tomato soup 247 grams, bread 19 grams, butter 16.5 grams, sweet potato 105 grams, farina croquette 115 grams, syrup 60 grams, coffee 150 grams, milk 50 grams, sugar 14 grams.

Dinner.—Consommé 250 grams, bread 52 grams, butter 20 grams, Hamburg steak 117 grams, boiled potato 150 grams, coffee 150 grams, cream 50 grams, sugar 14 grams.

Food.	Grams.		Per cent N	litrogen.	Total Nitrogen.
Banana	90	×	0.23	=	0.207 grams.
Cream $.125 + 50 + 50 =$	225	×	0.46	=	1.035
Sugar $. 21 + 14 + 14 =$	49	X	0.00	=	0.000
Coffee $150 + 150 + 150 =$	450	X	0.06	=	0.270
Roll	64	X	1.66	=	1.068
Butter $11 + 16.5 + 20 =$	47.5	X	0.15	=	0.071
Rice	150	X	0.75	=	1.125
Syrup $50 + 60 =$	110	×	0.024	=	0.026
Tomato soup	247	X	0.41	=	1.013
Bread	19	×	1.60	=	0.304
Sweet potato	105	X	0.32	=	0.336
Farina croquette	115	×	1.09	=	1.690
Bread	52	×	1.80	=	0.936
Consommé	250	X	0.88	=	0.950
Hamburg steak	117	×	8.64	=	4.259
Potato	150	×	0.88	=	0.570
Total nitro	gen in fo	nd.			13.855 grams.
Total nitro	•				10.030
~ ANT THE O	9~~ ·~ W		• • •		

Fuel value of the food . . . . 2946 calories.

# Thursday, May 19, 1904.

- Breakfast. Banana 158 grams, roll 122 grams, butter 15 grams, boiled hominy 150 grams, syrup 48 grams, coffee 150 grams, cream 100 grams, sugar 28 grams.
- Lunch. Bread 50 grams, butter 12 grams, fried hominy 100 grams, syrup 48 grams, boiled onions 80 grams, omelette 60 grams, coffee 150 grams, cream 50 grams, sugar 14 grams.
- Dinner. Tomato soup 200 grams, bread 43 grams, butter 10 grams, potato 50 grams, baked macaroni 202 grams, coffee 150 grams, sugar 14 grams, cream 25 grams.

Food.	Grams.	Per	cent Nitrogen.	Total Nitrogen.
Roll	122	X	1.54 =	1.879 grams.
Boiled hominy	150	X	0.20 =	0.300
Butter $15 + 12 + 10 =$	37	×	0.15 =	0.056
Banana	158	×	0.23 =	0.368
Sugar $28 + 14 + 14 =$	56	×	0.00 =	0.000
Cream . $100 + 50 + 25 =$	175	×	0.47 =	0.823
Coffee . $.150 + 150 + 150 =$	450	×	0.06 =	0.270
Syrup $48 + 48 =$	96	×	0.024 =	0.023
Bread	50	×	1.60 =	0.800
Potato	150	×	0.49 =	0.735
Fried hominy	100	×	0.67 =	0.670
Onions	80	×	0.27 =	0.216
Omelette	60	×	1.58 =	0.948
Tomato soup	200	×	0.53 =	1.060
Bread	43	×	1.74 =	0.748
Potato	50	×	0.38 =	0.190
Baked macaroni	202	X	0.93 =	1.879
Total nitrogen i	n food .			10.960 grams.
Total nitrogen i				10.150

Fuel value of the food . . . . 3018 calories.

# Friday, May 20, 1904.

Breakfast. — Coffee 150 grams, cream 25 grams, sugar 14 grams, roll 70 grams, butter 9 grams.

Lunch. — Potato 100 grams, butter 8 grams, lima beans 50 grams, hominy 69 grams, syrup 48 grams, coffee 150 grams, cream 25 grams, sugar 14 grams.

Dinner. — Consommé 150 grams, bread 28 grams, butter 15 grams, string beans 56 grams, potato 200 grams, rice croquette 65 grams, syrup 48 grams, coffee 150 grams, sugar 14 grams, cream 25 grams.

Evening. — Beer 450 grams.

Food.		Grams.	Per c	ent Nitrogen.	Total Nitrogen.
Coffee . 150	+ 150 + 150 =	450	×	0.06 =	0.270 grams.
Cream 25	5 + 25 + 25 =	75	×	0.44 =	0.830
Sugar 14	+ 14 + 14 =	42	×	0.00 =	0.000
Roll		70	×	1.72 =	1.204
Butter	.9 + 8 + 15 =	27	×	0.15 =	0.041
Potato		100	×	0.80 =	0.800
Lima beans		50	×	0.76 =	0.380
Fried hominy .		69	×	0.57 =	0.398
Syrup	48 + 48 =	96	×	0.024 =	0.023
Consommé		150	×	0.59 =	0.885
String beans .		56	×	0.86 =	0.202
Potato		200	×	0.84 =	0.680
Bread		28	×	1.97 =	0.552
Rice croquettes		65	×	1.06 =	0.689
Beer		450	×	0.069 =	0.311
	Total nitrogen	in food			. 6.260 grams.
	Total nitrogen	in urine			. 8.640

Fuel value of the food . . . . 1748 calories.

# Saturday, May 21, 1904.

- Breakfast. Banana 78 grams, bread 63 grams, butter 7 grams, coffee 150 grams, cream 50 grams, sugar 28 grams.
- Lunch. Potato 150 grams, tomato 145 grams, fried Indian-meal 81 grams, syrup 48 grams, coffee 150 grams, sugar 21 grams, cream 25 grams, water ice 165 grams.
- Dinner. Bean soup 150 grams, bread 29 grams, butter 16 grams, bacon 15 grams, fried potato 150 grams, cake 36 grams, coffee 150 grams, sugar 14 grams, cream 45 grams.

Evening. - Beer 600 grams.

Food.	Grams.	Per	ent Nitrogen.	Total Nitrogen.
Banana	73	X	0.28 =	0.168 grams.
Sugar $28 + 21 + 14 =$	63	X	0.00 =	0.000
Cream $50 + 25 + 45 =$	120	X	0.43 =	0.516
Bread	68	X	1.65 =	1.040
Butter 7 + 16 =	23	X	0.15 =	0.035
Coffee . $150 + 150 + 150 =$	450	X	0.06 =	0.270
Potato	150	X	0.71 =	1.065
Tomato	145	X	0.17 =	0.247
Fried Indian-meal	81	X	1.09 =	0.883
Syrup	48	X	0.024 =	0.012
Water ice	165	X	0.012 =	0.020
Bean soup	150	×	1.21 =	1.815
Bread	29	X	1.62 =	0.470
Bacon	15	X	8.05 =	0.458
Fried potato	150	×	0.60 =	0.900
Cake	36	×	0.86 =	0.810
Beer	600	X	0.069 =	0.414
Total nitrogen	in food .			. 8.628 grams.
Total nitrogen				. 8.460

Fuel value of the food . . . . 2393 calories.

# Sunday, May 22, 1904.

- Breakfast. Oatmeal 200 grams, sugar 28 grams, coffee 150 grams, roll 60 grams, butter 6 grams.
- Lunch.—Fried rice 140 grams, syrup 48 grams, potato 100 grams, macaroni 155 grams, boiled onions 80 grams, butter 5 grams, coffee 150 grams, cream 25 grams, sugar 14 grams, ice cream 185 grams, cake 34 grams.
- Dinner. Cream of celery soup 150 grams, mashed potato 134 grams, butter 11 grams, spinach 100 grams, strawberry short-cake 185 grams, cream 70 grams, sugar 28 grams.

Food.	Grams.	Per	cent Nitrogen.	Total Nitrogen,
Roll	<b>6</b> 0	×	1.67 =	1.002 grams.
Oatmeal	200	×	0.43 =	0.860
Sugar $28 + 14 + 28 =$	70	×	0.00 =	0.000
Coffee $150 + 150 + 150 =$	450	×	0.06 =	0.270
Butter $6+5+11=$	22	×	0.15 =	0.033
Potato	100	×	0.30 =	0.300
Macaroni	155	×	0.46 =	0.718
Cream $25 + 70 =$	95	×	0.45 =	0.428
Onions	80	×	0.25 =	0.200
Fried rice	140	×	0.75 =	1.050
Ice cream	185	×	0.53 =	0.981
Cake	84	×	1.20 =	0.408
Syrup	48	×	0.024 =	0.012
Cream of celery soup	150	×	0.88 =	0.495
Mashed potato	134	×	0.37 =	0.496
Short-cake	185	×	0.50 =	0.925
Spinach	100	×	0.55 =	0.550
Total nitrogen	in food			. 8.723 grams.
Total nitrogen				. 7.960

Fuel value of the food . . . . 2812 calories.

# Monday, May 23, 1904.

Breakfast. — Banana 115 grams, wheat griddle cakes 87 grams, syrup 48 grams, butter 7 grams, coffee 150 grams, sugar 28 grams, cream 50 grams.

Lunch. — Fried potato 100 grams, rice croquette 115 grams, syrup 48 grams, apple sauce 125 grams, coffee 150 grams, sugar 14 grams.

Dinner. — Macaroni 270 grams, potato croquette 134 grams, coffee 150 grams, sugar 14 grams, water ice 154 grams.

Food.		Grams.	Per cent	Nitrogen.	Total Nitrogen.
Banana		115	× 0.5	23 =	0.265 grams.
Coffee 150	+ 150 + 150 =	450	× 0.0	06 =	0.270
Sugar 28	+ 14 + 14 =	56	× 0.0	00 =	0.000
Cream		50	× 0.4	<b>15</b> =	0.225
Griddle cakes .		87	× 0.9	91 =	0.792
Syrup	48 + 48 =	96	× 0.0	024 =	0.023
Butter		7	× 0.:	15 =	0.011
Apple sauce .		125	× 0.0	)20 =	0.025
Rice croquette.		115	× 0.0	31 =	0.702
Fried potato .		100	× 0.6	30 =	0.600
Macaroni		270	× 0.8	37 <b>=</b>	2.349
Potato croquette		18 <b>4</b>	× 0.3	77 =	1.032
Water ice		154	× 0.0	006 =	0.009
	Total nitrogen	in food			6.808 grams.
	Total nitrogen	in urine			7.490

Fuel value of the food . . . . 2224 calories.

# Tuesday, May 24, 1904.

- Breakfast. Fried rice 115 grams, syrup 48 grams, roll 60 grams, butter 14 grams, coffee 150 grams, sugar 14 grams.
- Lunch. Celery soup 150 grams, farina croquette 108 grams, syrup 48 grams, fried potato 200 grams, bread 22 grams, butter 7 grams, coffee 150 grams, sugar 14 grams.
  - Dinner. Fried potato 200 grams, cream pie 167 grams, coffee 150 grams, cream 25 grams, sugar 14 grams.

Food.	Grams.	Per	ent Nitrogen	. Total Nitrogen.
Fried rice	115	×	0.36 =	0.414 grams.
Coffee $. 150 + 150 + 150 =$	450	×	0.06 =	0.270
Roll	60	×	1.64 =	0.984 ·
Butter $14 + 7 =$	21	×	0.15 =	0.032
Sugar $14+14+14=$	: <b>42</b>	×	0.00 =	0.000
Syrup $48 + 48 =$	: 96	×	0.024 =	0.028
Celery soup	150	×	0.48 =	0.720
Farina croquette	108	×	0.74 =	0.799
Fried potato	200	×	0.26 =	0.520
Bread	. 22	×	1.66 =	0.865
Fried potato	200	×	0.46 =	0.920
Cream pie	. 167	×	0.98 =	1.553
Cream	25	×	0.45 =	0.118
Total nitrogen	in food			. 6.718 grams,
Total nitrogen	in urine	a)		. 6.600

Fuel value of the food . . . . 2824 calories.

# 424 PHYSIOLOGICAL ECONOMY IN NUTRITION

## NITROGEN BALANCE. - W. L. Anderson.

	Nitrogen Taken in.		Output. Weight of Fæces (dry).
May 18	13.855 grams.	. 10.03 gram	s. 42:3 grams.
19	10.960	10.15	• • •
20	6.260	8.64	17.5
21	8.628	8.46	• • •
22	8.728	7.96	42.7
28	6.303	7.49	41.0
24	6.718	6.60	29.2
			172.7 grams contain
			6.30 % N.
	61.487	59.33 +	10.880 grams nitrogen.
	61.487 grams	nitrogen. 70.2	10 grams nitrogen.

Nitrogen balance for seven days = -8.778 grams. Nitrogen balance per day = -1.253 grams.

#### Average Intake.

Calories per day . . . . . . . . . . . . . . . . 2494.

Nitrogen per day . . . . . . 8.777 grams.

Here, we find a minus balance of 8.773 grams of nitrogen for the seven days' period. The total intake of nitrogen amounted to 61.437 grams, while there were excreted through the urine 59.33 grams, and through the fæces 10.880 grams of nitrogen, making a total output of 70.210 grams as contrasted with an intake of 61.437 grams of nitrogen. loss of body material is to be attributed to the small fuel value of the food, - only 2494 calories as the day's average, - though perhaps in part to the relatively small intake of nitrogen. In this connection it is to be noted that the average daily excretion of metabolized nitrogen for the seven days' period amounted to only 8.777 grams, while the average daily excretion for the last two months of the experiment was as high as 10.07 grams. Undoubtedly, the subject did not eat as much food during the week of this balance trial as was needed to maintain equilibrium, under the conditions of bodily activity then prevailing.

The same statement applies to Mr. Bellis, whose balance trial likewise shows a deficiency of ingested nitrogen over the nitrogen output. Here, however, the deficiency is more manifestly due to the small fuel value of the daily food, which averaged only 2174 calories. Bellis showed an average daily excretion of metabolized nitrogen amounting to 8.45 grams for the last six weeks of the experiment, while in the balance period the excretion of metabolized nitrogen was 8,19 grams per day. The daily intake of nitrogen in the food, however, averaged only 7.76 grams, obviously too small a quantity to meet the wants of the body, especially with the low fuel value of the food. It is quite plain that during the week of this balance trial, the amount of food consumed was not equal to the necessities of the body, neither was it equal in nitrogen or fuel value to what the subject had been taking during the last few months of the experiment, and on which he had practically maintained body-weight for at least the last month of the experiment. It is further noticeable that during the balance week the body-weight dropped off somewhat.

Owing to the absence of Dr. Callahan from New Haven during this period, no attempt was made to determine experimentally whether he was in nitrogen equilibrium or not.

The following tables give the data in the experiment with Bellis:—

#### BELLIS.

## Wednesday, May 18, 1904.

Breakfast. — Banana 94 grams, wheat roll 53 grams, butter 11 grams, coffee 150 grams, cream 75 grams, sugar 21 grams.

Lunch. — Soup 150 grams, farina croquette 100 grams, syrup 50 grams, string beans 75 grams, fried sweet potato 117 grams, bread 86 grams, butter 7 grams, coffee 150 grams, sugar 14 grams.

Dinner. — Hamburg steak 58 grams, potato 250 grams, spinach 100 grams, bread 55 grams, butter 10 grams, coffee 150 grams, cream 75 grams, sugar 21 grams, apple pie 142 grams.

Food.	,	Grams.	Pe	r cent Nitro	gen.	Total Nitrogen.
Banana	•	94	X	0.28	=	0.216 grams.
Cream 75 + 75	=	150	X	0.46	=	0.690
Sugar $21 + 14 + 21$	=	56	X	0.00	=	0.000
Coffee $. 150 + 150 + 150$	=	450	×	0.06	=	0.270
Roll (wheat)		58	X	1.66	=	0.880
Butter $.11 + 7 + 10$	=	28	X	0.15	=	0.042
Soup		150	X	0.41	=	0.615
Farina croquette		100	X	1.09	=	1.090
Fried sweet potato		117	×	0.32	=	0.374
String beans		75	X	0.84	=	0.255
Syrup		50	X	0.024	=	0.012
Bread		86	×	1.60	=	0.576
Hamburg steak		53	×	3.64	=	1.929
Potato		250	×	0.88	=	0.950
Spinach		100	X	0.53	=	0.530
Bread		55	×	1.80	=	0.990
Apple pie		142	×	0.48	=	0.611
Total nitrog	en i	in food				. 10.080 grams.
Total nitrog	en i	in urine	• .		•	. 8.850

Fuel value of the food . . . . 2686 calories.

## Thursday, May 19, 1904.

- Breakfast. Banana 155 grams, roll 53 grams, butter 10 grams, coffee 150 grams, sugar 28 grams, cream 70 grams.
- Lunch.—Fried hominy 60 grams, syrup 48 grams, potato 100 grams, boiled onion 82 grams, coffee 150 grams, cream 35 grams, sugar 21 grams, bread pudding 134 grams.
- Dinner. Tomato soup 200 grams, macaroni 75 grams, mashed potato 54 grams, spinach 70 grams, boiled Indian-meal 100 grams, coffee 150 grams, cream 80 grams, sugar 21 grams.

Food.	Grams.	Per c	ent Nitrogen.	Total Nitrogen.
Roll	53	×	1.54 =	0.816 grams.
Banana	. 155	×	0.23 =	0.857
Butter	. 10	×	0.15 =	0.015
Coffee $. 150 + 150 + 150 =$	450	×	0.06 =	0.270
Sugar $28 + 21 + 21 =$	70	×	0.00 =	0.000
Cream $. 70 + 35 + 80 =$	185	×	0.47 =	0.870
Potato	100	×	0.49 =	0.490
Fried hominy	60	×	0.67 =	0.402
Boiled onion	. 82	×	0.27 =	0.221
Bread pudding	. 134	×	0.28 =	0.375
Syrup	48	×	0.024 =	0.012
Tomato soup	200	×	0.58 =	1.060
Macaroni	. 75	×	0.98 =	0.698
Mashed potato	54	×	0.38 =	0.205
Spinach	. 70	×	0.56 =	0.392
Boiled Indian-meal	. 100	×	0.20 =	0.200
Total nitrogen	in food			. 6.388 grams.
Total nitrogen	in urine			. 9.600

Fuel value of the food . . . . 2075 calories.

# Friday, May 20, 1904.

Breakfast.—Orange 150 grams, roll 57 grams, butter 17 grams, coffee 159 grams, sugar 14 grams.

Lunch. — Fried farina 74 grams, syrup 48 grams, potato 250 grams, lima beans 50 grams, coffee 150 grams, sugar 14 grams.

Dinner. — Rice croquette 92 grams, syrup 48 grams, string beans 93 grams, mashed potato 352 grams, bread 40 grams, butter 8 grams, coffee 150 grams, sugar 14 grams.

Evening. — Beer 450 grams.

Food.	Grams.	Per	Total Nitrogen.		
Orange	. 150	×	0.20	=	0.300 grams.
Roll	. 57	×	1.72	=	0.980
Coffee $. 150 + 150 + 150 =$	450	×	0.06	=	0.270
Sugar . $14 + 14 + 14 =$	<b>= 42</b>	×	0.00	=	0.000
Butter 17 + 8 =	<b>25</b>	×	0.15	=	0.088
Potato	250	×	0.80	=	0.750
Lima beans	. 50	×	0.76	=	0.880
Fried farina	. 74	×	0.57	=	0.422
Syrup $48 + 48 =$	96	×	0.024	=	0.023
String beans	93	×	0.86	=	0.835
Bread	. 40	×	1.97	=	0.788
Mashed potato	352	×	0.34	=	1.197
Rice croquette	92	×	1.06	=	0.975
Beer	450	×	0.069	=	0.311
Total nitrogen	in food				6.769 grams.
Total nitrogen					10.670

Fuel value of the food . . . . 1980 calories.

## Saturday, May 21, 1904.

- Breakfast Banana 69 grams, baked potato 57 grams, bread 59 grams, butter 8 grams, coffee 150 grams, sugar 14 grams.
- Lunch. Fried Indian-meal 80 grams, syrup 48 grams, potato croquette 152 grams, tomato 147 grams, coffee 160 grams, sugar 14 grams, water ice 168 grams.
- Dinner.— Bean soup 150 grams, bacon 18 grams, fried egg 50 grams, fried potato 206 grams, lettuce salad 45 grams, bread 38 grams, butter 8 grams, coffee 150 grams, sugar 14 grams.

Evening. — Beer 600 grams.

Food.	Grams.	Per	cent Nitro	gen.	Total Nitrogen.
Banana	. 69	×	0.23	=	0.159 grams.
Bread	. 59	×	1.65	=	0.974
Coffee $. 150 + 150 + 150$	= 450	×	0.06	=	0.270
Sugar. $14 + 14 + 14$	= 42	×	0.00	=	0.000
Baked potato	. 57	×	0.40	=	0.228
Butter 8 + 8	= 16	×	0.15	=	0.024
Potato croquette	. 152	×	0.71	=	1.079
Tomato	. 147	×	0.17	=	0.250
Fried Indian-meal	. 80	×	1.09	=	0.872
Syrup	. 48	×	0.024	=	0.012
Water ice	. 163	×	0.012	=	0.020
Bread	. 88	×	1.62	=	0.616
Bacon	. 13	×	8.05	=	0.397
Fried egg	. 50	×	2.27	=	1.135
Bean soup	. 150	×	1.21	=	1.815
Lettuce salad	. 45	×	0.21	=	0.095
Fried potato	. 206	×	0.60	=	1.236
Beer	. 600	×	0.069	=	0.414
Total nitroge	n in food				9.596 grams.
Total nitroge	n in urine				0.480

Fuel value of the food . . . . 2071 calories.

## Sunday, May 22, 1904.

Breakfast. — Orange 100 grams, oatmeal 100 grams, roll 50 grams, butter 8 grams, coffee 150 grams, cream 40 grams, sugar 21 grams.

Lunch. — Macaroni 112 grams, potato 200 grams, onions 143 grams, coffee 150 grams, sugar 7 grams, ice cream 170 grams, cake 81 grams.

Dinner. — Cream of celery soup 150 grams, mashed potato 182 grams, spinach 100 grams, coffee 150 grams, sugar 7 grams, strawberry short-cake 97 grams.

Food.	1	Grams.	Per Cent	Nitrogen.	Total Nitrogen.
Orange		. 100	× 0.2	0 =	0.200 grams.
Roll		. 50	× 1.6	7 =	0.835
Butter		. 8	× 0.1	5 =	0.012
Oatmeal		. 100	× 0.4	3 =	0.480
Coffee . 150 -	+ 150 + 150	= 450	× 0.0	6 =	0.270
Sugar 21 -	+ 7+ 7	= 85	× 0.0	0 =	0.000
Cream		. 40	× 0.4	5 <b>=</b>	0.180
Macaroni		. 112	× 0.4	6 =	0.515
Potato		. 200	× 0.8	0 =	0.600
Onions		. 143	× 0.2	5 =	0.858
Ice cream		. 170	× 0.5	8 =	0.901
Cake		. 81	X 1.2	<b>20</b> =	0.372
Cream of Celery	soup	. 150	× 0.8	3 =	0.495
Mashed potato		. 182	× 0.3	7 =	0.673
Spinach		. 100	× 0.5	5 =	0.550
Strawberry short	t-cake	. 97	× 0.5	0 =	0.485
	Total nitrog	en in food			6.876 grams.
	Total nitrog	en in urine			7.710

Fuel value of the food . . . 1929 calories.

## Monday, May 23, 1904.

- Breakfast.—Banana 219 grams, coffee 150 grams, cream 80 grams, sugar 28 grams.
- Lunch. Rice croquette 148 grams, syrup 45 grams, potato 200 grams, coffee 150 grams, sugar 7 grams, apple sauce 250 grams.
- Dinner. Vegetable soup 150 grams, bacon 37 grams, string beans 100 grams, potato 101 grams, macaroni 86 grams, coffee 150 grams, water ice 184 grams.

Food.		Grams.	Per	Total Nitrogen.	
Banana		219	×	0.23 =	0.504 grams.
Cream		80	×	0.45 =	0.360
Sugar	28 + 7 =	85	×	0.00 =	0.000
Coffee . 150 +	150 + 150 =	450	×	0.06	0.270
Apple sauce		250	×	0.02 =	0.050
Rice croquette .		143	×	0.61 =	0.872
Syrup		45	×	0.024 =	<b>0</b> .011
Potato		200	×	0.60 =	1.200
Macaroni		86	×	0.87 =	0.748
Vegetable soup .		150	×	0.70 =	1.060
Bacon		87	×	3.28 =	1.214
String beans	<b>.</b>	100	×	0.22 =	0.220
Potato croquette .		101 ~	×	0.77 =	0.778
Water ice		184	×	0.006 =	0.011
	Total nitrogen				. 7.288 grams.
, <b>!</b>	Fotal nitrogen	in urine			. 5.980

Fuel value of the food . . . . 2226 calories.

# Tuesday, May 24, 1904.

Breakfast. — Orange 100 grams, baked potato 188 grams, butter 5 grams, coffee 150 grams, sugar 14 grams.

Lunch. — Celery soup 150 grams, farina croquette 91 grams, syrup 48 grams, boiled onions 110 grams, potato 200 grams, stewed prunes 113 grams.

Dinner. — Tomato soup 150 grams, farina croquette 107 grams, syrup 48 grams, potato 200 grams, spinach 180 grams, cream pie 140 grams.

Food.						•	Frams.	Pe	er cent	Nitro	gen.	Total Nitrogen.
Orange							100	×	0.2	20	=	0.200 grams.
Baked potato .							138	X	0.5	25	=	0.845
Coffee							150	×	0.0	06	=	0.090
Sugar							14	×	0.0	00	=	0.000
Butter							5	×	0.1	15	=	0.008
Boiled onions .							110	×	0.8	<b>30</b>	=	0.330
Celery soup .							150	×	0.4	18	=	0.720
Potato							200	×	0.2	26	=	0.520
Farina croquette							91	×	0.7	74	=	0.673
Syrup		48	+	48	:	=	96	×	0.0	)24	=	0.023
Stewed prunes							118	×	0.1	17	=	0.192
Spinach							180	×	0.6	54	=	0.972
Tomato soup .							150	×	0.1	19	=	0.285
Potato							200	×	0.4	16	=	0.920
Farina croquette							107	×	0.3	76	=	0.813
Cream pie							140	×	0.9	93	=	1.302
	To	tal	ni	itro	ge	n i	n food .					7.393 grams.
	To	tal	ni	itro	ge	a ir	urine					6.610

Fuel value of the food . . . . 2254 calories.

## NITROGEN BALANCE. - Bellis.

	Nitrogen Taken in.	Outp Nitrogen in Urine.	ut. Weight of Fæces (dry).			
May 18	10.030 grams	. 8.85 grams.	• • •			
19	6.388	9.60	• • •			
20	6.769	10.67	• • •			
21	9.596	8.46	51.0 grams.			
22	6.876	7.71	46.0			
23	7.288	5.98	27.2			
24	7.398	6.61	57.5			
			181.7 grams contain			
			6.88 % N.			
	54.385	57.88 +	11.592 grams nitrogen			
	54.835 grams	nitrogen. 68.97	2 grams nitrogen.			

Nitrogen balance for seven days = -14.637 grams. Nitrogen balance per day = -2.091 grams.

#### Average Intake.

Calories per day . . . . . . . . . . . . . . . . . 2174.

Nitrogen per day . . . . . . . . . . . . . . . . . 7.762 grams.

THE PHYSICAL CONDITION OF THE SUBJECTS.

Having considered the marked decline in the extent of proteid metabolism which these subjects have exhibited for a period of five months, and having shown the possibility of their maintaining body-weight and nitrogen equilibrium on a low proteid intake, coupled with a relatively small amount (low fuel value) of non-nitrogenous food, it is appropriate to consider next their physical condition under this changed mode of living. So much has been written upon the necessity of a rich proteid diet, with a corresponding rate of proteid metabolism, for the maintenance of bodily strength and vigor, that it becomes a question of vital importance to obtain data bearing upon the effect of a lowered proteid intake upon bodily strength. If, as is so widely believed, diminishing the daily proportion of proteid food below the standards set by Voit and other physiologists will result in a weakening of the muscles of the body, in decreasing the strength, vigor, and endurance of the individual, then obviously physiological economy in this direction would in the long run be uneconomical, and indeed injurious. The maintenance of body-weight and of nitrogen equilibrium on a small amount of proteid food would count for little, when compared with a gradual loss of bodily strength and vigor.

It was truly a great surprise when the systematic strength tests applied month after month to the soldiers indicated a marked gain in muscular power, which seemingly increased as the rate of proteid metabolism diminished, coincident with the decrease in the amount of proteid food fed. The dynamometer tests were applied primarily to make sure there was no falling off in strength, and when the marked gains already referred to were recorded, it was thought at first that they must be the result mainly of the systematic training the soldiers were undergoing in the gymnasium. Undoubtedly, this daily training, with the more regular and systematic methods of living, did contribute in some measure to the beneficial results obtained, but as the improvement and general gain in

strength became more and more apparent, it was equally clear there were other factors involved than mere training.

The opportunity presented by the present subjects therefore was particularly desirable. These men had been in training for many months, some of them for several years, and naturally had acquired a high degree of proficiency in all kinds of athletic work, in the handling of themselves and in the handling of the apparatus, by use of which the strength tests are made.

The tests, etc., were applied exactly in the same manner as in the case of the soldier detail, description of which will be found on pages 259 and 260.

The following tables give the results of the tests — made at the Yale Gymnasium, and reported by Dr. Anderson — from January to June, for the eight men. It will be noted, however, that the record of Mr. Bellis is incomplete. This was owing to an injury to his hand, which prevented his working with the apparatus during the months of May and June.

The results presented by these tables are very important and suggestive. Every man, without exception, showed a decided improvement in his muscular power as measured by the strength tests. With many of the men the gain was progressive, with others there was noticeable - as in the case of W. L. Anderson and G. W. Anderson in the March test — a drop in some one test. This could generally be explained by some temporary cause. Thus, the March test taken by W. L. Anderson was at a time when he was under great strain in connection with an intercollegiate meet, etc. However, it is clear from the figures presented that all these men, though living on a greatly reduced amount of proteid food, and with certainly no increase in the quantity of non-nitrogenous food, showed at the end of the experiment a decided gain in muscular power. Note for example the great gain in strength shown by Schenker; in January his dynamometer tests, etc., indicated a total of 5728, while at the close of the experiment in June his record was 7135. Again, Bellis increased from 5993 to 8165, and W. L. Anderson from 6016 to 9472. Further, the

# 436 PHYSIOLOGICAL ECONOMY IN NUTRITION

## STRENGTH OR DYNANOMETER TESTS.

					Ann	erso	ж, G.	w.					
	Weight.	Lang Capacity.	Bight Hand.	Left Hand.	Obest.	Back.	Legs.	Pall up.	Push up.	Vault and Ladder.	Run.	Product.	Total.
Jan. 20 Feb. 20	168 163	500 535	108 109	97 105	85 70	425 380	900 570	8 10	11 14	E	1.05 1.07	3300 3972	4913 5206
Mar. 20 Apr. 20 May 26	159 161 159	530 500 582	110 101 120	103 97 100	75 85 80	410 490 475	720 810 885	10 18 9	9 11 15	E	1.07 1.05	3021 3864 3816	4489 5387 5476
June 17	157	580	100	90	80	580	840	12	14		:::	4082	5722
	Anderson, W. L.												
Jan. 20	139	420	95	88	145	410	625	18	20	E	0.48	4553	6016
Feb. 20 Mar. 20	136 185	440 480	95 95	80 85	137 140	550 510	560 570	15 11	25 15	E	0.45 0.47	5520 3510	<b>694</b> 2 <b>489</b> 0
Apr. 20	138	460	93	75	165	650	730	12	30		1.04	5806	7519
May 26	134	450	90	70	155	570	880	25	31			7504	9267
June 17	137	450	95	85	160	600	860	30	26	• •		7672	9472
						Brl	LIS.						
Jan. 20	180	550	130	180	155	550	825	10	13	E	1.06	4653	5993
Feb. 20	174	500	155	135	135	580	925	11	15	E	1.08	4615	6445
Mar. 20	176	550	150	135	155	500	900	12	20	E	1.08	5728	7568
Apr. 20	177	510	160	150	150	560	1111	13	21		• •	6035	8165
					(	CALL	ahan.						
Jan. 20	204	565	105	110	145	450	620	2	1	D	1.20	724	2154
Feb. 20	198	560	120	120	145	440	590	4	8	F	1.18	1865	2780
Mar. 20 Apr. 20	185 181	565 580	125 120	115 120	145 145	420 530	650 685	5 5	4	E	1.15 1.16	1692 1629	8142 8229
June 18	184		120	115	140	520	890	7	5		1.14	2208	3983

					]	Dona	HUE.						
	Weight.	Lung Capacity.	Right Hand.	Left Hand.	Chest.	Back.	Lega.	Pull up.	Push up.	Vault and Ladder.	Run,	Product.	Total.
Jan. 20	142	820	95	100	74	330	500	12	12	E	1.04	3480	4584
Feb. 20	136	345	105	115	95	340	490	13	14	E	1.04	3758	4905
Mar. 20	187	400	110	95	100	<b>340</b>	710	15	16	E	1.06	4809	5664
May 1	138	420	95	80	115	360	650	17	16	• •	• • •	4554	5854
May 26	137	425	108	90	185	430	570	17	16		• • •	4589	5917
							·'				·		<u>.                                    </u>
						Jaco	BUS.						
Jan. 20	126	350	97	93	96	250	800	14	14	E	1.00	8712	4548
Feb. 20	124	380	97	97	75	460	470	15	15	E	1.02	3683	4888
Mar. 20	125	870	95	95	80	430	490	15	15	E	1.08	8810	5000
May 2	124	350	85	90	100	850	510	15	14	• •	1.03	8610	4740
May 26	125	360	95	95	110	860	600	14	17	$ \cdot \cdot$		8875	5185
June 16	125	875	95	85	120	400	700	15	19		• • •	4267	5667
		•				Всни	NEER			<u></u>	•		,
		1	1	·	<del></del>	1	i	1	· · · ·	1	1		1
Jan. 20	161	485	112	90	185	410	440	14	14	E	1.06	4771	5728
Feb. 20	159	490	115	90	125	530	730	14	15	E	1.07	4748	6338
Mar. 20	158	500	100	95	120	550	780	15	15	E	1.12	4830	6475
Apr. 20	164	490	90	90	127	410	580	17	15	1	1.10	5248	6545
May 26	159	500	125	105	155	470	850	16	15	• •	1.08	5104	6809
June 9	160	500	110	110	155	570	910	17	16			5280	7135
					i	·	· · · · ·		•	•	·	·	-
						TAPL	BTON	•					
Jan. 20	170	520	105	105	155	455	570	11	12	E	1.21	8961	5351
Feb. 20	167	520	105	110	160	550	690	6	10	E	1.22	2704	4319
Mar. 20	170	500	100	100	145	350	600	7	12	E		3268	4563
Apr. 20	162	500	100	105	155	410	670	9	14			8726	5886
May 26	164	515	100	95	150	470	770	12	20			5248	6838

men all agree in the good effect the changed conditions have had upon them, and they have, without exception, been able to do their athletic work and maintain their athletic supremacy.

Naturally, in the case of these men the gain in strength recorded cannot be assigned to systematic training. The only change in their mode of living which can in any sense be considered as responsible for the improvement is the change in diet. The main fact to be emphasized, however, is that these men—trained athletes, accustomed to living on relatively large amounts of proteid food—for a period of five months reduced their intake of proteid food more than fifty per cent without loss of bodily strength, but, on the contrary, with a marked improvement in their muscular power.

Most striking is this gain in strength when compared with the very marked decline in the rate of proteid metabolism. Thus, in the case of Jacobus, the excretion of metabolized nitrogen was reduced to 7.43 grams per day as the average for the last two months of the experiment, yet his strength test showed an increase from 4548 in January to 5667 for June. Further, it must be recalled that an excretion of 7.43 grams of nitrogen means the metabolism of only 46.4 grams of proteid Similarly, in the case of Donahue, a very active man whose work on the Varsity basket-ball team called for vigorous exercise, his strength test rose from 4584 to 5917 on a daily diet which led to the metabolism of only 7.39 grams of nitrogen per day, or about 46 grams of proteid matter. Further, Donahue frequently referred to the far greater freedom from fatigue he experienced on the low proteid diet, and he was clearly conscious of a distinct improvement in his physical condition.

The following letter from Dr. Anderson, the Director of the Yale Gymnasium, gives his estimate of the men at the end of the first three months of the experiment:

YALE UNIVERSITY GYMNASIUM, NEW HAVEN, CONN., April 12, 1904.

Professor Russell H. Chittenden,
Director of the Sheffield Scientific
School, Yale University.

DEAR SIR, — Herewith find a brief report of the physical condition and ability of the eight Yale students who are taking the special diet while engaged in active exercise.

These men, with one exception, Dr. Callahan, are experts in their special lines of avocation.

- Mr. G. W. Anderson is a foot-ball, base-ball, and basket-ball player, as well as a crew man (not Varsity), well built and an all round athlete.
- Mr. W. L. Anderson, a "Y" athlete (hurdler), the captain of the Yale Gymnastic Team, University Gymnastic Champion, and American Collegiate Gymnastic Champion.
- Mr. H. S. Bellis, a member of the Y. G. A., a gymnast and acrobat and in constant training.
- Dr. W. H. Callahan, Medical Assistant at the Gymnasium, in daily practice in the gymnasium; bowling, hand-ball, and running.
- Mr. M. Donahue, a very muscular and versatile athlete, a football player and a Varsity basket-ball player.
- Mr. C. S. Jacobus, a "Y" athlete, a noted long-distance man, and one of the best University runners.
- Mr. H. R. Schenker, an active member of the Y. G. A., a point winner and intercollegiate competitor in gymnastics.
- Mr. John Stapleton, a wrestler and gymnast. A professional, a man of large body and great strength.

These eight men are in constant practice and in the "pink of condition." They were in "training form" when they began the changed diet. All have lost in weight, especially Dr. Callahan, who has dropped from 204 pounds to 185 pounds in two months. Dr. Callahan is not an athlete, but is a vigorous worker in the gymnasium, being in daily and constant practice. He is liberally supplied with adipose tissue and can well afford to drop in weight.

As to the loss of weight in the other cases, it would not be wise to attribute this to the diet alone. We find that most athletes who represent the University in the big contests lose in body-weight, but I attribute this loss as much to worry and responsibility as to strict bodily activity.

These students are in a different class from the soldiers, first, because they are well educated young men, secondly, because their development was towards a specific end, the attainment of strength and skill as representative Yale athletes, and thirdly, on account of college requirements of fifteen hours per week, which time stands for study and laboratory attendance aside from the recitations. We have here a double drain on the body energy. All mental work is expensive, hence the demand upon the corporeal machinery has been very constant and strenuous.

I notice little change in the condition of the men over that of a year ago, when I had most of them with me and under like physical training.

In the case of W. L. Anderson, captain of the Yale Gymnastic Association, there was a noticeable falling off in the strength tests in February and March, but I believe the worry incident to the intercollegiate contests, the steady training, and the business cares of the Association went far towards producing a fatigued state. W. L. Anderson is only a freshman in the Medical School; he did his studying at night, and this combined with his youth doubtless caused the loss of weight as much as any change in diet. He has shown the same symptoms before this year. At this writing he is in good physical condition. I speak with certainty in his case because I have had good opportunity to study him at home. It was while under the restricted diet that he won both championships, these being the Collegiate and All-around Intercollegiate Championship of America.\*

Schenker won points for the first time in the intercollegiate contests while on the diet; he showed no falling off, rather to the contrary, made a steady gain in ability.

Jacobus complained of a pain in his side but in spite of this he has entered a number of events, has kept up his training and is in good condition. Jacobus is a long-distance runner; great endurance is required for these events, and this endurance he has

<sup>\*</sup> Italics inserted by R. H. C.



W. L. ANDERSON BELLIS

Photographs taken prior to the experiment.



kept up. He tells me his stomach is in better condition than it has been during his three years of work at Yale.

Donahue has steadily improved in ability. He has kept his position on the Varsity Basket Ball Team, and has put up strong and aggressive games, and says he is as well as ever.

Stapleton shows no falling off at all. He keeps up wrestling, which is a drastic exercise; he works at heavy gymnastics and gains steadily.

One matter must be reported in reference to the strength tests. The first trial was made when all members of the squad were present. College men are very sensitive to competition, hence the great exertion put forth. The other trials were made when the men were by themselves. The "spur" was missing.

I have watched the efforts of these men with interest and care, especially as two of them live in my own family. I fail to see any falling off in strength, the case of W. L. Anderson excepted. The fellows report being in satisfactory shape and claim that the "ups and downs" are no more in evidence this year than in the past.

These picked men, representing several kinds of competitive sports, have gained in ability and skill on the more limited diet they are now using, and are not showing any signs of deterioration from the diminished intake of proteid food. I pronounce them, from a physical standpoint, in good shape.

Respectfully yours,

(Signed) WILLIAM G. ANDERSON.

It must be remembered that this letter from Dr. Anderson was written after the March strength test was taken, and prior to the test of April 20. Dr. Anderson was not in New Haven at the close of the experiment, consequently it was not possible to obtain his estimate of the men at that date, but there can be no question that there was a distinct improvement from the middle of April to the middle of June; certainly as marked as the improvement from the beginning of the experiment in January, to April 12, the date of Dr. Anderson's letter.

Finally, attention may be called to the photographs of Messrs. Stapleton, Bellis, and W. L. Anderson, which are introduced primarily to show the physical make-up and muscular development of the men composing this student group. The photographs of Stapleton were taken in April, 1904, after he had been under experiment for three months. The photographs of Bellis and W. L. Anderson were taken prior to the experiment. They all afford a good illustration of the highly developed muscular mechanism of different types, with a corresponding adaptability for different lines of muscular effort.

## REACTION TIME.

Through the courtesy and kind co-operation of Dr. Charles H. Judd, in charge of the Yale Psychological Laboratory, these students were subjected to the same careful tests during the five months of their experiment as were applied to the soldier detail. The results which are presented in the following report, kindly prepared by Dr. Judd, indicate quite clearly that there was no general nervous change in the reactors as a result of the low proteid diet. The data presented by Dr. Judd in this connection will be found in the accompanying tables, which, while indicating no noticeable improvement in the nervous condition of the men, make it quite plain that no deterioration whatever occurred as the result of the lowered proteid metabolism.

## REPORT ON REACTION

Reaction tests with the group of University students were conducted in essentially the same way as were the reaction tests with the soldiers. Details in regard to the method and apparatus employed need not be repeated. They can be found on pages 274 to 276.

Two new tests were added to the regular reaction determinations. One of these consisted in taking a record of the number of taps which could be executed in ten seconds. The reactor was seated before a table on which was fastened a telegraph key. He held the key between his thumb and first two fingers, and at a given signal began tapping as rapidly as possible until told to stop. Each time he tapped he closed an electric circuit. The 医马尔氏血管医

W. L. ANDERSON BELLIS

Photographs taken prior to the experiment.

,				
	•			
	-			
			1	
			1	
		•	ı	

current thus made was carried through a marker which indicated on a smoked paper each make and break at the key. A time line from a rod vibrating at the rate of once every twentieth of a second was traced on the smoked paper parallel with the marker record. By a comparison of the time line with the marker record, it was easy to determine the number of taps made in ten seconds. At first, the number of taps per second were counted, but the results showed such uniformity from second to second on a given day that only the net results for the whole ten seconds are given in the tables.

The second test added to the reaction tests was undertaken to determine the steadiness of the subjects. Two brass rods 40 cm. long were held in a vertical position at a distance of 7 mm. from each other. The subject took in his right hand a brass rod 40 cm. long and 5 mm. in diameter and tried to pass the end of this rod up and down between the vertical rods without touching them. The subject's relation to the vertical rods may be further defined by saying that he stood directly in front of them and reached out nearly at arm's length. The vertical rods were at about the height of his chest. In order to get a record of the accuracy with which the subject moved the hand-rod up and down without touching the vertical rods, the vertical rods and the hand-rod were connected with the two poles of an electric circuit. Whenever they touched they closed the circuit, and a marker placed in the same circuit recorded the fact on a smoked paper record. The time was recorded in parallel with this record, so that any long continued contact could be measured. For the most part, contacts were only of very brief duration. In reporting the results of this test, every contact made while passing the rod once downward and once upward is counted as at least one. Continued contacts are recorded as two, three or more, according to the period of duration. Where there are such added counts because of continued contacts, a second quantity is given in the tables after the first. This second quantity, which is enclosed in parenthesis, indicates merely the number of contacts without reference to whether they are long or short.

No special comments are necessary to explain the tables. Tables 1-5 report in sigmas, or thousandths of a second, the average time of ten reactions on the date, and for the subject,

indicated. In the third column, is the mean variation for the series of ten reactions.

Table 6 shows the general averages by the month for a given individual, and in the sixth column the general average of all the determinations for each reactor. The fourth column in each of the earlier tables shows how much the results of a given day vary from the final general average. All the tables give, at the bottom, group averages whenever the full series is present.

Table 7 gives the number of taps executed in ten seconds at each successive test. The dates are not given in detail, but are the same as those of the reaction tests.

Table 8 gives the monthly averages of taps.

Table 9 presents the results of the steadiness tests.

The three sets of results, namely, those from reaction, tapping, and steadiness, differ from each other. On the whole, the reactions grow longer; the tapping varies, but shows neither decided improvement nor deterioration; while steadiness improves very decidedly.

The comment made on the results obtained with the soldiers applies here so far as the reaction tests are concerned. These tests were not repeated with sufficient frequency to reduce the reaction to automatic performance.

The tapping is such a simple performance that improvement is not to be expected. The absence of any general improvement or deterioration argues for an absence of any general nervous change in the reactors.

The improvement in steadiness is in part at least, probably in very large measure, due to the fact that the subjects became more familiar with the test and approached it with less of the embarrassment which attends a new and unfamiliar test.

The lengthening of the reaction times indicates a less intense concentration of the subject upon the work in hand. The slightest relaxation of attention puts the subject behind in responding to the signal. It was clear to superficial observation, especially in certain individual cases, that the subject was giving less attention in the later experiments. This lack of concentration is not obviously related to the changes in diet. Indeed, the fact that no corresponding falling off appears in the tapping would seem to argue that the lack of attention in the reaction tests was not due

to deep-seated nervous conditions, so much as to growing impatience on the part of the reactors with the ordeal of being tested. The tapping experiment is less likely to be affected by lack of interest on the part of the subject, because here the subject is called upon to be constantly active, and there is no such opportunity for attention to lapse as is furnished by the intervals which intervene between successive reactions. The tapping is accordingly perhaps the best series on which to base final judgment as to the nervous condition of the men. Here, there appear the variations which show in any ordinary series, but there is no steady improvement through growing familiarity with the test, nor any laxness of attention to produce relatively unfavorable results.

Mr. Steele and Dr. McAllister are largely responsible for the actual collection of the data on which this report is based.

(Signed) C. H. Judd.

Name.	Date.	Avg.	M. v.	Var. from G. A.	Date.	Avg.	M.v.	Var. from G. A.
I. Anderson, G. W	19	153.5	13.8	81.4	26	192.2	19.4	7.3
II. Anderson, W. L	18	261.0	82.5	82.5	27	298.6	71.0	65.1
III. Bellis	16	189.3	40.0	8.2	26	182.3	18.5	0.5
IV. Donahue	17	173.6	43.0	28.4	24	182.7	14.5	19.8
V. Jacobus	17	197.8	14.7	27.6	24	178.2	18.3	46.7
VI. Schenker	17	162.8	29.7	67.6	24	209.8	46.6	20.1
VII. Stapleton	19	185.1	31.6	23.9				
Group averages	•	188.8						
VIII. Callahan *	18	156.2	10.4	37.0	25	210.8	27.2	17.6

TABLE 1. - FEBRUARY, 1904.

<sup>\*</sup> Kept separate because series of tests is not complete.

TABLE 2. - MARCH, 1904.

Name.	Date.	Avg.	E. v.	Var. from G. A.	Dete.	Avg.	M. v.	Var. from G. A.	Date	Avg.	M. v.	Var. from G. A.	Date.	Avg.	M. v.	Var. from G. A.
I. Anderson. G. W.		:	:	:	11	181.6	88.3	8.8	18	181.8	25.1	8.6	88	170.5	28.1	14.4
II. Anderson, W. L.	∞	286.4	50.4	0.00	91	207.8	82.2	21.2	11	189.3	41.3	89.2	:	:	:	:
III. Bellis	<b>∞</b>	1740	44.4	7.8	16	180.8	21.8	1.0	83	198.5	16.5	11.7	88	181.0	8.5	0.8
IV. Donahue	64	201.6	24.8	9.0	6	214.7	29.3	12.7	क्ष	224.9	58.7	877	:	:	:	:
V. Jacobus	81	207.5	14.4	17.4	6	189.9	11.9	35.0	16	239.6	41.0	14.7	83	196.7	27.2	28.2
VI. Schenker	63	283.5	0.86	63.6	8	296.9	74.9	67.0	16	281.9	80.8	62.8	:	:	:	:
VII. Stapleton	4	211.1	41.7	2.1	=	206.4	14.6	8.6	:	:	:	:	:	:	:	:
Group averages	188	:				210.9										
VIII. Callahan	ဇာ	202.0	23.9	8.8	10	197.0	29.0	8.8	17	200.9	23.0	7.7	24	179.9	20.6	18.8

TABLE 8. — APRIL, 1904.

Name.	Date.	Avg.	M. V.	Var. from G. A.	Date.	Avg.	<b>W</b> . V.	Var. from G. A.
I. Anderson, G. W.	22	202.8	10.6	17.9	29	182.9	6.7	2.0
II. Anderson, W. L.					29	195.0	20.8	33.5
III. Bellis	19	193.0	27.2	11.2	26	182.5	18.2	0.7
IV. Donahue	20	198.4	7.0	36	27	191.6	37.6	10.4
V. Jacobus	20	258.1	70.7	28.2	27	221.3	14.7	8.6
VI. Schenker	21	229.4	83.0	0.5	28	250.1	78.8	20.2
VII. Stapleton			<b> </b>		29	217.1	80.5	8.1
Gro	up av	erages	• •	• • •		205:8		
VIII. Callahan	21	205.5	12.5	12.8				

ABLE 4. - MAT. 1904.

Name.	Date.	Avg.	M.v.	Var. from G. A.	Date.	Avg.	K, v,	Var. from G. A.	Date.	Avg.	М. ч.	Var. from G. A.	Dote.	Avg.	K	Var. from G. A.
I. Anderson, G. W.	:	:	:	:	18	181.2	16.5	8.7	ଛ	199.6	8.2	14.7	27	190.7	12.1	5.8
II. Anderson, W. L.		:	:	:	91	202.1	12.1	26.4	:	:	:	:	র	246.4	84.9	17.9
III. Bellis	:	:	:	:	22	186.6	19.4	8.7	18	185.8	8.4	4.0	75	171.8	10.3	10.6
IV. Donahue	:	:	:	:	:	:	:	:	18	217.2	18.0	16.2	28	207.1	19.2	5.1
V. Jacobus	4	240.6	42.4	15.7	11	234.2	40.0	9.3	18	235.5	39.6	10.6	25	246.4	29.7	20.5
VI. Schenker	9	214.3	24.0	15.6	:	:	:	:	19	234.3	40.5	4.4	98	196.9	16.9	33.0
VII. Stapleton	<u>:</u>	:	:	:	=	224.4	16.9	16.4	18	201.1	11.7	4.9	25	217.8	16.5	8.8
	ð	Group averages	erages								:			210.8		

TABLE 5. - JUNE, 1904.

Name.	Date.	Avg.	M. v.	Var. from G. A.
I. Anderson, G. W	3	197.4	16.8	12.5
II. Anderson, W. L	7	226.8	20.3	1.7
III. Bellis	7	162.0	8.4	19.8
IV. Donahue	1	208.4	31.7	6.4
V. Jacobus	1	284.6	45.6	59.7
VI. Schenker	8	228.6	48.3	1.3
VII. Stapleton	7	210.4	34.2	1.4
Group averages		216.9		

TABLE 6.

Name.	Avg. of all Tests for the Month.	Avg. of all Tests of for the Month.	Avg. of all Tests of for the Month.	Avg. of all Tests for for the Month.	Avg. of all Tests g for the Month.	General Avg. of all Testa.
I. Anderson, G. W	177.9	177.8	192.9	190.5	197.4	184.9
II. Anderson, W. L	277.3	210.7	195.0	224.3	226.8	228.5
III. Bellis	185.8	182.3	187.8	178.5	162.0	181.8
IV. Donahue	178.1	213.4	195.0	212.2	208.4	202.0
V. Jacobus	187.8	208.4	237.2	238.9	284.6	224.9
VI. Schenker	186.1	200.7	289.8	215.2	228.6	229.9
VII. Stapleton	185.0	208.3	217.1	214.4	210.4	209.0
Monthly averages	196.8	200.2	209.8	210.6	216.9	
VIII. Callahan	188.5	194.9	205.5			193.2

Ŀ
闰
Ħ
Ţ
_

	February.	Ė		March.	ند		Ψ	April.		7	May.		June.	
Name.	Mo. of Tape In 10 sec.			Mo. of Tape in 10 sec.			sucT to off	Mo. of Tape one of ni	·	sady to off	Mo. of Tape oee Of at		Mo. of Tape in 10 sec.	
I. Anderson, G. W.	<b>8</b>	78	:	49	92	72	78	27	:	78	n	72	76	75.0
II. Anderson, W. L	81	88	92	2	11	:	:	88	:	78	:	88	82	80.8
III. Bellis	\$	8	78	82	8	92	78	8	:	8	28	67	72	78.0
IV. Donahue	8	8	8	88	8	:	88	8	:	:	8	88	88	86.8
V. Jacobus	20	88	22	11	22	92	4	82	22	8	76	8	74	77.8
VI. Schenker	8	8	2	\$	2	:	8	8	49	:	8	2	8	65.6
VII. Stapleton	81	:	8	Ŀ	:	:	:	#	:	18	22	#	74	77.1
Group averages	78.8	:	:	74.4	:	:	:	76.8	:	:	:	78.7	76.4	
VIII. Callahan	88	75	88	88	18	88	88	:	:	:	:	:	:	88
		İ	l			l	l				l			

TABLE 8.

	February.	March.	April.	May.	June.
Name.	Avg. of all Determina- tions for the Month.	Avg. of all Determina- tions for the Month.	Avg. of all Determina- tions for the Month.	Avg. of all Determina- tions for the Month.	Avg. of all Determina- tions for the Month.
I. Anderson, G. W.	88	71	75	74	76
II. Anderson, W. L.	84	79	88	81	78
III. Bellis	87	78 <sup>.</sup>	69	74	72
IV. Donahue	85	83	85	90	88
V. Jacobus	76	74	78	80	74
VL Schenker	64	66	65	69	74
VII. Stapleton	87	70	79	78	74
Monthly averages .	81	74	76	81	75
VIII. Callahan	79	85	86		

j j	) Be	February.		March.	<i>ਬ੍</i> ਹ		•	April.		Ę.	May.			June.
	Date.		Date.		Date.		Date.		Date.		Date.		Date.	
I. Anderson, G. W. 19 34 (28)	19	34 (28)		11 27 (23)	88	20 (17)	8	20 (15)	ន	16 (13)	:	:	∞	مد
II. Anderson, W. L.	27	27 48 (87)	10	72 (69)	17	49 (41)	:	:	9	28 (19)	:	:	~	12 (9)
III. Bellis	16	67 (43)	80	48 (37)	83	80 (24)	8	40 (24)	18	23 (18)	:	:	~	16 (12)
IV. Donahue	11	17 45 (40)-	87	27 (21)	প্ত	86 (30)	27	24 (22)	32	18 (16)	:	:	_	0
V. Jacobus	11	87 (30)	6	76 (68)	16	86 (81)	ଛ	24 (20)		11 27 (18)	26	20 (15)	-	18 (14)
VI. Schenker	11	75 (66)	6	62 (88)	:	:	21	63 (54)	19	46 (89)	:	:	8	30 (24)
VII. Stapleton	19	44 (83)		11 82 (26)	· :	:	8	80 (22)	16	11 (10)	:	:	~	•
VIII. Callahan		18 29 (26)		17 24 (20)	:		<u>:</u>	:	:	• • • •		:		:

## GENERAL SUMMARY.

It is quite evident from a study of the results obtained in the foregoing experiments that young, vigorous men of the type under observation, trained in athletics, accustomed to the doing of vigorous muscular work, can satisfy all the true physiological needs of their bodies and maintain their physical strength and vigor, as well as their capacity for mental work, with an amount of proteid food equal to one-half, or one-third, that ordinarily consumed by men of this stamp. As the results show, all these men reduced their rate of proteid metabolism in such degree that the amount of nitrogen excreted daily during the period of the experiment averaged 8.8 grams, implying a metabolism of about 55 grams of proteid matter per day.

In other words, these athletes were able to reduce their nitrogenous metabolism to as low a level as many of the men of the professional group and of the soldier group, and this with not only maintenance of health and strength, but with a decided increase in their muscular power.

Metabolized nitrogen per kilo of body-weight for all these men, with one exception, during the experiment amounted to 0.108 to 0.134 gram per day, fully as low as was obtained with the members of the soldier detail on their prescribed diet. It is clear, therefore, that physiological economy in nutrition is as safe for men in athletics as for men not accustomed to vigorous exercise. There is obviously no physiological ground for the use of such quantity of proteid food, or of total nutrients, as the prevalent dietary standards call for.

The athlete, as well as the less active man (physically), or the professional man, can meet all his ordinary requirements with an intake of proteid food far below the quantities generally consumed, and this without increasing in any measure the amount of non-nitrogenous food.

# IV. THE SYSTEMIC VALUE OF PHYSIOLOGICAL ECONOMY IN NUTRITION.

It is one of the axioms of physiology that the majority of the diseases of mankind are due to, or are connected with, perversions of nutrition. General or local disturbances of metabolism are broadly responsible for disease, and with a due recognition of this fact it may be well to consider more specifically whether greater economy in the consumption of food, i. e., a restriction of the daily diet to amounts more commensurate with the physiological needs of the body, may not be of value in preventing disease, or prove of use in combating disease when the latter has manifested itself.

Broadly speaking, the extent and character of the metabolic processes of the body are dependent in large measure upon the amount and character of the diet. Further, it is equally certain that the chemical composition of the blood and lymph is quickly affected by the amount and character of the food materials absorbed from the alimentary canal. Even in the matter of secretion of the digestive juices, we have learned, through the recent experiments of Pawlow, that the chemical composition and solvent action of these fluids may be modified by the amount and character of the food fed. How much more, then, may we expect the intricate processes of cell and tissue metabolism to be modified by changes in the chemical composition of the blood and lymph that bathe them.

Further, recognizing as we must the extreme sensitiveness of the central and peripheral parts of the nervous system to changes in the composition of the blood, we see suggested indirect ways by which metabolism, both general and local, may be modified by influences exerted upon the nervous system, whereby the nutritive condition of individual structures may undergo change. Vasomotor influences, controlled as they are by nerve fibres, which in turn are sensitive to the conditions of their environment, likewise indirectly affect the rate and character of tissue metabolism; a fact which may serve to emphasize the many ways whereby the metabolism of an organ or tissue may be modified through the primary influence of a diet which, controlling in a measure the volume and character of the circulating blood and lymph, must of necessity exert an influence more or less extended.

The one factor above all others that tends to increase the extent of proteid katabolism is the amount of proteid food ingested. Increase in the amount of the albuminous foodstuffs is at once, or speedily, followed by an increase in the output of nitrogenous waste products, the latter constituting a good measure of the extent of proteid metabolism going on in the body. We have been taught to believe that the healthy adult under ordinary conditions of life needs for the maintenance of health, strength, bodily and mental vigor, about 118 grams of proteid food daily. This amount of albuminous food, if metabolized, means at least 16 grams of nitrogen in the urine, in the form of urea, uric acid, creatinin, purin bases, and other nitrogenous products more or less closely related. Under the stress of modern conditions and following the dictates of an acquired taste, the daily intake of proteid food in many individuals at least far exceeds the above figures, with an increase of proteid katabolism equal to 18 or more grams of nitrogen in the 24 hours' urine.

When we recall that these 18 grams, or more, of nitrogen in the urine reach the final stage of urea, etc., only by passing through a series of stages, each one of which means the using up of a certain amount of energy, to say nothing of the energy made use of in digestion, absorption, etc., we can easily picture to ourselves the amount of physiological labor which the daily handling by the body of such amounts of proteid food entails. Further, it needs very little imagination to see that a large amount of energy is used up in passing on these nitrogenous waste products from organ to organ, or from tissue to tissue, on the way to elimination, and we can fancy that liver and kidneys must at times rebel at the excessive labor they are called upon to perform.

Moreover, the thought suggests itself that possibly these waste products of proteid metabolism, the leucomaines so abundantly formed in the breaking down of proteid material, are not wholly free from objectionable features. If so, an excess of such products might be advantageously dispensed with. Indeed, we have what seems abundant evidence tending to show that many of the nitrogenous waste products elaborated in the body through the breaking down of proteid materials are possessed of more or less physiological action. that direct antecedent of urea, ammonium carbamate, which we have reason to believe is formed more or less generally throughout the tissues of the body, is not above suspicion. To be sure, Nature has provided a mechanism in the hepatic cells whereby it is quickly transformed into the harmless urea, but it is only necessary to join the portal vein to the hepatic vein, thereby throwing the liver out of the circuit, in order to see the effect of an excess of proteid food. Under such conditions, this is followed by the appearance of all the symptoms of poisoning with ammonium carbamate, i. e., convulsions ending in death.\*

Further, we may refer to the observations of Mallet† with creatin and creatinin, two conspicuous nitrogenous bases present in muscle, which show unmistakably that these bodies tend to retard slightly the action of the heart. This seems to be their most decided physiological effect, although large doses likewise cause a slight frontal headache, with some general nervous agitation. Attention may also be called to the extremely important experiments of Minkowski, ‡ in which he found that adenin — one of the purin bases formed in the breaking down of cell nuclei — has a most marked toxic action, both on man and on dogs. Adenin affects the circulatory ap-

<sup>\*</sup> See Hahn, Massen, Nencki, und Pawlow: Archiv f. exper. Pathol. u. Pharm. Band XXXII. (1898), p. 101. Also, Nencki, Pawlow, und Zaleski: Ibid. Band XXXVII., p. 26.

<sup>†</sup> The physiological effect of creatin and creatinin, etc. Bulletin No. 66. U. S. Department of Agriculture, Office of Experiment Stations.

<sup>†</sup> Untersuchungen zur Physiologie und Pathologie der Harnsäure bei Säugethieren. Archiv f. exper. Pathol. u. Pharm. Band XLI., p. 406.

paratus very strikingly, increasing the heart's action, etc.; it acts on the mucous membrane of the duodenum, causing an acute inflammation, thus leading to continuous vomiting, and in addition it has a local action on the kidneys, giving rise to a deposition in the kidney itself of spheroliths of uric acid, or urates, which leads to an acute nephritis with albuminuria, from which the animal speedily dies.

The alloxuric bases likewise cause fever when injected into the circulation or taken per os,\* and according to the recent observations of Mandel † there is a very noticeable relationship between the amount of alloxuric bases eliminated through the urine and the temperature of the body in cases of aseptic fevers, indicating that these substances, with possibly other incomplete products of tissue metabolism, are important factors in the production of febrile temperatures.

Reference may also be made to our general knowledge regarding the relationship between uric acid and gouty affections, including rheumatism, to say nothing of the possible relationship between uric acid and many other diseases less clearly established. The broader question deserving attention just here, however, is that all of the so-called leucomaines which, as Gautier states, are being formed continuously in the animal tissues side by side with the formation of urea and carbonic acid, and at the expense of the nitrogenous elements or proteid matter, are more or less toxic in their properties, at least under certain conditions of the body. is perfectly clear that there are a large number of leucomaines, or nitrogenous waste products, which are indissolubly connected with the metabolism of cell protoplasm, and the formation of these substances is augmented by a diet rich in proteid matter.

It is well understood that the excretions of all living organisms, both plant and animal, are more or less poisonous to the

<sup>\*</sup> See Burian and Schur., Archiv f. die] gesammte Physiologie. Band LXXXVII., p. 239.

<sup>†</sup> The alloxuric bases in aseptic fevers. Amer. Journal of Physiology. Vol. X., p. 452.

organisms which produce and excrete them. The substances so formed originate in the metabolic changes by which complex organic molecules are broken down into simpler compounds. As stated by Vaughan and Novy,\* "we have good reason for believing that the proteid molecule has certain lines of cleavage along which it breaks when certain forces are applied, and that the resulting fragments have also lines of cleavage along which they break under certain influences, and so on until the end-products, urea, ammonia, water, and carbon-dioxide, are reached; also that some of these intermediate products are highly poisonous has been abundantly demonstrated." It would therefore seem self-evident that the nitrogenous waste products of the body, i. e., the products of proteid katabolism, may be more or less dangerous to the welfare of the body, and consequently there would seem to be reason in the assumption that greater freedom from disease - especially from the so-called autogenous diseases - might be expected where greater care is exercised in the amount of proteid food consumed.

It is generally understood, or at least is frequently stated by medical writers, that certain febrile conditions are autogenous, and Brunton has made the assertion that the condition termed "biliousness," and which is most prone to occur in persons who eat largely of proteid foods, is due to the formation of poisonous alkaloidal-like substances which might well be classified under the broad term of leucomaines. To repeat, there are a great many observations and some facts which warrant the view that the nitrogenous waste products of the body — the products of proteid katabolism — are more or less dangerous to the well-being of the organism, and hence there seems justification for the belief that there is greater safety for health and longevity in adopting dietetic habits that are more in accord with the real needs of the body.

The writer's opinion upon this question has been greatly strengthened by the large numbers of letters he has received

<sup>\*</sup> Ptomaines and Leucomaines, or the Chemical Factors in the Causation of Disease. Third Edition, 1896. Lea Brothers. p. 550.

- during the course of this inquiry - from persons all over the world, many of whom in their search for health and strength have adopted more frugal methods of living, and who have found relief in an abstemiousness which, compared with ordinary dietetic standards, would seem quite inadequate to support life, yet they have recovered health and strength, and by the judicious practice of physiological economy in their diet have maintained health and vigor, with capability for work that has proved a perpetual surprise to themselves and their friends. The writer's faith in these spontaneous statements made by persons wholly unknown to him has been augmented by his personal knowledge of people suffering with various troubles, who have found relief by the simple use of reason and judgment in the taking of food, with a view to lowering the rate of proteid metabolism. There is no question in the mind of the writer that excessive proteid decomposition within the body entails possible danger.

If it is true, on the other hand, that the healthy organism needs a daily intake of 118 grams of proteid food more or less, in order to maintain physiological equilibrium, to keep up physical and mental vigor, and to preserve the normal power of resistance to the incursions of disease, then we must consider that the good overbalances the evil, and that evil exists in order that good may be accomplished. We are certainly justified, however, in saying, on the basis of our daily observations made on a large number of individuals and extending over many months, that there is no apparent need for any such amount of proteid food as is ordinarily consumed by the average individual.

We can point to various persons who, for periods ranging from six months to a year, have metabolized daily 5.5 to 7.5 grams of nitrogen instead of 16 to 18 grams, i. e., they have subsisted quite satisfactorily on an amount of proteid food daily, equal to one-third or one-half the amount ordinarily considered as necessary for the maintenance of health and strength, and this without unduly increasing the amount of non-nitrogenous food. Further, our observations have shown

that with this great reduction in the consumption of proteid food, with corresponding diminution of proteid katabolism, body-weight can be maintained at a stationary figure, after the body has once adjusted itself to the new conditions. Moreover, there is marked increase in physical strength as demonstrated by repeated dynamometer tests on many individuals, which may perhaps be ascribed to the greater freedom of blood and lymph, as well as of muscle-plasma, from nitrogenous extractives. Lastly, we have failed to find any falling off in physical or mental vigor, any change in the hæmoglobincontent of the blood, or in the number of erythrocytes. fact, all our observations agree in showing that it is quite possible to reduce with safety the extent of proteid katabolism to one-third or one-half that generally considered as essential to life and strength. In other words, there is perfect safety in a lowered proteid metabolism, and we are inclined to raise the question whether a daily diet containing one-half, or even less, the amount of proteid food ordinarily consumed does not come nearer to the normal and natural requirements of the healthy body than the more elaborate standards we have gradually adopted.

Here, then, we have suggested a radical change in diet which experiment shows is perfectly safe, and we are disposed to urge that there is great systemic value, both in health and in many forms of disease, in such a change. It is obvious, as previously stated, that the smallest amount of food that will serve to maintain bodily and mental vigor, keep up bodily strength, and preserve the normal powers of resistance to disease, is the ideal diet. Any excess over and above what is really needed for these purposes imposes just so much of an unnecessary strain upon the organism. It entails a wasteful expenditure of energy that might better be preserved for future emergencies. It imposes upon the excretory organs the needless labor of removing waste products which could well be dispensed with, to say nothing of the possible physiological action of these products as they circulate through the body.

Dr. Walker Hall,\* in his interesting article in "The Practitioner" on "Metabolism in Gout," states that "under normal circumstances a man weighing eleven stone and performing average work requires twenty grams of nitrogen and three hundred grams of carbon per day." This statement is in perfect harmony with generally accepted views, but I should like to emphasize the fact that all of the twenty-six men we have been experimenting with at New Haven, representing different types, ages, and degrees of activity, have been able to maintain health, strength, and vigor, from six months to a year on a daily quantity of nitrogen equal to one-half, one-third, and even one-quarter the amount of this so-called necessary twenty grams. Further, nitrogenous equilibrium was easily maintained on such quantities of proteid food, and, as before stated, there was great gain in physical strength. Are we not justified, therefore, in raising the broad question whether such a radical change in diet as these facts suggest might not be of systemic value in gout, and especially in cases where there is a predisposition to gout. Speaking as a physiologist, the writer is strongly of the opinion, based in part upon his own observations and in part upon both the voluntary and unconscious testimony of others, that there is possible great gain to the gouty and rheumatic individual by a practice of physiological economy in nutrition.

Physiological economy, as the writer defines it, is not prohibition, but temperance. Moderation in diet, especially in the taking of proteid foods, means a great saving in the wear and tear of the body machinery. It must presumably mean greater freedom from many diseases in which individual organs, such as the liver and kidneys, are frequently involved. It suggests, likewise, greater freedom from many disturbances of general metabolism which eventually terminate in a perversion of nutrition, so marked as to constitute a serious condition of disease. More specifically, lowered proteid metabolism means diminished introduction and diminished

<sup>\*</sup> The Practitioner. London. July, 1903. p. 61.

formation of nitrogenous products of the purin type, such as xanthin, hypoxanthin, guanin, adenin, etc., as well as of other nitrogenous bodies less clearly defined. Consequently, we have as one of the results of such a systemic change in diet a decreased formation of uric acid, or at least a diminished output of uric acid through the urine.

Obviously, a lowered proteid intake means, in some measure at least, a decreased consumption of meat and similar products more or less rich in free and combined purin bases. quite plainly must result in a diminished production of uric acid, but the writer is strongly of the opinion that we do not as yet possess sufficiently full knowledge regarding all the ways in which uric acid results in the body. It is true, we differentiate between endogenous and exogenous uric acid, and further, we understand quite clearly that variations in the intake of free and combined purin bases exercise a potent influence upon the output of uric acid through the urine. We still lack, however, concise information as to the various ways in which uric acid may be produced, and its ultimate fate in the body. This is well illustrated by a recent paper from the Marburg laboratory, in which Kutscher and Seemann\* point out the possibility of a production of uric acid in the animal body synthetically, and likewise suggest that uric acid may be utilized for the formation of nuclein bases, i.e., a reversal of the oxidative process by which uric acid results from the ingestion of free or combined nuclein bases, suggesting indeed the possibility of uric acid and the nuclein bases being produced from each other, according to the circum-Thus, when nucleins or free purin bases are taken with the food, the organism may utilize this material at once in the synthesis of nucleins for the use of the body cells. There is no need of a reduction of the formed uric acid to nuclein bases, and consequently there is an increased excretion of uric acid through the urine, but this does not result from a direct transformation of the ingested purin material into uric

<sup>\*</sup> Centralblatt für Physiologie. Band XVII, p. 715. 1904.

acid, but is the result of a sparing of the already formed uric acid. The nuclein bases thus act as sparers of uric acid. This view explains, according to Kutscher and Seemann, why feeding with nuclein bases increases the output of uric acid, and feeding with uric acid—a sudden overflow of uric acid into the circulation—is followed, as a rule, by an increased elimination of urea, the uric acid being thus transformed by energetic oxidation. This hypothesis is brought forward not merely because it is an interesting suggestion, but mainly because it illustrates that we do not as yet know fully all the steps in the production of uric acid, nor do we know how far the uric acid we find and determine in the urine is a measure of the formation of uric acid in the body.

Taking our knowledge on these matters as it stands to-day, however, we find by experiment that lowering the intake of proteid food, with its consequent corresponding diminution in proteid katabolism, is followed at once by a marked decrease in the output of uric acid. Let us consider a few of the data obtained in our experiments. The first case I will refer to is that of a college athlete (Callahan). For a period of ten days on his ordinary diet, the average amount of nitrogen in the urine per day was 22.8 grams, equal to the metabolism of 142.5 grams of proteid food. During this same period the average daily output of uric acid was 1.103 grams. For the following four months and a half, on a more restricted diet, with a marked cutting down of the proteid food, but with no exclusion of meat, the average daily output of nitrogen through the urine was 9.04 grams. In other words, for this period of over four months the extent of proteid katabolism was reduced considerably more than 50 per cent. The average daily output of uric acid for this same period was 0.624 gram, equal to a reduction of about 40 per cent from his normal excretion.

Another college athlete (Stapleton), on his normal diet, showed an average excretion of nitrogen through the urine per day, for ten days, of 19.70 grams, while the daily average

excretion of uric acid for the same period was 0.893 gram. On a more restricted diet, with diminished proteid katabolism, the daily average excretion of nitrogen through the urine for a period of over four months was 11.06 grams, while the daily average excretion of uric acid for the same period fell to 0.699 gram. In the first of these two cases the average daily ratio of uric acid to total nitrogen during the period of lowered proteid metabolism was 1:14. In the second case the ratio was 1:16.

A third college student (G. W. Anderson), on his ordinary diet, excreted through the urine for a period of nine days 17.17 grams of nitrogen as the daily average, while the average daily output of uric acid for the same period was 0.956 gram. On the more restricted diet of the next four or five months his average daily excretion of nitrogen fell to 9.37 grams per day, while the average daily excretion of uric acid was reduced to 0.632 gram. On his ordinary diet, the ratio of uric acid to nitrogen was 1:18, while later with the diminished proteid metabolism the ratio was 1:14.

Turning to another class, viz., professional men, reference may be made to the writer, whose average daily nitrogen excretion through the urine for a period of nearly nine months was 5.699 grams, corresponding to the metabolism of 85.6 grams of proteid per day. During this same period of nearly nine months the average daily excretion of uric acid amounted to 0.392 gram, the ratio of uric acid to total nitrogen being 1:14. In passing, it may be repeated that the subject of this experiment succeeded in maintaining a constant body-weight, and he further avers that in physical and mental vigor he can find no evidence of deterioration, although the amount of proteid food consumed daily during this long period was less than 40 grams per day. Further, he was in nitrogenous equilibrium during this period, although the nitrogen metabolized daily amounted to only 99 milligrams per kilo of body-weight. Another case in this same group may be mentioned, principally because the subject for over a year became a vegetarian, abstaining from all meat. During the last nine months, this man (Beers) eliminated 8.28 grams of nitrogen through the urine as the daily average, indicating a metabolism of 51 grams of proteid material per day. During this same period, the average daily excretion of uric acid was 0.349 gram, the ratio of uric acid to total nitrogen being 1:23.

The main point to be emphasized in these results is that they show quite conclusively how greatly the daily output of uric acid may be reduced by diminishing the intake of proteid food, and thereby restricting the extent of the proteid metabolism. The ratio of uric acid to the total nitrogen excreted may or may not be altered; this will depend in large measure upon the character of the diet, the relative proportion of free and combined purin bases introduced with the food, etc. As already stated, we do not know with certainty how far the excreted uric acid represents the formation of uric acid in the body, but presumably there is a more or less close relationship, and hence we are doubtless warranted in saying that the formation of uric acid is diminished, in essentially the same proportion as its excretion is reduced, with a lowered proteid intake. Certain it is that several of the persons under observation, who had troubles of a gouty and rheumatic nature in the past, have during the course of the experiment experienced relief, with complete and permanent abeyance of all symptoms. The writer is firmly of the opinion that ordinary gout and rheumatism are entirely preventable by reasonable care and judgment in the matter of diet. Whether, when once firmly established, in aggravated form, they will prove amenable to dietetic treatment is not so certain, but undoubtedly mild cases will respond to the beneficial influences of a rational diet, reinforced by treatment adapted to the removal of urates already deposited. In any event, due regard for the well known deleterious effects of purin-containing foods as a source of exogenous uric acid, and with restriction of proteid metabolism to the true necessities of the body, should serve as an effective means of preventing all those troubles for which uric acid is generally held responsible.

The two following tables give a summary of results bearing

upon the excretion of uric acid and its relation to nitrogen and body-weight, for all the subjects belonging to the "professional group" and the "student group." Emphasis should be laid upon the fact that these figures represent the average daily excretion for the different individuals through the entire period of the experiment.

AVERAGE DAILY EXCRETION THROUGH THE URINE FOR SEVEN-NINE MONTHS. — PROFESSIONAL GROUP.

Name.	Body- weight.	Total Nitrogen.	Uric Acid.	Ratio of Uric Acid to Mitrogen.	Uric Acid per kilo of Body- weight.	Phosphoric Acid P <sub>2</sub> O <sub>8</sub> .
	kilos	grams	gram		grams	grams
Chittenden	57.0	5.69	0.892	1:14	0.0068	0.90
Mendel	70.0	6.58	0.419	1 : 15	0.0060	1.46
Underhill	65.0	7.43	0.516	1:14	0.0079	1.28
Dean	65.0	8.99	0.386	1:28	0.0059	1.73
Beers	61.5	8.58	0.865	1:23	0.0059	1.49

AVERAGE DAILY EXCRETION THROUGH THE URINE FOR FOUR-FIVE MONTHS. — STUDENT GROUP.

Name.,	Body- weight.	Total Nitrogen.	Uric Acid.	Ratio of Uric Acid to Nitrogen.	Uric Acid per kilo of Body- weight.	Phosphoric Acid P <sub>2</sub> O <sub>2</sub> .
	kilos	grams	gram		grame	grams
Anderson, G. W.	71.0	9.37	0.682	1:14	0.0089	1.75
Anderson, W. L.	61.0	10.41	0.516	1:20	0.0084	2.14
Bellis	78.0	8.88	0.531	1 : 16	0.0068	1.98
Callahan	83.0	9.04	0.624	1:14	0.0075	1.74
Donahue	62.0	7.47	0.395	1 : 19	0.0068	1.79
Jacobus	56.0	7.58	0.423	1:17	0.0075	1.67
Schenker	73.0	10.09	0.624	1:16	0.0085	2.20
Stapleton	75.0	11.06	0.699	1 : 16	0.0093	2.64

#### 468 PHYSIOLOGICAL ECONOMY IN NUTRITION

Turning now to the third group of men, i. e., the soldier detail, under observation for a period of six months, during five months of which time they lived on a prescribed diet with diminished content of proteid food, but with no exclusion of animal food, the following average results are to be noted:

AVERAGE DAILY EXCRETION THROUGH THE URINE FOR FIVE MONTHS. — SOLDIER DETAIL.

Name.	Body- weight.	Total Nitrogen.	Uric Acid.	Ratio of Uric Acid to Nitrogen.	Uric Acid per kilo of Body- weight.	Phosphoric Acid P <sub>2</sub> O <sub>5</sub> .
	kilos	grams	gram		gram	grams
Oakman	62	7.42	0.405	1:18	0.0065	1,39
Morris	59	7.03	0.450	1 : 15	0.0076	1.25
Broyles	60	7.26	0.398	1 : 18	0.0066	1.41
Coffman	58	8.17	0.379	1 : 21	0.0065	1.23
Sliney	60	8.39	0.647	1 : 18	0.0107	1.82
Steltz	58	7.13	0.416	1:17	0.0078	1.24
Henderson	71	8.91	0.488	1 : 18	0.0068	1.42
Fritz	72	7.84	0.642	1 : 12	0.0089	1.58
Cohn	62	8.05	0.512	1 : 15	0.0082	1.28
Loewenthal	59	7.88	0.872	1:19	0.0068	1.28
Zooman	55	8.25	0.457	1 : 18	0.0083	1.19
Bates	65	8.08	0.387	1:20	0.0059	1.23
Davis	57	8.61	0.414	1:20	0.0072	1.42

These figures are interesting in many ways. First, they make clear that on the diet prescribed, these men were manufacturing or excreting about the same amount of uric acid per kilo of body-weight as the men of the two preceding groups, living more or less with free choice of food. In other words, all these men, with one and possibly two exceptions, were practically throwing out only uric acid of endogenous origin, i. c., that which came from the breaking down of the man's

own tissue cells. Second, it is to be noted that the ratio of uric acid to nitrogen in the men of this group varies only within narrow limits.

It is very evident from these figures, reinforced by those of the previous groups, that we can diminish greatly the output of uric acid by simply restricting the extent of proteid katabolism, through reduction in the amount of proteid food. Further, we now know that this general lowering of proteid metabolism can be accomplished not only without danger to the body, but with a distinct betterment of the physical condition.

Just here I should like to emphasize one point that appears to me of primary importance in any consideration of the influence of diet in gouty affections, and in so doing I merely echo a statement made by Sir Dyce Duckworth\*, viz., "that the subject of gout, either by inheritance or acquirement, is so far peculiar in his constitution that he reacts differently to various agencies, such as climate, food, etc., from persons not so disposed." In this connection, let me refer again to the foregoing table of results obtained with the soldier detachment, remembering that these thirteen men were living under exactly the same conditions and consuming the same kind of food each day, and in essentially the same amounts. Yet notice the striking variation in the output of uric acid by one of these men (Sliney), - a variation which shows itself especially when the uric acid is calculated per kilo of bodyweight. How can this variation be accounted for except on the assumption that there may be personal idiosyncrasies, personal coefficients of nutrition, natural or acquired, that modify to some extent the production of uric acid, the oxidation of uric acid, or the elimination of uric acid from the body?

Lastly, in advocating the possible systemic value of a lowered proteid metabolism as of value in the prevention of gout, and of other disorders which have their origin in per-

<sup>\*</sup> The Practitioner, July, 1903, p. 88.

verted nutrition, I am inclined to emphasize the desirability of using common-sense in the application of dietetic rules, remembering that man is an omnivorous animal, and that Nature evidently never intended him to subsist solely on a "cereal diet," or on any specific form of food to the exclusion of all others. On matters of diet every man should be a law unto himself, using judgment and knowledge to the best of his ability, reinforced by his own personal experiences. Vegetarianism may have its virtues, as too great indulgence in flesh foods may have its serious side, but there would seem to be no sound physiological reason for the complete exclusion of any one class of food stuffs, under ordinary conditions of life. Far more rational is temperance in place of prohibition, and I am inclined to emphasize the systemic value of a daily diet so reduced in quantity that the metabolic processes may be largely decreased, in closer harmony with true physiological needs, especially those which involve the breaking down of proteid matter; and in making this suggestion I can add the assurance, based upon these observations on many individuals, that there is not only perfect safety but gain to the body, in diminishing proteid metabolism to a level somewhere near the actual requirements of the individual.

# V. ECONOMIC AND SOCIOLOGICAL IMPORTANCE OF THE RESULTS.

The importance of the foregoing results from an economic and sociological standpoint is perhaps worthy of a brief consideration. We have learned that a much smaller amount of albuminous or proteid food than is ordinarily consumed will suffice for the daily needs of the body. It remains to be seen whether this fact will gain the popular recognition it would seem to deserve. Ignoring for the time the matter of physiological economy and its possible bearing upon health and strength, it is a fair question to ask why should people indulge in such wasteful extravagance in the matter of diet when there is no real physiological need for it? Why not accustom the body to a smaller consumption of food, thereby saving for other purposes the expenditure which this excess of food involves?

The question of the daily diet is one of the most important for the family of small means, and there is no reason why the family treasury should be so heavily drained for this imaginary need. Simplicity of living might well be given more careful consideration, and now that we have convincing proof of much smaller dietetic requirements on the part of the body, it might be well to consider the practical application these results naturally suggest. It is obvious from our data, that it is quite safe to diminish by one-half the amount of albuminous or proteid food ordinarily consumed, and this without any apparent detriment to health, and with even gain to the economy. The ordinary forms of proteid food are, as a rule, the most costly of dietetic articles, and since this restriction of albuminous food calls for no great increase in the amount of nonnitrogenous food, it is quite apparent that a great saving in the daily expenditure can be accomplished.

Obviously, however, there must be a decided change in the attitude of the public on this question before any great im-

provement can be hoped for. Habit and sentiment play such a part in our lives that it is too much to expect any sudden change of custom. By a proper system of education commenced early in life it may, however, be possible to establish new standards, which in the may prevail and eventually lead to more enlightened methods of living, whereby there will be less drain upon the resources of the people. With habits firmly fixed and palates calling for new sensations, reinforced by the prevalent opinion that by hearty eating lies the road to health and strength, it is easy to foresee difficulty in the advance of new doctrines along the lines indicated. The pleasure of eating is not to be minimized. The palate serves as the gateway through which food passes, and its sensitiveness and power of appreciation are not to be despised.

Simplicity of diet, however, does not diminish but rather increases the pleasure of eating, especially when daily restriction in diet — indulged in until a new habit is formed — has created a greater keenness of appetite, since under such conditions the palate takes on a new sensitiveness, and manifests a fuller appreciation of the variations of even a simple dietary. There is therefore no hardship, nor curtailment of the pleasure of eating in the restriction of the diet to the real needs of the body. Neither is there implied any cessation of that kindly hospitality that delights in the 'breaking of bread' with one's friends. With enlightened methods of living, on the other hand, will come a truer appreciation of the dignity of the body, and a lessened desire to manifest one's feelings of hospitality by a lavish intemperance that is as unphysiological as it is wasteful.

For the rich, as well as for the poor, there is need for careful consideration of this question of intemperance in the daily dietary. Were this the proper place, it would be easy to adduce figures showing the great waste which the consumption of food beyond the physiological requirements of the body entails. It needs no great imagination to picture the enormous saving per capita, in dollars and cents, by a reduction of the daily food to a true physiological basis.

The saving to the community, to the family, might well amount to enough to constitute the difference between pauperism and affluence. The resources of a community, as well as the resources of the family, are not to be lightly thrown away. We count the cost of this or that necessity, of this or that luxury, with careful consideration of the relative need and expense, but in the matter of living we pay little heed except it may be to exclude certain dietetic luxuries which seem beyond our purse. We are prone to fancy that health and strength are fostered by great liberality in the amount and variety of the daily food provided, and we are apt to express great concern if all the family and our guests do not avail themselves to the utmost of the foods so lavishly spread before them. The poorer man emulates his richer neighbors as soon as his circumstances will permit, and resources that could be much more advantageously expended for the good of the family and the home life are practically wasted - to say nothing of possible injury to health — under the mistaken idea that this more generous method of living is the surest road to health and strength.

Further, there is ground for thought in the possible economy of time which an improved condition of health would result in for the working members of the family. If greater economy in diet will diminish the number of sick days in the year, thereby increasing the working power of the wage earner, and if greater strength and efficiency can be acquired at the same time, the economic value of the proposition is at once apparent.

Finally, happiness and contentment, which usually appear in direct proportion to the health and prosperity of the individual, may be counted upon as becoming more conspicuous in the life of the community. So we see suggested various ways in which the application of the principles herein laid down, if consistently adopted and followed, may lead to a betterment of economic and sociological conditions. The writer, however, leaves to others, more familiar with sociological problems, the fuller development of this line of thought.

#### VI. GENERAL CONCLUSIONS.

When this investigation, the results of which have been detailed in the foregoing pages was first planned, it was intended to be simply a physiological study of the minimal proteid requirement of the healthy man, extended over sufficient time to render the results of scientific and practical value. There were no special theories involved, no special system of dietetics in view, but the object was simply to ascertain experimentally the minimum amount of proteid or albuminous food necessary for the maintenance of health and strength, under ordinary conditions of life. The impression in the mind of the writer was that there was no satisfactory scientific evidence to support the views held by most, if not all, physiologists regarding the needs of the body for food, especially nitrogenous or proteid food, and that the dietary standards universally adopted by scientific men were of very questionable accuracy, being founded mainly upon the customs and habits of mankind rather than upon any systematic study of what the actual necessities of the body are.

The results attained have certainly thrown a great deal of light upon this question of minimal proteid requirement, and the experimental study has been throughout a purely physiological one; but as the work has progressed the writer has been more and more impressed with the importance and significance of the results in their bearing upon the broader problem of general physiological economy in nutrition. There is no question, in view of our results, that people ordinarily consume much more food than there is any real physiological necessity for, and it is more than probable that this excess of food is in the long run detrimental to health, weakening rather than strengthening the body, and defeating the very objects aimed at.

Confining our conclusions to general statements, it may be said that our results, obtained with a great diversity of sub-

jects, justify the conviction that the minimal proteid requirement of the healthy man under ordinary conditions of life is far below the generally accepted dietary standards, and far below the amounts called for by the acquired taste of the generality of mankind. Expressed in different language, the amount of proteid or albuminous food needed daily for the actual physiological wants of the body is not more than one-half that ordinarily consumed by the average man. Bodyweight (when once adjusted to the new level), health, strength, mental and physical vigor, and endurance can be maintained with at least one-half of the proteid food ordinarily consumed; a kind of physiological economy which, if once entered upon intelligently, entails no hardship, but brings with it an actual betterment of the physical condition of the body. It holds out the promise of greater physical strength, increased endurance, greater freedom from fatigue, and a condition of wellbeing that is full of suggestion for the betterment of health.

Physiological economy in nutrition means temperance, and not prohibition. It means full freedom of choice in the selection of food. It is not cereal diet nor vegetarianism, but it is the judicious application of scientific truth to the art of living, in which man is called upon to apply to himself that same care and judgment in the protection of his bodily machinery that he applies to the mechanical products of his skill and creative power.

Food requirements must of necessity vary with changing conditions, but with due recognition of this fundamental principle, all the results so far obtained in this investigation, with a great variety of persons, point to the conclusion that the real demands of the body for proteid food do not exceed fifty per cent of the amount generally consumed. One-half of the 118 grams of proteid food called for daily by the ordinary dietary standards is quite sufficient to meet all the real physiological needs of the body, certainly under ordinary conditions of life; and with most individuals, especially persons not leading an active out-of-door life, even smaller amounts will suf-

fice. Excess means waste, but of far greater importance is the unnecessary strain placed upon the body by this uncalledfor excess of food material, which must be gotten rid of at the expense of energy that might better be conserved for more useful purposes.

Further, the total consumption of food by the average individual, non-nitrogenous as well as nitrogenous, is considerably greater than the real needs of the body demand, although here we must give closer heed to the varying requirements of the body incidental to varying degrees of activity. The man whose work is mainly mental has no real need for high fuel values in his daily ration. For such a man, a high potential energy in the daily intake of food is an incubus and not a gain. Body equilibrium can be maintained on far less than 3000 calories per day by the brain worker, and in the interest of health, strength, and vigor, as well as scientific truth, why teach the doctrine that a healthy man needs, on an average, foodstuffs to furnish 3000 calories or more per day, with 16 to 18 grams of nitrogen in the form of proteid? Moreover, as our experiments have clearly indicated, even the man who is called upon to perform considerable physical work has no apparent need for a fuel value in his food of 3000 calories per day. No doubt, the man who works at hard labor for ten or twelve hours a day will require a larger intake of fats and carbo-hydrates, sufficient to yield even more than 3000 calories, but this is not true of the moderate worker, nor of the average man whose work is in large measure mental rather than physical.

Finally, the writer may be permitted to express the hope that the outcome of this experimental work will serve to arouse scientific and intelligent interest in a subject which promises fruitful results for the individual, and for the community.

#### VII. DESCRIPTION OF ILLUSTRATIONS

Photographs of the soldiers were taken a few days prior to the close of the experiment, just before the men left New Haven at the termination of their work. Consequently, the pictures show the physical condition of the men after their long period of low nitrogen diet. Study of these photographs, especially those of the individuals, gives a correct idea of the appearance of the men, and shows the character of their muscular development at the close of their experimental work.

In considering these photographs, it must be remembered that the men as a class, as stated by Dr. Anderson in his Report, were not particularly well set up. It is evident, however, that the subjects were in good physical condition and had not lost any undue amount of flesh or fat. The two photographs of Fritz, facing pages 198 and 203, show him to have been in fine physical condition, with even a superabundance of fat. Steltz, on the other hand, whose photograph is shown facing page 211, was somewhat fine. This man, however, is of quite different build from his companion, Coffman, and was in excellent physical condition for certain lines of gymnastic work.

It may be well at this point to refer the reader to the photographs of W. L. Anderson and Bellis, facing pages 440 and 442. These men, typical Yale athletes, were in prime physical condition, and the photographs were taken prior to the experiment, at a time when they were consuming their ordinary, rich proteid diet. It is plain, by a comparison of these photographs, that Steltz was not trained to a much finer point than W. L. Anderson, although he does lack the full muscular development characteristic of the Yale athlete.


Sliney, whose photograph is found facing page 272, was likewise in a somewhat fine condition. He, however, like Steltz, was in splendid physical shape, so far as can be judged by his general health, spirits and aptitude for work. The

other men of the soldier group, whose photographs are shown, were not trained down to quite the same degree. Both Sliney and Steltz, however, had essentially the same body-weight at the close of the experiment, as on their arrival in New Haven. Steltz, indeed, weighed a trifle more in April, 1904, than he did in October, 1903. Sliney, on the other hand, had lost about one pound in weight. It is obvious, therefore, that these two men do not owe their spare condition to the low proteid diet.

The photographs facing pages 136, 261, 284 and 296 illustrate some of the methods employed in attempts to improve the bodily movements of the soldiers.

Among the group of University athletes, the photographs of Stapleton, facing pages 328 and 366, show the muscular development of a typical athlete endowed with more than the usual amount of muscular tissue. These two photographs of Stapleton were taken in April, after the subject had been for several months on a low proteid diet. There is in the photographs certainly no suggestion of any loss of muscle tissue, and no evidence of physical weakness. Stapleton, as has been previously stated, was an expert in wrestling and events of that character, for which his heavy muscular build well fitted him.

The photographs of W. L. Anderson and Bellis, facing pages 440 and 442, show, on the other hand, two athletes whose characteristic build is indicative of ability as gymnasts. More graceful in form, with smaller joints, and less heavy musculature, these men, at the time the photographs were taken, were in the pink of condition, and in a high degree of training for their special fields of athletic work. Emphasis should be laid upon the fact that at the time these two photographs were taken, the men in question had not commenced to lower their daily amount of proteid food. These two photographs are introduced especially to illustrate the general physical makeup of the men belonging to the group of University athletes made use of in the experiments.



# DATE DUE

		1	
1-7-1	74 CL	A	
	1 105	VE	

DEMCO 38-297





### DIGEST OF THE

## LIBRARY REGULATIONS.

No book shall be taken from the Library without the record of the Libraryan.

No person shall be allowed to retain more than five volumes at any one time, unless by special vote of the Council.

Books may be kept out one calendar month; no longer without renewal, and renewal may not be granted more than twice.

A fine of five cents per day accurred for every volume not returned within the time specified by the rules.

The Librarian may demand the return of a book after the expiration of ten days from the date of borrowing.

Certain books, so designated, cannot be taken from the Library without special permission.

All books must be returned at least two weeks previous to the Annual Meeting.

Persons are responsible for all injury or loss of books charged to their name.



5.M.1905.3 Physiological economy in nutrit1907 Countway Library BDR6445 3 2044 045 475 134